ARCTIC PERNAAFROST ATLAS

ARCTIC PERMAFROST ATLAS

Disclaimer

The views and interpretations in this publication are those of the author(s). They are not attributable to GRID-Arendal nor any of the institutions to which the contributors are affiliated with, and do not imply the expression of any opinion concerning the legal status of any country, territory, city or area of authority, or concerning the delimitation of frontiers or boundaries, or the endorsement of any product.

Copyright © 2023 GRID-Arendal

ISBN: 978-82-7701-217-9 DOI: https://doi.org/10.61523/KPJI4549

GRID-Arendal would appreciate receiving a copy of or the link to any material that uses this publication or its graphics as a source. None of the photos (Katie Orlinsky) or artwork (Oluko) may be reproduced. No part of this publication may be reproduced for resale or any commercial purpose without prior permission in writing from GRID-Arendal. Applications for such permission, with a statement of the purpose and extent of the reproduction, should be addressed jointly to grid@grida.no and nunataryuk@awi.de.

This publication and its maps are available in electronic form at www.grida.no/activities/404, www.awi.de and www.nunataryuk.org/news/atlas.

Recommended citation:

Westerveld, L., Kurvits, T., Schoolmeester, T., Mulelid, O. B., Eckhoff, T. S., Overduin, P. P., Fritz, M., Lantuit, H., Alfthan, B., Sinisalo, A., Miesner, F., Viitanen, L.-K., and the NUNATARYUK consortium (2023). *Arctic Permafrost Atlas*. GRID-Arendal, Arendal.



The Arctic Permafrost Atlas was written and edited by GRID-Arendal together with all NUNATARYUK project partners, scientists, and early-career scientists



This project is funded by the European Union's Horizon 2020 research and innovation programme under grant agreement No. 773421.

ARCTIC PERMAFROST ATLAS

Contents

Foreword	6
Prologue	7
Earth's Freezer: Introduction to Permafrost	9
Frozen grounds: Permafrost in the Arctic	11
Permafrost in profile: Landscape features	12
Frozen in time: The history of permafrost	15
An icy balance: Arctic permafrost physiography	17
What lies within: Organic carbon in permafrost	19
When ice grows up: Pingo Canadian Landmark	21
Drilling down: Learning the secrets of permafrost	22
Portrait: Annett Bartsch	25
Un/settled: Life on frozen ground	26
Frozen States I: Russian Federation	29
Portrait: Vyacheslav Shadrin	31
Frozen States II: North America	32
Portrait: Jessi Pascal	35
Frozen States III: Nordic region	36
Portrait: Palle Jerimiassen	39
Awakening Giant: Permafrost and Climate Change	41
Warming up, warming down: Increasing ground temperatures	42
The chill is gone: Thickening of the active layer	44
Disappearing act: Declining permafrost extent	47
Microorganisms, macro effects: Permafrost carbon cycle	49
Faster, deeper, stronger I: Speed of thaw in North America	50
Faster, deeper, stronger II: Speed of thaw in Scandinavia and the Russian Federation	53
Crossing the threshold: Future scenarios of carbon release	54
Portrait: Dmitry Streletskiy	57

Moving Grounds: Permafrost Changes	59
Frost and flora: The role of vegetation in permafrost landscapes	61
Fire on ice: Peat, permafrost, and fire	63
State of matter: Water, snow, and permafrost	64
The rivers run through it: Arctic rivers, deltas and hydrology	67
Along the edge of the world: Arctic coastal classification	68
Wear and tear: Erosion of Arctic permafrost coasts	70
Eating into the landscape: Retrogressive thaw slumps	73
Portrait: Angus Alunik	75
Losing ground: Projected rates of Arctic coastal erosion	76
Beneath the waves: Changes in subsea permafrost	79
	~ 4
Arctic Ripples: Impacts of Permafrost Thaw	81
Arctic Ripples: Impacts of Permafrost Thaw Feeling the heat: Permafrost thaw impacts on infrastructure	81 82
	_
Feeling the heat: Permafrost thaw impacts on infrastructure	82
Feeling the heat: Permafrost thaw impacts on infrastructure Risky business I: North American Arctic and Kalaallit Nunaat (Greenland)	82 84
Feeling the heat: Permafrost thaw impacts on infrastructure Risky business I: North American Arctic and Kalaallit Nunaat (Greenland) Risky business II: The Russian Federation and Scandinavian Arctic	82 84 87
Feeling the heat: Permafrost thaw impacts on infrastructure Risky business I: North American Arctic and Kalaallit Nunaat (Greenland) Risky business II: The Russian Federation and Scandinavian Arctic Terra infirma I: Coastal infrastructure in Yamalo-Nenets	82 84 87 89
Feeling the heat: Permafrost thaw impacts on infrastructure Risky business I: North American Arctic and Kalaallit Nunaat (Greenland) Risky business II: The Russian Federation and Scandinavian Arctic Terra infirma I: Coastal infrastructure in Yamalo-Nenets Portrait: Susanna Gartler	82 84 87 89 91
Feeling the heat: Permafrost thaw impacts on infrastructure Risky business I: North American Arctic and Kalaallit Nunaat (Greenland) Risky business II: The Russian Federation and Scandinavian Arctic Terra infirma I: Coastal infrastructure in Yamalo-Nenets Portrait: Susanna Gartler Terra infirma II: Reinforcing runways in Paulatuk	82 84 87 89 91 92
Feeling the heat: Permafrost thaw impacts on infrastructure Risky business I: North American Arctic and Kalaallit Nunaat (Greenland) Risky business II: The Russian Federation and Scandinavian Arctic Terra infirma I: Coastal infrastructure in Yamalo-Nenets Portrait: Susanna Gartler Terra infirma II: Reinforcing runways in Paulatuk Terra infirma III: Keeping cold food cold in Alaska	82 84 87 89 91 92 94
Feeling the heat: Permafrost thaw impacts on infrastructure Risky business I: North American Arctic and Kalaallit Nunaat (Greenland) Risky business II: The Russian Federation and Scandinavian Arctic Terra infirma I: Coastal infrastructure in Yamalo-Nenets Portrait: Susanna Gartler Terra infirma II: Reinforcing runways in Paulatuk Terra infirma III: Keeping cold food cold in Alaska Terra infirma IV: Urban planning in Ilulissat	82 84 87 89 91 92 94 97

104

107

109

110

.

_

_

Coming back to life: Reemerging pathogens Frozen assets I: The formal economy

Cultural homeland: Alaas landscapes in Yakutia

Frozen assets II: Traditional and subsistence activities

-

- -

Holding Tight: Adaptation to Permafrost Thaw	113
Bumpy road ahead: Transportation infrastructure and permafrost	114
Undermined: Mining infrastructure and permafrost	117
Keeping the light on: Energy infrastructure and permafrost	118
No time to waste: Waste management and permafrost	120
Modern history: Preserving Svalbard's cultural heritage	123
Portrait: Ingrid Rekkavik	125
Going South: Permafrost in Other Areas	127
A planetary perspective: Permafrost outside the Arctic	129
Frozen giants: Permafrost in the mountains	130
The view from the top: The Qinghai-Tibetan Plateau, Hindu Kush Himalaya, and Andes	132
Europe's frozen heart: Permafrost in the Alps	134
The ends of the Earth I: Permafrost in Antarctica	136
The ends of the Earth II: Antarctic Peninsula	138
The ends of the Earth III: Queen Maud Land, Victoria Land, and the McMurdo Dry Valleys	140
Over the Horizon	143
Glossary	146
Acronyms	147
References	148
Authors and acknowledgments	166
Artist spotlight: Olga Borjon-Privé (Oluko)	168
Artist spotlight: Katie Orlinsky	170

Foreword

The word "atlas" was coined by the renowned Flemish geographer and cartographer, Gerardus Mercator. In 1595, he published a series of maps intertwined with text and epitaphs, much like a modern-day atlas. Seeing this comprehensive effort as more than just a collection of maps, he aptly named his work *Atlas, or Cosmographical Meditations Upon the Creation of the Universe*. Our endeavour in this Arctic Permafrost Atlas is not so grandiose, but it directly stems from the vision of Mercator to use maps and text to empower the reader through a holistic view of the Earth.

Maps are powerful instruments for conveying timely information to society in a visually pleasing way. Where words might fall short, maps provide an added layer of understanding. On the other hand, words and stories can provide a depth and emotional context that maps may be unable to represent. In this atlas, we tried to deliver maps and graphics of the highest quality mixed with text, art, and stories to explain some of the complex processes associated with permafrost. We aimed to evoke the experiences of people living on and researching permafrost. The 82 maps and graphics, more than 50,000 words and nine stories cannot entirely do justice to the richness and complexity of the "country of permafrost", a term coined by Edward A.G. Schuur and colleagues in 2022, but together they form an unprecedented effort to capture the essence of permafrost characteristics and the issues at stake for local, regional, and global policy in a single document. It was made possible through the involvement of researchers and consortium partners, as well as the results from the European Union Horizon 2020 project NUNATARYUK (2017–2023).

Indeed, this atlas comes at a time when the word "permafrost" has morphed from a relatively unknown academic subject into a topic referenced across news outlets and discussed by the global public. People are aware of it or have heard about it somewhere. Over the past 20 years, the number of web searches for the term "permafrost" has nearly doubled in proportion to the total amount of searches. Part of this growing interest comes from permafrost's relevance to global climate policy and local community adaptation. Climate scenarios show that thawing permafrost will emit between 50 billion and 200 billion tons of carbon dioxide equivalents by the end of the century. Adaptation costs for communities built on permafrost could skyrocket to USD 182 billion by mid-century. The European Union made permafrost one of its eight focal areas in its Arctic policy published in 2021, highlighting the urgency of tackling the issue from both research and policy perspectives.

This atlas is an attempt to translate and consolidate the available knowledge on permafrost. It is a timely book suffused with the compelling enthusiasm of its authors and contributors. Close to a hundred individuals participated in its making, and it does a magnificent job at describing permafrost with maps, words, art, and stories. Far from being an academic product in the traditional sense, it gathers knowledge from the voices of scientists, Indigenous Peoples, northern residents, and local practitioners to provide a holistic and inclusive view of today's challenges in the "country of permafrost".

Prof. Dr. Hugues Lantuit Head Arctic Coastal Research Group Alfred Wegener Institute

J. Hanin

Dr. Peter Harris Managing Director GRID-Arendal

Prologue

The cryosphere is the part of the Earth that is frozen at any time. Snow, ice, glaciers, ice sheets, and ice caps are all part of the cryosphere –and so is permafrost. All the components of the cryosphere contribute to and are evidence of the Earth's cooling system. A sensitive sentinel, the cryosphere reacts to fluctuations in temperature. However, as humans warm the Earth at an unprecedented rate, this cooling system can no longer cope and is shrinking.

The changes in snow and ice across land, glaciers, ice caps, and sea ice have long been evident because they are readily observable and measurable. Reflective sea ice in the Arctic Ocean is being replaced by open waters, while melting ice bodies on land – the Earth's "water towers" – are accelerating global sea level rise. The effects on permafrost, however, are less conspicuous and have received less attention beyond the scientific community and Arctic residents. Nevertheless, permafrost is undergoing rapid changes with implications for the entire planet.

Lying beneath the surface of the land and the ocean, permafrost is seldom visible to us. Yet it is comparable in size to other components of the cryosphere, rivalling Antarctica in spatial extent. Like other parts of the cryosphere, permafrost is absorbing heat. The overall volume of ice melting from permafrost today is similar to the amount of ice annually lost in Antarctica: about 150 billion tonnes of water each year. This is enough water to increase sea level by approximately 0.4 millimetres. The extent to which this water reaches the ocean is uncertain, but the melting ice within permafrost means that the land surface in many ice-rich permafrost regions is "falling" much faster than the sea is rising, creating challenges for communities across the Arctic and along the Arctic coast. What truly distinguishes permafrost from its cryosphere counterparts is not so much its water content but the amount of carbon it contains. The frozen remains of plants and animals in permafrost have long made it a significant carbon sink for hundreds of thousands, or even millions, of years. It is our global freezer! But as permafrost thaws, this organic material begins to decompose, resulting in the production and release of greenhouse gases. Given the size of the permafrost system and the vast quantity of organic carbon it contains, the implications for global warming are serious.

The latest assessment report of the Intergovernmental Panel on Climate Change (IPCC) clearly states the importance of permafrost for the Earth's climate. As permafrost thaw expands with global warming, both its area and volume continue to decrease, leading to broad and irreversible changes in Arctic ecosystems that threaten not only flora and fauna but also the lives, livelihoods, and cultural identities of Arctic peoples. The report also emphasises that the ongoing release of greenhouse gases from permafrost thaw will feed back into climate change, aggravating warming now and for centuries to come, thus increasing the challenges we face as a society – a global society.

What is certain is that over the past 50 years, change has been accelerating in the Arctic. The Arctic has warmed nearly four times faster than the global average because of a combination of climate feedbacks, a process known as Arctic amplification. Over the same period, permafrost has warmed by 2 to 3 °C. If global warming continues, the Arctic will experience further warming and permafrost thawing.

As more attention is directed towards understanding the role of permafrost in the climate system and the various aspects of loss and damage associated with permafrost thaw, an ever-growing body of literature has emerged from both the natural and social sciences. The primary goal of the Arctic Permafrost Atlas is to consolidate these recent findings and offer insights into the diverse aspects and impacts of permafrost and permafrost thaw. In doing so, it aims to provide greater insight into the role and impacts of permafrost thaw, share knowledge to help mitigate uncertainties, and facilitate adaptation to changing conditions, especially with respect to infrastructure, food security, and mobility. It endeavours to be both informative and motivating, contributing to a deeper understanding of this largely invisible yet vitally important component of the Earth's ecosystem.


The Arctic Permafrost Atlas also strives to assess the impacts of permafrost change on Indigenous Peoples of the Arctic. While people across the Arctic may have adopted a more modern lifestyle, Indigenous Peoples continue to maintain close ties to the land as a means of preserving their culture, knowledge, and traditional food supply. However, the dramatic impacts of permafrost thaw on Arctic ecosystems now place thousands of years of cultural tradition at risk.

Finally, there is no single definition of the Arctic. Across this vast area, there are great social, cultural, physical, ecological, and economic differences. Permafrost change itself is highly diverse and specific to each location. In this atlas, the many dimensions of permafrost change are addressed across the northern hemisphere without adhering to a strict definition of the Arctic or associating it solely with the occurrence of permafrost, for, as the climate warms and permafrost thaws, the Arctic as we currently see it will also diminish.



Earth's Freezer

Introduction to Permafrost



Frozen grounds *Permafrost in the Arctic*

Permafrost, ground that remains frozen for at least two years, occurs in Earth's coldest reaches: the Arctic and boreal zones, Antarctica, and high-altitude regions. This vast but largely unseen landscape characteristic influences not only the flora and fauna of these regions, but also the lives, livelihoods, and cultures of the people who live and work there.

The phenomenon of "permanently" frozen ground was known in Medieval Russia but was only scientifically described in 1757 by the Russian scientist Mikhael Lomonosov. The first maps delineating the extent of the so-called "permanent frost" began appearing in the late 1880s for Siberia and then for the entire Arctic in 1928. The term "permanent frost" was eventually replaced by "perennially-frozen ground". Now, the accepted definition for permafrost is: "ground (soil or rock and included ice and organic material) that remains at or below o °C for at least two consecutive years".

Globally, permafrost underlies between 14 and 16 million square kilometres of the Earth's exposed land surface, roughly the area of Antarctica. Most of the world's permafrost is found in the northern hemisphere where it underlies about 15 per cent of the exposed land area. It is found virtually everywhere in the Arctic. The largest expanses of permafrost occur in Alaska, Canada, Kalaallit Nunaat (Greenland), Russia, and on the Tibetan Plateau. Smaller areas of permafrost occur in Iceland, Scandinavia, Svalbard, and in high mountain areas of Europe. In the southern hemisphere, permafrost is found in the Andes and Antarctica. For permafrost on land, scientists distinguish four different permafrost zones based on the proportion of the ground that is permanently frozen: continuous, discontinuous, sporadic, and isolated.

In the **continuous zone**, 90 per cent or more of the landscape is underlain by permafrost. It extends under most surfaces, except for rivers and deep waterbodies, and may reach hundreds of metres deep.

In the **discontinuous zone**, permafrost underlies 50–90 per cent of the landscape. As the name suggests, this zone is broken into areas with and without permafrost, where permafrost does not exist under south-facing slopes. The transition from continuous to discontinuous permafrost generally follows the transition from tundra to subarctic boreal vegetation, although there is no sharp boundary between the two zones. In the discontinuous permafrost zone, the mean annual ground temperature usually measures above –2 °C. Permafrost thickness has a much greater range in the discontinuous zone, varying from a few metres to several hundred metres.

Near the southern extent of the discontinuous zone, permafrost occurs in the **sporadic and isolated patches zones**, underlying 10–50 per cent and less than 10 per cent of the landscape, respectively. This permafrost is patchy in its distribution, tending to occur under north-facing slopes and in peatlands, where permafrost-free areas are common. The thickness of the permafrost ranges from a few metres to over one hundred metres. Permafrost in Finland, Sweden, and mainland Norway largely occurs as discontinuous or sporadic permafrost. Permafrost also exists in vast areas below the seabed. **Subsea permafrost** occurs in the Arctic continental shelves in a large swath extending offshore of the Russian coast from the Barents Sea to the Chukchi Sea and beyond to the Alaskan and north-western Canadian continental shelves. The extent of this subsea permafrost is estimated to be 2.5 million square kilometres. The greatest areas, more than 60 per cent of this type of permafrost, are found in the Kara, Laptev, East Siberian, and Chukchi Seas. In these seas, the mean thickness of permafrost can reach over 400 metres below the sea floor.

Permafrost in profile *Landscape features*

There are numerous unique landscape features and geomorphological processes associated with permafrost environments. The ground above permafrost that is subject to annual freezing and thawing is called the **active layer**. The active layer plays an important role in Arctic ecosystems as this is where most hydrological, biogeochemical, and soil processes occur. The thickness of the active layer primarily depends on the summer heating of the ground as well as the properties of the soil. In general, thicker or deeper active layers occur in areas with warmer summers, although local conditions such as vegetation, duration, thickness of snow cover, and soil moisture also have an influence. In the High Arctic, where summers are cool and short, the active layer may only be a few centimetres thick. Where summers are warmer, the depth of the active layer may reach

a metre or more. In peat-rich landscapes, the active layer may be very thin because the peat insulates the permafrost in summer. The change in active layer thickness is one of the features of climate warming in the Arctic and is thus a focus of scientific monitoring.

Ice wedges are common in permafrost landscapes, mainly seen as polygonal networks on the landscape surface. The ice wedges vary in size from a few centimetres to 3 metres across at the surface and taper down to a point 1–10 metres or more deep. Ice wedges form when vertical cracks open in the ground during rapid winter cooling (causing soil contraction) and subsequently fill with spring meltwater. As the water refreezes, it creates a vertical ice vein in the ground and serves as a natural zone of weakness for repeated cracking in successive winters. Because only the active layer on top of the

permafrost thaws, the wedge shape is maintained over the summer. As the process can repeat each year, the ice wedge widens over time, pushing the soil around the wedge to create ridges. When the soil material contracts during rapid freezing, the stress is ultimately relieved by the formation of frost cracks that intersect each other in a polygonal shape creating ice-wedge polygons. The polygons can range in size from a few to 50 metres across, may host a pond, and are most common in the continuous permafrost zone. They play a role in controlling microclimate, water movement, and greenhouse gas flux from permafrost. When ice wedges melt, they can form networks of trenches or gullies with mounds of the remaining sediment up to several metres high in between. These mounds are home to several species of animals because they provide a unique habitat that is warmer and drier than the surrounding area.

Permafrost is ground (soil or rock) that remains at or below 0°C for at least 2 years. The **active layer** is topsoil that thaws in the summer and refreezes in the winter. It is thinner at high latitude, where summers are shorter and cooler. Ice wedges form when the frozen ground cracks and meltwater freezes inside. When they melt, ponds can form. They vary in size from 10 centimetres to 3 metres across at the surface and taper down to a point at 1 to 10 or more metres deep.

Arctic Ocean

Subsea permafrost occurs under the sea floor of the Arctic Ocean.

BY LEVI WESTERVELD | GRID-ARENDAL (2023)

UNFROZEN

CONTINUOUS 90% or more of the land surface underlain by permafrost DISCONTINUOUS 50-90% permafrost **Pingos** are large conical hills comprising a massive ice core covered with soil and vegetation. They form where water is injected into the permafrost or when a talik refreezes beneath a former lake, leading to growth of an ice core which pushes up the overlying soil. Occurring in both the continuous and discontinuous permafrost zones, pingos range in size from three to more than 50 metres in height and can reach 1,000 metres in diameter. They have either a circular or oval base and a fissured top that may also contain a crater.


Pingo-like features also occur offshore in some shelf areas of the Arctic Ocean. Superficially similar to terrestrial pingos, they typically measure less than 30 metres in height and 300 metres in diameter. The formation of offshore pingo-like features is related to liquid and gas flow driven by permafrost thaw or the decomposition of gas resulting from marine transgression across a permafrost landscape.

Taliks are areas of unfrozen ground within permafrost. Even in the winter, this material does not freeze. This may be due to local thermal anomalies, such as heat storage in surface waters (rivers, lakes, etc.) which keeps the material underneath unfrozen, or to mineralised groundwater flowing through permafrost. In the continuous permafrost zone, taliks are often found under thermokarst lakes and rivers that do not freeze all the way to the bottom.

Thermokarst is an erosional process whereby the thawing of permafrost leads to the settling, collapsing, or slumping of the ground when excess ground ice, such

as ice wedges, melts. Approximately 3.6 million square kilometres, or 20 per cent, of the northern permafrost region is covered by thermokarst landscapes.

Numerous landforms are associated with thermokarst regions. Thermokarst lakes and ponds are found over much of the Arctic landscape, formed in depressions created by the thawing of ice-rich permafrost. They are important ecosystems, providing valuable habitat for many Arctic species. As the permafrost below continues to thaw, these lakes may drain and disappear, as part of the thaw-lake cycle. Other typical thermokarst landforms include retrogressive thaw slumps, erosion gullies, and active layer detachment slides. They occur in hilly terrain, when erosion hits icerich permafrost resulting in the retreat of a hillslope, or when water-saturated soils flow downslope.






Frozen in time *The history of permafrost*

To understand modern permafrost, we must understand the conditions under which it formed and how they have changed. Although the oldest permafrost on Earth likely formed 2.4 million years ago, most permafrost found today was formed during or since the last ice age which ended about 12,000 years ago. Since then, the Earth's climate has alternated between cold glacial periods and warmer interglacial periods affecting the occurrence, distribution, and thickness of permafrost.

The Last Glacial Maximum (LGM), the time when ice sheets were at their greatest extent in the most recent geological past, occurred approximately 23,000 to 19,000 years ago in the northern hemisphere. At that time, the sea level was about 125 metres below current levels and ice sheets covered about 32 per cent of the Earth's total land area (compared with 10 per cent today). Permafrost likely reached its maximum spatial extent (the Last Permafrost Maximum) in the northern hemisphere towards the end of the last ice age. Subsea permafrost formed during periods of glaciation when the sea level was much lower. In the Beringia region during the LGM, for example, the lower sea level exposed large areas of the continental shelf forming a land bridge between North America and Asia. The very cold climate created deep and cold permafrost beneath the exposed land. With the onset of warmer conditions, the ice caps receded, and the sea level rose again, submerging the land bridge. The cold bottom water of the Arctic shelves and ice in the sediment has allowed this permafrost to persist for over 10,000 years after being submerged. As a result of warming from seawater above and geothermal heat flux from below, subsea permafrost has been slowly thawing since it was flooded.


Glacial isostatic adjustment (GIA) plays a critical role in determining the rate of local sea level change and the presence or absence of both subsea and saline permafrost. GIA is the rebound of the Earth's crust after the retreat of large ice sheets. Land that was covered by large ice caps at the end of the last ice age (e.g., much



of Scandinavia and the Canadian Arctic Archipelago), is emerging today. Subsea permafrost is absent in these areas. In contrast, where the land was free of ice (e.g., extensive stretches of the Arctic Coastal Plain in western North America and Eurasia), vast areas of terrestrial permafrost were created and subsequently flooded by the sea, leading to the occurrence of subsea permafrost. Additionally, the influence of GIA on hydrology, temperature, and topography can alter the accumulation and distribution of salt and the formation of frozen ground leading to the occurrence of saline permafrost.




Last Glacial Maximum (LGM). Ice caps and glaciers cover large parts of the northern hemisphere, especially in North America and Scandinavia. Permafrost is present in other areas, especially Asia.




As temperatures rise, ice caps and glaciers retreat (A) resulting in sea level rise (B). Rising seas floods terrestrial permafrost to create susbsea permafrost (C).



Present day. Ice caps and glaciers continue to melt. Climate change drives the degradation and thawing of terrestrial, coastal, and subsea permafrost (D).





An icy balance Arctic permafrost physiography

The way in which permafrost responds to atmospheric warming depends on numerous factors including local soil properties, topography, hydrology, and ground-ice content. A dry organic rich surface layer insulates the ground and can delay thawing, particularly if the permafrost is rich in ground ice. On the other hand, bedrock is a much better conductor for heat, so permafrost in rocky areas responds much faster to warming.

Landscape features such as hills and valleys affect ground temperatures and therefore influence permafrost processes. The temperature of sloping ground depends on its steepness and aspect (orientation towards the sun). In the northern hemisphere, steep south-facing slopes absorb more heat than gentle slopes or slopes that face north. Likewise, on the leeward side of a slope, snow may linger longer than in exposed locations. Snow acts as insulation, keeping the ground warmer rather than cooling it down. Flat depositional landscapes are more prone to the formation of thermokarst lakes while areas with greater relief are more susceptible to slumping and erosion.


Ground ice refers to ice that forms in freezing and frozen ground. Ground-ice content has a tremendous influence on permafrost thaw, with considerable differences between ice-rich permafrost (i.e., permafrost containing large amounts of ice) and ice-poor permafrost (i.e., permafrost that contains little or no ice). The melting of excess ground ice leads to surface subsidence and the formation of ponds and lakes, a process known as thermokarst. In ice-rich permafrost, increased rates of thermokarst development can impact hydrology, ecology, and biogeochemistry as well as infrastructure such as buildings and transportation routes, while in areas with less ice, the effects will be smaller


As the Arctic climate warms and permafrost thaws, the landscape may be subject to great changes, especially in areas with ice-rich permafrost. When ice-rich permafrost thaws, hillsides may collapse resulting in "retrogressive thaw slumps". The thawed sediment becomes saturated with meltwater and starts sliding downhill, where it can threaten infrastructure. If it enters nearby waterbodies, the increased sediment can make the water cloudier and affect aquatic ecosystems. If a thaw slump reaches the margin of a lake, it may rapidly drain as the water runs over the headwall of the slump.

The Peel Plateau in Canada's Northwest Territories may be experiencing some of the most dramatic changes in North America. A rugged landscape with many fastmoving streams combined with ice-rich permafrost is resulting in large-scale landscape changes as the ground ice melts and summer precipitation increases. As ice-rich sediment in the upper permafrost is exposed to thawing, the ground subsides, exposing more permafrost to warming and releasing sediments into nearby waterways and eventually reaching the Arctic Ocean. These processes are happening around the Arctic wherever there is ice-rich permafrost.

Although the appearance and disappearance of thermokarst lakes is not uncommon, the overall trend is towards a decrease in their number. Thermokarst lakes may drain because of the presence of taliks, areas of unfrozen ground in the permafrost that connect to the groundwater underneath, or due to the erosion of drainage channels. As the Arctic warms, thermokarst lakes and shallow lakes without thermokarst origin may become more common in the continuous permafrost zone, while they may disappear in the discontinuous zone and in lake-rich coastal plains.

Also characteristic of some permafrost landscapes are ice-wedge polygons, which form when frozen ground shrinks and cracks. These cracks occur in a polygonal pattern and the resulting ice wedges grow to form a honeycomb pattern across the tundra. As the climate continues to warm, the ice wedges may melt and the polygon centres may be emptied of water. Over time, this can lead to a draining of the landscape and further ground subsidence. In low-lying coastal areas, ice-wedge polygons will become flooded by sea level rise or by rising waters in rivers.





What lies within Organic carbon in permafrost

As long as permafrost stays frozen, the organic carbon within it will remain locked in the ground. But when permafrost thaws, microorganisms become active and start to break down the organic matter releasing greenhouse gases such as carbon dioxide, methane, and nitrous oxide. Methane and nitrous oxide are relatively short-lived but very potent greenhouse gases, while carbon dioxide is much more abundant and can persist in the atmosphere for thousands of years. As the climate warms, the thickness of the active layer (the seasonally thawing layer) increases, accelerating microbial decomposition and increasing greenhouse gas emissions. This in turn stimulates further global warming and permafrost thawing in a positive feedback loop.

Arctic permafrost soils are among the largest carbon stores in the world. Even though near-surface permafrost soils represent only 15 per cent of the total global soil area, they contain almost one-third of the world's soil carbon. Permafrost soils contain almost twice as much carbon as is currently found in the atmosphere. The surface layer (o–3 metres deep) contains 65–70 per cent of the total permafrost soil carbon, while another 25–30 per cent of permafrost carbon is stored at depths greater than 3 metres. The Yedoma region of Siberia and Alaska, which remained ice free during the last glaciation, has rich carbon stores in deep, thick deposits. Additional permafrost soil carbon, less than 10 per cent of the total, is contained in Arctic river deltas.

Subsea permafrost also contains carbon, both as organic carbon frozen in the permafrost and as gas, either in pockets of free gas or trapped in gas hydrates. Most of the gas found in this setting is methane. Though the amount of organic carbon stored in subsea permafrost is poorly understood, it is likely to be considerable, as permafrost extends as deep as 700 metres in some places.

Gases may be prevented from escaping into the atmosphere by layers of overlying permafrost and seawater. However, warming temperatures, increasing stormy weather, and decreasing sea ice have led to rising water temperatures at the sea floor. As the overlying subsea permafrost thaws, methane and other gases may find new pathways to escape through the sediment into the water and then to the atmosphere. Whether and how quickly this ends up increasing greenhouse gas concentrations in the Earth's atmosphere is still being studied.

To understand how permafrost is changing over time, scientists monitor the thickness of the active layer, the amount of groundwater in it, and the temperature of the ground. Measurements taken over many years provide valuable information on how permafrost may be changing and offer insights into its future. Together with satellite observations and laboratory experiments, as well as computer modelling, scientists are building a more complete picture of permafrost and its changes across the Arctic.

Many questions remain unanswered about permafrost thaw and greenhouse gas emissions. Researchers continue to work on getting better estimates of the amount of carbon in permafrost soils and in subsea permafrost; how much and how quickly permafrost will thaw under different climate warming scenarios; how much carbon will be released into the atmosphere from thawing soils, and how much methane, carbon dioxide, or nitrous oxide will be released.. Ice-wedge polygons are highlighted here. They are the product of water infiltrating the soil and freezing year after year, pushing the soil laterally to form these striking geometric features.

SOURCE: MODIFIED FROM VIEIRA ET AL. (2019). BY LEVI WESTERVELD | GRID-ARENDAL (2023)

Thawing permafrost resulted in thermokarst basins, or thaw lakes. The Ibyuk Pingo in Canada is the second tallest pingo in the world at 49 metres. It formed about 1,300 years ago in a drained lake. It is the result of frozen water in the permafrost generating upward pressure pushing the soil into a dome-like feature.

The upper part of the pingo has started degrading.



In the past, a large lake occupied the area. Coastal erosion happening further north has drained the lake. Saline water and the flooding regime have resulted in two types of fundra vegetation, as seen in this image.

> Dead trees transported by the Mackenzie River and redistributed along the shore can help identify the high water mark of floods and storm surges. In this image, logs are marked by white lines.

When ice grows up Pingo Canadian landmark


The greatest concentration of pingos in the world – about 1,350 – is found on the Tuktoyaktuk Coastal Plain in the western Canadian Arctic. In 1984, the Government of Canada formally designated 16 square kilometres of this region as the Pingo Canadian Landmark. Eight pingos are found within this unique protected area, including Ibyuk Pingo.

Standing at 49 metres in height and 300 metres across its base, Ibyuk Pingo is the tallest in Canada and second tallest in the world after the Kadleroshilik Pingo in Alaska which is 54 metres tall. Like most pingos, Ibyuk Pingo grew in a drained lake bottom and is cored by massive ground ice. Ibyuk Lake existed more than 12,000 years ago but probably drained less than 2,000 years ago.

Ibyuk and the other pingos of this region have served as the basis of scientific research for over 50 years but they also carry cultural significance for Indigenous Peoples. The word "pingo" itself comes from the local Inuit language, Inuvialuktun, and means "small hill". The first recorded use of "pingo" in English was in 1938 by the Danish-Canadian botanist Alf Erling Porsild. The name "Ibyuk" is another Inuvialuktun word meaning "two mounds which are close together" and was how local Inuit collectively referred to Ibyuk Pingo and another large pingo, Split Pingo, located to the north of Ibyuk.

Ibyuk is estimated to be about 1,300 years old. Observations show that the upper part was still growing at approximately 2 centimetres per year in the 1980s. At the top is a crater that formed when the overburden at the summit collapsed. The bottom of the crater contains a small pond, which recent observations show may have drained.

All pingos reach their maximum height within several centuries but are also susceptible to erosion. Although still growing, Ibyuk is starting to show signs of collapse. There is already evidence of slumping and erosion on its slopes, giving the appearance that it is splitting into two parts.



Drilling down Learning the secrets of permafrost

Scientists use a range of techniques to study permafrost and how it responds to changing environmental conditions. Core sampling is a technique used to provide information on the vertical distribution of organic matter, ice, and sediments in permafrost landscapes. The corer consists of a long hollow tube screwed into the ground to remove long cylinders, or cores, of permafrost material (sediment, rock, soil, etc.). In the lab, the samples can be analysed for their physical properties, chemical composition, microbial content, age, and other characteristics.

Ground-based measurements using cores are the only way to measure the vertical distribution of ground-ice content, which is critical to developing an accurate understanding of the effects of climate change on permafrost. As temperatures increase, ground ice in the upper layer of permafrost melts, eventually leading to soil subsidence. The sinking ground affects both human settlements and infrastructure (roads, electric poles, pipelines) which were originally built upon solidly frozen ground. As the ground thaws, organic matter that had been stored in permafrost for millennia is broken down by microbial activity and released into the atmosphere as carbon dioxide, methane, and nitrous oxide, which furthers warming. Thawing permafrost also contributes to further climate warming.

The Yukon coastal plain is an example of an area with particularly ice-rich permafrost. Core samples taken here, however, reveal a great variability in groundice content in different deposits, with the highest ice content found in moraines and the lowest in glaciofluvial deposits. More accurate measurements help scientists to better forecast the impacts of climate warming on subsidence, erosion, and the mobilisation of organic carbon in different landscapes.

In addition to core sampling, scientists also conduct continuous, long-term monitoring of ground temperature in boreholes to understand the changes occurring in permafrost. The first systematic observations of permafrost temperature started in the 1930s in eastern Siberia. Early investigations were performed manually and required multiple measurements each day. This meant that researchers could only carry out transient monitoring studies at research stations. Over time, technology evolved, and it is now possible to monitor a range of permafrost parameters (e.g., temperature, active layer depth, moisture content) year- round, either manually or remotely (see graphic on changing permafrost temperatures in section 2.1).

Permafrost temperature is commonly measured using temperature sensors installed in boreholes drilled into the permafrost. Boreholes are classified according to their depth: surface (less than 10 metres), shallow (10-25 metres), intermediate (25-125 metres), and deep (greater than 125 metres). The use of data loggers (small automated digital recording devices) in the boreholes reduces the need for site visits to take measurements. There are now networks of boreholes for long-term permafrost temperature monitoring across the Arctic and in other permafrost regions around the world. As of 2015, there were 1.074 boreholes in the Global Terrestrial Network for Permafrost, with the greatest numbers in Russia (294), United States/Alaska (201), and Canada (194). The oldest borehole in the network is in Russia, drilled in 1957 to a depth of 85 metres.


Permafrost core taken in the spring of 2019 from the Yukon coastal plain. The core was taken to explore the distribution of organic matter, ice, and sediments in permafrost.

The top of the permafrost core is composed of frozen and decomposing organic matter that form a ~6 centimetre-thick brown layer.

> The horizontal stripes are an artefact of the coring technique which involves screwing a metal tube into the soil.

This sample is limited to 60 cm because the coring technique could not penetrate the harder, rocky underlying soil.







Computed tomography (CT) scan A CT scan is similar to an X-ray used by doctors in hospitals. It allows permafrost researchers to see inside the permafrost without

Darker stripes show ice interveined in the sediments.

breaking it apart.

 Dense objects, such as this 3 centimetre-long pebble, appear brightest on the CT scan of the permafrost borehole.





Classified CT scan Sediments in orange (frozen mixture of mostly minerals and organic material with pore ice) and ice in blue (mainly ice, mixed with a small amount of sediment). This allows permafrost researchers to calculate their respective overall contributions.

The top 22 cm of this borehole is the active layer. It undergoes seasonal thawing cycles and hosts plants and organisms which live and grow through the warmer spring and summer months.

Permafrost table

Ice
Sediment

Ice dominates these deeper layers of the permafrost. But smaller pebbles and denser sediments (orange) are present throughout the core.




3D Visualization This 3D visual of the core shows mainly the excess ice content, including ice-rich sediments. Visualizing the data this way helps scientists better understand the structure of the core and the history of the landscape it was taken from.

The cryostructure of ice veins is visible a the top of the core.

Lenticular cryostructure with ice lenses visible above and below the core break.

This suspended ice lens occurs right at the break between the second and third core segment.

These holes in the core are filled with sediments.



© Olga Borjon-Privé (Oluko)

PORTRAIT CEO AND SENIOR SCIENTIST AT B.GEOS GMBH, AUSTRIA Annett Bartsch

Since my school days, I've been interested in the Arctic, drawn in by stories of the vast cold landscapes and research expeditions there. As a geography student, I travelled to Svalbard where I gave presentations about permafrost and geomorphology to other students. Today, I'm a remote sensing specialist, leading international projects and finding novel ways to use satellite imagery to analyse permafrost in the Arctic.

One of the main challenges of mapping permafrost from satellite imagery is that permafrost is a sub-ground phenomenon. But despite this, there are different ways imagery can be used to help map and study permafrost thaw. First, land surface temperatures can help model the presence of permafrost. Second, land surface features associated with permafrost can be seen and analysed on imagery. The mass of lakes across the Arctic, for example, are the result of permafrost thaw. Permafrost can also be monitored from space by analysing ground subsidence and upheaval, which are also tied to soil properties: the more ice in the ground, the greater the change at the surface over a season. An added challenge of mapping landscape features is the very frequent presence of clouds in the Arctic. Scientists like me make use of radar satellite data which can see through the clouds, but this type of data is still expensive and hard to obtain at a very high resolution.

In the summer, I spend many nights in Siberia or Scandinavia sleeping in tents and collecting field measurements. These measurements help connect what is measured on the satellite images with what is happening on the ground. Collaboration with local experts is very important. In Siberia, we work with Russian scientists who often have camps next to boreholes that contain instruments to measure permafrost. Nowadays, drones are also used to close the gap between satellite imagery and field observations.

One of the main goals when applying remote sensing to permafrost studies is to improve climate models and to assess the impacts of permafrost thaw. Permafrost is thawing across the entire Arctic and the implications are complex, both locally and globally. With remote sensing alone, we can make very interesting maps, as we've done in recent work mapping all infrastructure found within 100 kilometres of the Arctic coastline. But if we want to answer the big questions, we need to go one step further. For this, you need specialists from different fields to work with remote sensing experts, including Earth climate modellers and field specialists, that have a good understanding of local processes.

Many questions remain open, but technology has already come a very long way in helping to answer them. During my studies, it was very difficult to get access to data, and the resolution was very poor compared with today. Now, there is so much data that it's challenging to find the methods to process everything. It requires a lot of computing power. We need artificial intelligence and machine learning to deal with these data, which in the coming years will be central to cloud computing.

Un/settled *Life on frozen ground*

Humans have lived in the Arctic for thousands of years, adapting to life in this harsh environment and passing their knowledge on to successive generations. Today, approximately 5 million people reside in the Arctic circumpolar permafrost area, the area defined by the northern circumpolar permafrost extent within the eight Arctic nations. Communities within this extent are known as permafrost settlements. There are currently 1,162 permafrost settlements in the Arctic. Almost 90 per cent of them have less than 5,000 inhabitants, but most people in this region, approximately 4 million, live in the larger 123 settlements that range in population size from 5,000 to 360,000. Most of the large settlements, 85 per cent, are found in the Russian Arctic.

The greatest number of permafrost settlements, over 65 per cent of them, are in the sporadic permafrost zone, in the southern part of the permafrost region. Unless situated directly on permafrost, these communities may not be immediately affected by permafrost thaw, but all of them will be affected by the changes occurring in the surrounding environment. Settlements in the continuous and discontinuous permafrost zones may experience significant impacts as the underlying permafrost degrades because of climate change. This includes Yakutsk and Norilsk, two of the largest settlements in the Arctic with a combined population of 486,000.

Constructing long-lasting, stable structures on permafrost is a major challenge since permafrost is increasingly neither permanent nor stable. As human activity expands in the Arctic and as global warming continues, it becomes more important to understand how infrastructure may be affected and how it can be made more resilient. Building on frozen soils can cause the underlying permafrost to thaw and sink, resulting in damage to buildings. This can result in anything from minor cracks in the walls to major structural failure. Transportation infrastructure – roads, airstrips, railways, bridges, etc. – is also subject to significant damage from permafrost thawing. Engineering solutions, such as building road embankments that allow heat to escape, are required. In areas that are too wet for roads, people only drive on winter ice roads that are built on frozen marshes or on frozen lakes and rivers. The thick ice can support very large trucks bringing equipment and supplies to communities and industrial operations. These roads are seasonal and need to be rebuilt every winter; and as the climate continues to warm, they are becoming less reliable and shorter lived.

Unlike in other areas, most of the groundwater in permafrost is frozen. In areas of discontinuous permafrost, it may be possible to drill through unfrozen ground to reach groundwater. In areas of continuous permafrost, however, drinking water often comes from lakes, rivers, or melting ice and snow. Some communities build insulated and elevated water pipes to supply running water to their homes.


Industrial operations (e.g., mining, oil, and gas exploration) also require special considerations to prevent the permafrost from thawing. The deep wells required for oil and gas may be lined with cement to prevent them from collapsing, while the equipment at the surface can be built on concrete pads. Because the oil must be kept above 60 °C to allow it to flow, pipelines are often built above ground so as not to thaw the permafrost and risk damage to the pipes.


Indigenous Peoples and local communities depend on permafrost for more than just stabilising buildings and other infrastructure. Many still live traditional or semi-traditional lifestyles, where they depend on hunting, fishing, and gathering to support themselves, their families, and their communities. Knowing and understanding the characteristics of permafrost are vital for these needs. Reindeer herders, for example, rely on their knowledge of permafrost to determine how quickly the passage of their herds will turn the surface to mud. This determines where the herders will establish their camps, how long they can stay, and what herding operations they can perform. The effect of permafrost on the local landscape and vegetation also plays a role in influencing reindeer behaviour.

People also use permafrost as a nature-based solution for preserving food. For thousands of years people have harnessed the stable sub-zero temperatures of permafrost and built underground cellars in which to store fish and game. These traditional ice cellars are dug into the upper permafrost below the active layer with a doorway to protect the entrance area from warm summer air and discourage the accumulation of insulating snow in the winter.


Arctic permafrost settlements

NOTE: Only settlements with a population over 5,000 are labelled.









Frozen states I *Russian Federation*

Most of the permafrost in the Arctic is found in Russia. Approximately 65 per cent of the country is underlain by permafrost, mainly in the continuous zone. Russia encompasses 85 per cent of the Arctic's large permafrost communities, i.e., those with 5,000 inhabitants or more.


In contrast to most other parts of the Arctic, most Russian permafrost settlements are inland and on continuous permafrost. Of the 713 permafrost settlements, 401 are in the continuous zone, 241 in the sporadic zone, and 71 in the discontinuous permafrost zone.

Forty distinct groups of Indigenous Peoples live in the North, Siberia, and Far East of Russia. The largest of these are Dolgan, Nganasan, Nenets, Saami, Khanty, Chukchi, Evenk, Even, Enets, Yupik, and Yukagir. Many continue to rely on the traditional activities of reindeer herding, fishing, and hunting.

Of the estimated 2.2 million domesticated reindeer in the Arctic, 1.5 million are in Russia. The main reindeer herding areas are in Yamal-Nenets, Sakha Republic (Yakutia), and Chukotka. The Nenets people are thought to be the largest group of nomadic reindeer herders in the world. Almost half of the Nenets people (approximately 20,000), continue to herd reindeer today. Reindeer herders are often nomadic, moving with their herds from winter pasturelands in the forest tundra/northern taiga zone to the midsummer pasturelands on the coast of the Arctic Ocean. Each group of herders may tend to anywhere from hundreds to thousands of reindeer. The seasonal changes caused by the thawing and freezing of the ground influences the activities of the reindeer herders. In some areas, the thawing of the active layer in the summer can turn the ground to a semi-liquid mud layer, making travel difficult and even dangerous for both herders and reindeer. This sludge layer can occur from both natural causes and human or animal activity, such as trampling. By "reading" local conditions, experienced herders can determine the likelihood of this phenomenon occurring and move their herd to safer ground.

The permafrost landscape also influences travel. Landforms such as ice-wedge polygons must be traversed at the correct angle to avoid breaking the sledge. Likewise, the narrow ravines created on hillsides or riverbanks by permafrost thawing and sliding are used by herders to climb up and down steep slopes in a safer manner.

Herders also use their knowledge of the landscape and vegetation – both of which are influenced by permafrost processes – to monitor and predict the behaviour of their reindeer, moving them to desired locations as required. Similarly, the different waterbodies in permafrost landscapes also influence the movements of herders and their animals. Thermokarst lakes are important to herders, who believe them to be cleaner than other sources of water for drinking. These lakes also help to cool the reindeer and escape insects in the summer.





PORTRAIT CHAIR OF COUNCIL OF YUKAGHIR ELDERS, NELEMNOYE, RUSSIA Vyacheslav Shadrin

I was born in 1967 in the village of Nelemnoye in the Upper Kolyma region of the Sakha Republic (Yakutia). I am both a researcher focusing on the history and culture of Indigenous minorities of the Russian North, and the Chairman of the Council of Elders of the Yukaghir people. My role as Chairman means I travel a lot, visiting communities across the Russian Arctic.

Permafrost underlies 80 per cent of Yakutia, including my home village of Nelemnoye, which is located just below the Arctic Circle in an open, subtundra forest. This Yukaghir village is composed of 40 buildings situated a few hundred metres west of the Yasachnaya River, a tributary of the Kolyma River. About 300 people live in the village, 200 of which are Yukaghir, an Indigenous Peoples group from north-east Asia. According to the latest population census, only 1,603 Yukaghir remain today.

Changes in climate limit our traditional fishing and hunting practices. Autumn sea-ice fishing, which historically would start around the end of October, can no longer take place, or has become very dangerous, as any ice is still quite thin then. Hunting practices have been similarly affected. Musk deer hunting usually takes place in November when the animals migrate from their summer to winter pastures. But as the ice road is no longer accessible this early, traditional hunting grounds are impossible or very dangerous to access. Hunting that occurs later in the winter is less profitable, as the quality of fur is lower.

Reindeer herding, an important livelihood among traditional communities in the Arctic, is also affected by permafrost thaw. The lowlands are wetter and muddier, and reindeer sometimes lose weight and get ill. Herders are also increasingly dependent on technology such as GPS to find their way. Thirty or forty years ago, it was impossible to imagine that a herder could get lost; they knew the tundra extremely well. Losing their traditional knowledge makes them more vulnerable.

Traditionally, underground cellars dug in the frozen ground are used to store meat, fish, and berries. Now, these underground freezers are thawing due to the warming climate. Many are collapsing and no longer in use. They are being replaced with refrigerators, but these require constant electricity and are vulnerable to power outages. I estimate that 80 per cent of the diet of inhabitants in Nelemnoye now consists of food bought in shops. In my youth, only goods such as sugar, salt, and tea were bought from shops; the rest was harvested from nature. Cases of diabetes are increasingly frequent in northern Indigenous communities. We are victims of this system. We do not choose this lifestyle.

As the number of fishers and hunters dwindle, more people are now scavenging for mammoth tusks and bones, which can fetch high prices on the black market. These appear at the tundra's surface as permafrost thaws. But scavenging mammoth tusks is against traditional beliefs. Traditionally, we associate mammoths and their spirits with the underworld. Bad things come from there. In the past, remains of mammoths mostly appeared during floods and on eroding riverbanks, so they are associated with such events. When these tusks and bones resurface in the tundra, a door is opened between the underworld and the middle world where people live.

Politicians and scientists always talk about the need to develop adaptation strategies. But these aren't for us. It is our life, our everyday. We live in these conditions, and we are adapting all the time. For Indigenous Peoples, climate change is a real threat that brings about many challenges which impact our traditional livelihoods and lead to fewer people being involved in traditional activities.

Frozen states II North America

In North America, the Arctic circumpolar permafrost area encompasses the state of Alaska in the United States of America and Yukon, Northwest Territories, Nunavut, Nunavik, Nunatsiavut, and southern Labrador in Canada. Approximately 80 per cent of this region is underlain by permafrost. It is dominated by continuous and discontinuous permafrost in the western Arctic and the Arctic Archipelago, while sporadic and isolated permafrost are becoming more prevalent in the eastern Arctic and interior of Alaska.

Of the 300 permafrost settlements in the Canadian and Alaskan Arctic, 63 are in the continuous permafrost zone, 69 are in the discontinuous zone, and 168 are on sporadic permafrost. Almost half the settlements in this region are situated on the coastline and most of these are on continuous permafrost. While many Indigenous Peoples rely on subsistence hunting and fishing, resource extraction from mining and oil and gas development are important economic activities in both countries.


Permafrost creates unique challenges for the many communities in this region. Permafrost coastlines are exposed to high rates of coastal retreat due to longer periods of open water along the coast, warmer summers, and increased frequency and intensity of storms. Coastal communities, especially those in the western Arctic, are affected by erosion and subsidence, which affects local infrastructure, land cover, and cultural sites.


More than 20 communities in Canada and Alaska have already been declared threatened because


of coastal erosion. In Shishmaref, Alaska, the rate of coastal erosion is so high that the inhabitants have voted to move the town to a new location on the mainland. In the Hamlet of Tuktoyaktuk in the Northwest Territories (Canada), some houses deemed most at risk from collapsing into the Arctic Ocean have been moved further inland at significant cost. The hamlet is now implementing a plan to slow erosion, but eventually, the community will need to consider relocation.

Industrial and military facilities and critical community infrastructure along the western Arctic coast of North America are also threatened by accelerating coastal erosion and storm surges. Like all structures in the Arctic, they were designed for stable permafrost conditions but now many show signs of instability because of increased thaw settlement and seasonal frost heaving. Many will face costly retrofits if they are to remain viable in the future. Numerous sumps, designed to contain contaminants such as drilling mud in permafrost at former oil and gas exploration wells in the Mackenzie Delta region, are now unstable or collapsing, allowing toxic fluids to escape.

Subsea permafrost occurs offshore of the western Canadian and Alaskan Arctic, as well as off northern Eurasia. While there is currently limited offshore development, there is considerable potential for largescale oil and gas development in the future. Knowledge about the distribution and properties of subsea permafrost is still limited and more research is required to ensure the safety of operations in this environment.







PORTRAIT ENVIRONMENTAL COORDINATOR AT GWICH'IN RENEWABLE RESOURCES BOARD, AKLAVIK, NWT CANADA

Jessi Pascal

I would describe myself as independent, family oriented, and a lover of the outdoors. I have three siblings and we live in Aklavik, a hamlet with a population of approximately 500 located in the Inuvik region of the Northwest Territories, Canada. I recently moved back to Aklavik from Inuvik – the administrative centre of the Inuvik region – located approximately 50 kilometres east of Aklavik.

I would describe Aklavik as a community where everyone knows everyone, and where everyone helps each other. That's what I like most about Aklavik. The village is situated along the Peel River [Peel Channel of Mackenzie Delta], providing our community with all the water we use for drinking, showering, and washing. During the winter, it is cold and white. The mountains to the west are absolutely beautiful. A lot of people rely on wood stoves as a heat source, including my folks, because purchasing oil for the furnace is quite expensive. I haul wood for them whenever I can.

Spring is the time for goose hunting and carnivals. Each community in the Mackenzie River Delta has these carnivals, which include attractions like snowmobile races, dog sled races, and egg toss competitions. These were cancelled in 2021 due to COVID-19. In the summer, many people including my family go fishing, either in the Peel River or down at Shingle Point in the Arctic Ocean. In the fall, a lot of people hunt rabbits, either in the fields or over on the delta. There is always something happening.

One of my first encounters with permafrost was when my dog was digging in the ground and suddenly reached the frozen permafrost table. I wasn't very familiar with what permafrost was when I was a kid. But I thought it was pretty interesting, that a dog could just dig a couple of feet and suddenly bump into it!

Being located on permafrost means Aklavik has had to adapt in many ways. None of the pipes, for example, are underground. We get our sewage removed by a vehicle that comes in. All the buildings are built fairly high off the ground to protect them from flooding. But due to permafrost thaw, a lot of infrastructure is at risk. Recently we had to borrow some jacks and bumped my father's house to make it level again.

There is no road to get in or out of Aklavik. Instead, boats are used in the summer as a mode of transportation across the Mackenzie Delta. In the winter, snowmobiles and trucks can be driven on the ice road. But with climate change, these landscapes are changing. I go out with my friends to check our traps and it is really cold out, and then suddenly we are sinking in slush. It's very unpredictable. I am not sure if it is directly related to permafrost.

Reflecting on the impacts of permafrost thaw on my community, it's clear that people here know very little. It wasn't part of my high school curriculum, for example. I feel it is not treated as a priority for people to know about. I have interviewed many people as part of my work with the Joint Secretariat of the Inuvialuit Settlement Region about topics such as permafrost, fish, animals, weather, and so on. But the subject I've had most trouble with is permafrost. This is a subject that folks know and have heard about, but don't understand its science aspect and how it affects our lives. I think a little more education on the topic would be beneficial for our community. After all, it is literally right under our feet.

Frozen states III Nordic region

Permafrost in the Arctic's Nordic region is found in Kalaallit Nunaat (Greenland), Iceland, in the northern parts of Norway (Nordland, Troms, Finnmark), Svalbard, Sweden (Norrbotten), and Finland (Lappi). There are 149 human settlements in this region, most located in the sporadic permafrost zone. Only Kalaallit Nunaat and Svalbard have settlements in the continuous permafrost zone (23 and 2 settlements, respectively), while only Kalaallit Nunaat has settlements in the discontinuous zone (22 settlements). All permafrost settlements in Kalaallit Nunaat and Svalbard are coastal.


Although geopolitically part of Europe, Kalaallit Nunaat is geographically and geologically part of North America, and as such shares many characteristics with the eastern Canadian Arctic. As in the eastern Canadian Arctic, many homes here are built on bedrock, meaning they are unlikely to be destabilised by permafrost thaw and so are less vulnerable to collapse than sedimentary and ice-rich permafrost sediments. Other infrastructure, however, is often found on vulnerable permafrost. The area between the parking lot and runway at Kangerlussuaq airport has been severely affected by permafrost thaw. Damage to the surrounding community infrastructure, especially the road networks, has also been reported.

North-east Kalaallit Nunaat is home to the Zackenberg Ecological Research Operations, a research station established in 1995 with the goal of facilitating research into High Arctic ecosystems. Located 74 °N, the station is in the continuous permafrost zone with an active layer thickness of 20–100 centimetres. The remote location was believed to be ideal for conducting long-term ecological studies of at least 50 years. Research here contributes to a range of international programmes, including numerous studies on permafrost. Now, as elsewhere in the Arctic, the research station itself is under threat. Permafrost thawrelated erosion along a nearby riverbank has intensified. Now, rather than lasting 50 years, some buildings may need to be moved or rebuilt soon to avoid collapse.

The Svalbard archipelago is part of Norway but in many respects is a world away from it. Situated midway between mainland Norway and the North Pole, this remote archipelago was largely unpopulated until the first settlements - small outposts - started appearing in the 1600s. Today fewer than 3,000 people live on Svalbard. Like Kalaallit Nunaat, Svalbard is also an important location for permafrost research in the Nordic region of the Arctic. Permafrost thickness in Svalbard ranges from less than 100 metres in low-lying areas to more than 500 metres in the high mountains. Extensive permafrost monitoring takes place on Svalbard, and long-term observations of the active layer thickness and permafrost ground thermal regime are included in the Svalbard Integrated Arctic Earth Observing System.

Svalbard's permafrost also plays a vital role in helping protect the world's food supply. The Svalbard Global Seed Vault is built into the permafrost on a mountainside. The vault provides for the long-term storage of seeds from around the world, backing up plant genetic diversity in the case of disease or other disaster. Rather than a hindrance, the permafrost was considered an added benefit of the site because it would aid in preserving the seeds.

Although there is significantly less permafrost in southern Norway, it does exist in steep bedrock slopes, including at lower elevations on north-facing slopes. This area has been subject to more scientific study in recent years because of the increased frequency of rockfalls and rock-ice avalanches during hot summers. These occurrences can put both people and infrastructure at serious risk, either through direct impact or from associated events such as flooding.





© Olga Borjon-Privé (Oluko)

PORTRAIT MAYOR OF AVANNAATA MUNICIPALITY, KALAALLIT NUNAAT (GREENLAND) Palle Jerimiassen

To say it plainly: I love my job as mayor! When I get to work in the morning, I don't know what the day will bring. It's always something different, and new challenges emerge every day. Most of the work happens in front of the computer, working with different reports, future plans, and many meetings with different people. I have coffee meetings with the economic sector and soda meetings with young people. In politics, one must work to bring different opinions together. That is something I like to work with and something I think I'm good at.

Avannaata is a big municipality, the world's biggest, with 21 villages and four towns, and it's important that I am in contact with the whole population. There are rather big cultural differences within the municipality. Ilulissat is a multi-ethnic town with 5,000 people, and about 30–40 different ethnic groups. Up in the northernmost parts, in the Qaanaaq district, there are still communities that primarily live off hunting. The last village to get public electricity supply was two years ago. So, there are quite big differences in the way we live within the municipality, and I always have to think about how the decisions we take here in Ilulissat impact the people in other communities. It's all connected, and that's part of what makes it interesting.

Twenty years ago, some scientists came here and said that in 10 years' time there will no longer be ice in front of Ilulissat, but no one believed them, and neither did I. But I clearly see now that climate change has come to stay. I know that the fishers and hunters are very concerned about how the climate is changing, the unpredictable weather forecasts, and the changes that are happening in animal migration. It's something we all talk about daily because we base our whole lives on our surroundings.

Regarding infrastructure, we have many people complaining about the conditions of the roads, why their houses end up crooked, and why the sewage leaks. We are getting bigger and bigger challenges with climate change and permafrost thaw. The places where roads are built on permafrost become like the small roller coaster in Tivoli Gardens. The land around houses sinks and causes damage to their structures. Qaanaaq, for example, is built on land where the whole area has started moving. That means that everything becomes crooked. We've also had issues with the underground sewage system. Look at the old football pitch where sewage water started leaking out!

We must take climate change into account, we have to, especially up in the north where the whole way of life is completely changing. We made the United Nations Sustainable Development Goals a starting point for the municipality's development strategy. We have also considered permafrost in all our plans. In some places, new houses need to be built on bedrock. That's difficult in places like Qaanaaq where there aren't many suitable places around. We started with a road leading from Ilulissat to the new airport, but there have been challenges trying to figure out where to place the road on solid ground. In the future, we have to develop the sewage system above ground in places where there's permafrost so that we don't have the same issues we have in town now.

We get a lot of visits from all types of people. We say to them that we all must take responsibility. We are most impacted but even though we are only a small part of the whole world, we also have to take responsibility ourselves to reduce climate change.



Awakening Giant


Permafrost and Climate Change

Warming up, warming down Increasing ground temperatures

Permafrost is defined according to temperature: ground that remains at or below o °C for at least two consecutive years. Despite its name, permafrost is not permanent. Changes in the environment, both natural and human-caused/anthropogenic, may cause the ground temperature to rise above o °C, leading to permafrost thaw. Depending on the extent of the warming, permafrost thaw may trigger other environmental effects, including landscape changes and the release of greenhouse gases.

Temperatures at Deadhorse, Alaska

Mean annual values (° C) from 1987 to 2020



SOURCE: ROMANOVSKY, V. (2021). ADAPTED BY LEVI WESTERVELD | GRID-ARENDAL (2023)

Permafrost temperature is linked to ground surface temperature, which itself is linked to changes in air temperature and snow cover dynamics. As the temperature of the permafrost surface (i.e., the ground below the seasonally thawed active layer) increases, it conducts heat to deeper permafrost resulting in warming at depth as well. One way in which scientists monitor temperature change in permafrost is through sensors installed in boreholes drilled into the ground. This can be just a few metres deep in warm permafrost to more than 20 metres in cold permafrost and bedrock. Long-term monitoring of borehole temperatures shows widespread evidence of permafrost warming across the Arctic in response to global climate change, with dramatic increases in some locations.


During the 2007–2009 International Polar Year (IPY), ground temperature measurements were made at 575 locations around the circumpolar Arctic. Although longer-term records exist in some places, the IPY initiative added many new monitoring sites to provide a more complete network across a range of permafrost locales. Collectively, these sites provide an important baseline against which to measure permafrost change across the Arctic.

Measurements since the IPY reference period show that annual mean permafrost temperatures are increasing at most sites across the Arctic. Between 2007 and 2016, ground temperature in the continuous permafrost zone has increased by 0.39 °C on average. The greatest temperature increase in individual boreholes within continuous permafrost was recorded at Marre Sale on the Yamal Peninsula in north-western Siberia (+0.93 °C at a depth of 10 metres) and on Samoylov Island in the Lena River delta in north-eastern Siberia (+0.95 °C at a depth of 20.75 metres). Within the discontinuous permafrost zone, ground temperature was found to have increased by 0.2 °C with the greatest increase in Magadan in the far east of Siberia (+0.95 °C at a depth of 10 metres).


Permafrost can be described as either cold or warm, based on its temperature deep below the surface where it changes slowly from year to year. When cold permafrost, which is below -1 °C, is heated, it undergoes noticeable warming. In contrast, warm permafrost refers to permafrost with temperatures close to o °C. Within this type of permafrost, the presence of ground ice acts as a buffer against warming by absorbing heat and melting. However, once the ground ice has melted completely, the ground heats up and is no longer permafrost. The mountainous areas of Europe, as well as Central Asia and the Andes, tend to have warm permafrost that contains less ice.

The rate of permafrost warming is not uniform across the Arctic, varying according to local environmental conditions such as climate, snow cover, ground ice, vegetation, and organic layer thickness. The greatest changes have been observed in cold permafrost in the Arctic and High Arctic where warming may reach up to 1 °C per decade. While there have also been temperature increases in warm permafrost, these have been smaller, generally less than 0.3 °C per decade.

Warming permafrost has led to a global concern about the potential for the release of large amounts of greenhouse gases. Permafrost soils hold vast amounts



of organic material. When thawed, this material is exposed to microbial decomposition, producing either carbon dioxide, methane, or nitrous oxide. Current projections suggest that the combined release of carbon dioxide and methane from permafrost thaw will be equivalent to 100-200 billion tons of carbon dioxide by 2100. By comparison, global net greenhouse gas emissions from anthropogenic sources were equivalent to approximately 59 billion tons of carbon dioxide in 2019. The additional contribution from permafrost will make it even more difficult to limit global warming to 1.5 °C, as set out in the Paris Agreement of the United Nations Framework Convention on Climate Change. Warming permafrost can also bring about landscape changes such as erosion and landslides which impact natural ecosystems, traditional lifestyles (e.g., hunting and fishing), and built infrastructure.



The chill is gone Thickening of the active layer


The active layer is the ground above the permafrost that thaws each summer and refreezes with the return of lower temperatures in autumn. How deep the active layer thaws depends on local conditions, tending to be thinner in the colder high Arctic and thicker in the southern extent of permafrost where temperatures are higher. Seasonal thawing of the active layer is very important to Arctic ecology as this is where important biogeochemical processes take place. Because it is more responsive to short-term climatic variation, the depth of the active layer can vary considerably from year to year. However, long-term trends in active layer thickness are, together with the permafrost temperature itself, an important indicator of change in the thermal state of permafrost.

Active layer thickness has been monitored at numerous sites across the Arctic since the 1990s. It is most commonly obtained with an active layer probe, an instrument that can be pushed through the unfrozen sediment until it reaches the frost table. Because it is strongly influenced by seasonal snow cover and air temperatures, it can be more difficult to discern trends in active layer thickness than in permafrost temperature. Where longer data sets are available, observations generally show an increase in active layer thickness.


The Circumpolar Active Layer Monitoring network has been monitoring changes in the active layer and


near-surface permafrost since 1991. The greatest increases in active layer thickness have been observed in western Siberia and the northern European region of Russia where thickness increased by as much as 0.4 metres between 2009 and 2019. A similar trend has been observed at sites in Scandinavia, Svalbard, and Kalaallit Nunaat (Greenland), although with greater annual variation. The greatest changes in North America were recorded in the Alaskan interior where active layer thickness increased by more than 0.2 metres between 1996 and 2019. Active layer thickness from the Alaska North Slope, through Canada's Mackenzie River valley and into the eastern Canadian Arctic, has experienced both increases and decreases, but current measurements remain similar to those from the 1990s.

Challenges have been identified in assessing trend changes in permafrost using ground temperature and active layer thickness measurements. Active layer thickness is usually measured using the ground surface as a reference point. However, thawing in ice-rich locations may result in ground subsidence and a lowering of the permafrost table, and unless ground surface elevation is also monitored, active layer deepening is underestimated. Results can also be affected by the time of year that measurements are taken, particularly whether they are taken at the time of maximum thaw or not, which itself can vary from year to year.



SOURCE: OBU ET AL. (2021). BY LEVI WESTERVELD | GRID-ARENDAL (2023)






Disappearing act Declining permafrost extent

Terrestrial permafrost underlies approximately 14 million km² of the exposed land area in the northern hemisphere. When permafrost thaws, it can have significant impacts on terrestrial ecosystems, landscape processes, the hydrological cycle, greenhouse gas emissions, and infrastructure. The sensitivity of permafrost to climate warming and the potential for large-scale environmental changes make it important to understand the full extent of projected permafrost changes. All the latest climate models show continued warming leading to permafrost thaw, but the amount of permafrost that may be lost varies considerably under the different Earth system models due to a lack of data and limitations within the models themselves. Permafrost is generally considered to be lost when the active layer does not refreeze fully in the winter. This means that full thawing of the permafrost, down to hundreds of metres, can take much more time.

One approach for estimating future permafrost loss under different warming scenarios looks at the relationship between mean annual air temperature and permafrost occurrence. One study estimated that if the climate is stabilised at 1.5 °C above pre-industrial levels, between 2.6 and 6.8 million km² (over 30 per cent) of the present-day near-surface permafrost area could be lost. Stabilising the climate at 2 °C above pre-industrial levels would see that loss increase to between 4.4 and 8.6 million km² (over 40 per cent) of permafrost area. If warming increases even more, a large part of the existing near-surface permafrost will thaw. With 5 °C of warming, near-surface permafrost extent will be reduced to 0.3–3.1 million km², and with 6 °C of warming, it could disappear altogether.

The rate of permafrost degradation is not uniform across the Arctic. Permafrost degrades quickest in the western Russian Arctic where mean annual ground temperatures have increased from 0.03 °C to 0.06 °C per year in the continuous permafrost zone and the permafrost table has lowered by as much as 8 metres in the discontinuous zone. Between 1997 and 2018, scientists have observed distinct signs of permafrost degradation in this region: a progressive increase of the active layer thickness, enhanced permafrost degradation after ground ice has melted, and a lowering of the permafrost table.



Microorganisms, macro effects Permafrost carbon cycle

Permafrost contains vast amounts of organic carbon, the carbon found in all living organisms and their remains. The organic carbon in permafrost consists mostly of dead animals and plants that have accumulated and remained frozen for thousands of years. As temperatures increase and permafrost thaws, this organic matter starts to decompose and release carbon dioxide and methane into the atmosphere.

Most of the Earth's carbon is stored in rocks. The rest is found in the atmosphere, soils, oceans, and living organisms, and cycles through these reservoirs in a process known as the global carbon cycle. In the permafrost subcycle, carbon is transferred from the soil to vegetation and microbes, released into the atmosphere, transferred back into vegetation via photosynthesis, and finally returned to the soil where the process starts again. As long as the soil is frozen, microbial decomposition of soil organic matter is drastically slowed down and carbon is able


to accumulate. Many microorganisms are able to survive the freezing temperatures either by slowing their metabolism, or by entering a resting state. When permafrost thaws, the frozen water in the soil melts, making the organic matter accessible to soil microbial decomposition. While one group of microorganisms, methanotrophs, are able to consume methane (and are the only known sink of methane), most processes occuring during permafrost thaw increase the release of greenhouse gases into the atmosphere.

The presence or absence of oxygen during decomposition determines whether carbon is released as carbon dioxide or methane. In aerobic soils, where oxygen is available, such as collapsed and drained sites, the decomposition of organic carbon releases more carbon dioxide. In contrast. anaerobic decomposition in oxygen-limited and often waterlogged conditions (e.g., in permafrost thaw lakes) releases both carbon dioxide and methane. Although

the thawing of permafrost soils leads to deeper rooting vegetation and more carbon sequestration, this cannot compensate for carbon emissions from these soils while they thaw. Due to the potential vulnerability of large permafrost carbon stores, their degradation is globally important. Converting even a small portion of the permafrost carbon to greenhouse gases could increase the rate of climate change, spurring further permafrost warming and even more greenhouse gas emissions in a positive feedback loop.

Permafrost contains a lot of organic matter that is rich in carbon. At freezing temperatures. mircoorganisms are not that active, and the organic matter is not decomposed.









Different types of permafrost landscapes show varying responses to warming. Permafrost thaw is often divided into two main types: gradual thaw and abrupt thaw. Gradual widespread thaw occurs from the upper permafrost layer down into ever deeper layers. This exposes organic matter to decomposition over decades or even centuries. Abrupt thaw occurs where erosion takes place along riverbanks, lake shores, and coasts, and in ground containing large amounts of pure ice. The ice can make up much of the ground volume and also cements the frozen ground together. As warming penetrates the ground, permafrost collapses as the ice melts away. This ground surface collapse (called thermokarst) accelerates the thaw, and instead of thawing centimetres each year, several metres of permafrost can become destabilised within days or weeks. The landscape changes create new lakes, ponds, landslides, and other landforms.


Along the Alaska North Slope and into Canada, in an area that runs from the northern edge of the Brooks Range to the Beaufort Sea, the tundra landscape is rich in both ground ice and organic matter. In the foothills of the Brooks Range mountains and along river valleys, the thaw of ice-rich permafrost triggers slope failures and mass wasting of sediments. This not only leads to greenhouse gas production as frozen organic matter starts to thaw and decompose, but also releases large amounts of sediments and pollutants into aquatic ecosystems. On coastal lowlands, the degradation of ice-wedge polygon tundra leads to the formation of large thaw lakes as well as rapid coastal erosion.


Rapid thawing is also occurring along the Mackenzie River valley in the Northwest Territories (Canada), where much of the landscape is comprised of thick, ice- and carbon-rich sediments. Thick peat deposits have also accumulated on poorly drained and flat terrain. As the climate warms, these landscapes are affected by ground-ice loss, subsidence, and changes to surface hydrology.

The western Hudson Bay Lowlands are characterised by extensive and deep peat deposits which have accumulated since the retreat of the last inland ice sheet. These are the world's second largest peat deposits, after the west Siberian lowlands. Much of the peat accumulated in a slightly warmer climate during the Holocene thermal maximum, but as the climate cooled a few thousand years ago, the active layer became shallower and turned peat across the area into a frozen state. The frozen peatlands are rich in ground ice, and as they are close to the limits of their climate zone, they are vulnerable to permafrost thaw.









Faster, deeper, stronger II Speed of thaw in Scandinavia and the Russian Federation

In Scandinavia (excluding Svalbard), permafrost is dominated by two distinct types: mountain permafrost and lowland permafrost. Mountain permafrost occurs mainly in bedrock and does not contain much organic material, meaning that there are limited climate feedbacks. Thaw of mountain permafrost can destabilise the terrain causing rock falls, landslides, or other mass movements. Lowland permafrost occurs in relatively flat terrain where ice-rich peat and other sediments accumulate in ice-wedge polygons as well as in bogs. If permafrost thaws and the ice melts, these landscapes become vulnerable to collapse. This results in a shift to lake or wet fen ecosystems, which are a major source of methane.

Permafrost occurs across almost 65 per cent of Russia. Although relatively limited in the European part of Russia, it is extensive across Siberia. The Western Siberian Lowlands, which started developing 10,000–11,000 years ago, form the world's largest peatland region. Much of the area is affected by permafrost, forming complexes with uplifted peat plateaus and wetter fen areas. The uplift of permafrost is caused by large volumes of ground ice. When thaw occurs, the loss of ground ice causes abrupt thaw and ground subsidence. The rate of permafrost degradation in the western Russian Arctic is among the highest in the world. In response to increased temperature and precipitation since the mid-1970s, mean annual ground temperature has increased from 0.03 °C to 0.06 °C per year in the continuous zone. Signs of permafrost degradation are evident, including an increase in active layer thickness and a lowering of the permafrost table (up to 8 metres in the discontinuous zone). Permafrost is warming most rapidly in the tundra in western Russia (up to 0.06 °C per year) and most slowly in the northern taiga (up to 0.04 °C per year).

The landscape in north-eastern Siberia is characterised by Yedoma permafrost or ice complex deposits. These thick deposits were formed during previous ice ages and are rich in both ground ice and organic matter. Massive ice wedges make up much of the ground volume (often more than 50 per cent and up to 90 per cent) and the frozen soil around the ice is high in carbon. This combination of ice and carbon content makes the landscape extremely vulnerable to thaw, and high emissions of greenhouse gases have been measured from thaw lakes and eroding coastal permafrost.

Crossing the threshold *Future scenarios of carbon release*

Earth system models are tools used to quantify the response of the Earth's climate system to greenhouse gas emissions. The models also provide a "budget" of how much carbon can be emitted into the atmosphere before a particular warming threshold is crossed. Understanding how the Earth's climate will change under different scenarios (or emission levels) is vital to setting greenhouse gas emission targets to limit the global temperature increase to 1.5 °C, as recommended in the Paris Agreement.

The Earth's climate system is too complex to make projections with a high degree of certainty. Earth system models do not include all sources of carbon released to the atmosphere, and calculating the emissions from natural sources, such as terrestrial permafrost, has a much higher degree of uncertainty than from anthropogenic sources, such as industrial activity. As a result, IPCC did not include emissions from permafrost until its Sixth Assessment Report. Contributions from subsea permafrost are even more difficult to model due to the lack of critical data.


Net carbon flux is the difference between the carbon taken up by plants through photosynthesis and the carbon released into the atmosphere from sources such as respiration and land-use change. As permafrost thaws, more stored carbon becomes available for release into the atmosphere. Although there will also be greater carbon uptake by plants as the climate warms, modelling shows that more carbon will be released than can be used in primary production. The greatest carbon emissions are expected to occur under Representative Concentration Pathway 8.5, the highest emissions scenario up to 2100. An intermediate emissions as the increased emissions from permafrost thaw is compensated by an increase in uptake by vegetation.

According to the IPCC Sixth Assessment Report, for every degree of global warming, anywhere between 3.1 and 41 billion tons of carbon dioxide will be emitted from terrestrial permafrost by the year 2100. Although a great deal of uncertainty surrounds the scale of emissions, it is agreed that as more atmospheric warming occurs, more carbon will be emitted from thawing permafrost, contributing to even more global warming. Missing from these estimates are emissions from the abrupt thaw of permafrost, something which scientists are only just beginning to understand. Some estimates suggest that for every degree of warming, abrupt thaw will release almost twice as many emissions as from gradual thaw.

In current models, subsea permafrost thaw is similar across the different climate scenarios. Up to around the year 2100, all scenarios show permafrost thawing at about four times the rate prior to anthropogenic climate warming. After 2100, the loss of subsea permafrost continues at relatively similar rates. In the high emission scenario, however, subsea permafrost starts to thaw dramatically, about 15 to 18 times faster than before anthropogenic climate warming. The rapid thaw of subsea permafrost is tied to the loss of Arctic sea ice, which will cause the seabed on the Arctic shelves to warm up faster thus accelerating permafrost thaw. The continued reduction of sea ice is very likely. even under optimistic carbon emission reduction scenarios. Since the development of sea ice drives the formation of cold brine, which cools the seabed, the long-term effect of sea ice reduction will be warming of the permafrost below the shallow Arctic shelf areas.

Based on the limited knowledge of the quantity and quality of organic matter in subsea permafrost, models suggest that at the current rate of climate warming, methaneproducing microbes will decompose more than 400 million tons of carbon annually after the year 2100. Most of this carbon decomposition produces methane below the sea floor, over 200 million tons per year under the high emission scenario. However, it is difficult to predict how much methane will ultimately reach the atmosphere. In the sediments and water column above the thawing permafrost, methanotrophic microbes can oxidise the upward-migrating methane under both aerobic and anaerobic conditions, turning it into carbon dioxide before it reaches the sea floor and the atmosphere. The benthic methane sink is an extremely efficient microbial filter for upward-migrating methane and usually consumes a huge fraction of the methane flux. The current estimated global total of methane emissions from the ocean to the atmosphere is approximately 4–10 million tons per year. Even under the most extreme climate change scenarios, the maximum amount of methane released from thawing subsea permafrost, would only be about double what they are today.

Even if society reaches its carbon emission targets, emissions from permafrost may persist, which in turn could lead to overshooting those same targets. As permafrost thaws, it adds more greenhouse gases to the atmosphere, which drives further warming and risks exceeding warming predictions. In this scenario, permafrost carbon would be irreversibly lost because most types of permafrost do not recover quickly or easily. Therefore, permafrost thaw could also trigger a carbon loss that continues for centuries. These future emissions from permafrost need to be accounted for in modelling.



SOURCE: RIDOLFI, ARNDT ET AL (IN PREP). ADAPTED BY LEVI WESTERVELD | GRID-ARENDAL (2023)



© Olga Borjon-Privé (Oluko)

PORTRAIT ASSOCIATE PROFESSOR OF GEOGRAPHY AND INTERNATIONAL AFFAIRS AT THE GEORGE WASHINGTON UNIVERSITY, UNITED STATES **Dmitry Streletskiy**

I grew up in Moscow right next to a park, where I mostly spent my childhood outside climbing trees. Moscow winters were brutally cold with plentiful snow, so I did a lot of skiing and other activities related to cold climates. I think that was part of why I started to be interested in cold climate regions. The Arctic environment, especially the tundra, fascinated me with its low sky and pastel colours.

I've always enjoyed travelling. In high school, I realised that if you did geography, you could get paid to travel. That was my idea of a geographer and that's why I wanted to become one. These days, admittedly, I spend most of my time sitting in the office, but my favourite aspect of work is still being out in the field.

When I was a student, I was part of expeditions that focused on glaciers. To be honest, drilling in glaciers was a little bit too much for me. Permafrost research was more user-friendly since a lot of investigations are constrained by logistics and access to infrastructure. The research happens closer to people and doesn't involve sitting on glaciers in the middle of nowhere at high elevations. Permafrost research is also related to a lot of different job opportunities. You can do all sorts of things.

I realised that permafrost is essential for the people and communities living on it. I started looking at the connection between infrastructure and nature, and what the impacts will be once permafrost starts changing. After completing a PhD in climatology at the University of Delaware, I left for George Washington University, where I have been for more than 10 years researching and teaching classes in Arctic physical geography, natural hazards, and climate change. Lecturing students really keeps me motivated because they realise the problems we face, and they are anxious. They really want to do something about it and try to be involved. This helps keep my interest in the topic of permafrost. I think permafrost is an understudied topic and there is no shortage of new permafrost topics to research. We need geographers, biologists, social scientists, economists, engineers, material scientists – you name it.


Some of the things we permafrost scientists communicate do materialise and become policies, and I see some efforts going in the right direction. But sometimes it can get a bit frustrating when you have a good knowledge and understanding of something, but it doesn't necessarily ring a bell with others. I find myself knocking on a door, but there is no one to open it. I think permafrost doesn't get as much attention because it's not like an earthquake or a volcanic eruption with a sudden big event – it's gradual. But then an event like the oil spill in Norilsk happened, and that was like an earthquake. Suddenly I was getting calls and emails from all sorts of people: from municipalities, officers in the government, investors that sit somewhere in Australia or in London. Everyone suddenly wanted to know what permafrost is and what is happening to it.

It's very exciting to be part of a small community and to do something that not many people are doing. I'm at the forefront of a topic, asking new questions and trying to answer them, and then trying to communicate those answers to various groups. I also think it is very exciting to meet various stakeholders, not just scientists at conferences. Talking with these people helps me get a better sense of what's happening on the ground, and I love that.



Moving Grounds

Permafrost Changes




Frost and flora *The role of vegetation in permafrost landscapes*


Vegetation plays an important and complex role in permafrost processes. The seasonal thawing of the active layer provides a medium in which plants can grow. In turn, the presence of vegetation helps insulate permafrost, retain moisture in the surface soil, and moderate the effects of changes in air temperature. The influence of vegetation on permafrost is also closely linked with snow accumulation.

Where vegetation is thick, the shade it provides can cool the soil surface. Dense tree canopies contribute further to cooling by limiting the amount of snow reaching the ground. This affects the distribution of permafrost in the discontinuous zone. Conversely, where the canopy is more open, snow accumulates more readily, providing insulation from cold air temperatures, such as along the treeline. Areas with thick layers of organic matter also offer more insulation from changing air temperatures.

The Arctic tundra is dominated by low-growing vegetation such as grasses, sedges, rushes, small flowering plants, mosses, lichens, and shrubs. However, the warming of the Arctic is rapidly changing Arctic vegetation, spreading shrubs further north. Shrubs are more effective in trapping snow than lowgrowing tundra vegetation, mitigating the impacts of air temperature on permafrost. As such, areas with taller shrubs tend to have warmer surface temperatures than those with low-growing tundra vegetation. Some changes in vegetation are related to permafrost thaw, in addition to warmer and longer summers, increased precipitation, and increasing atmospheric carbon dioxide concentrations. Gradual permafrost thaw may stimulate vegetation growth through the release of nutrients as soil carbon decomposes. But when thaw is abrupt, plant communities may experience high mortality and changes in composition as the soil surface subsides or collapses. As permafrost thaws, changes also occur in the thickness of the active layer, local topography, and soil moisture content. The relationship between the change in tundra vegetation and permafrost, however, is still not well understood.

As Arctic summers become warmer and precipitation patterns change, climate-related disruptions such as extreme weather and vegetation fires are also becoming more common and are expected to increase in the coming decades. More frequent hot summers sometimes allow fires to continue smouldering over the winter and then flare up again in the subsequent spring. The loss of vegetation and the insulating surface organic layer to fire can exacerbate permafrost warming. Studies have found the mean annual ground surface temperature is 1-7 °C higher at sites that have burned.





Fire on Ice Peat, permafrost, and fire

Large areas of Arctic permafrost are covered by peatlands: wetland ecosystems that have a layer of partially decomposed organic matter (peat) at the surface. They are most widespread in the southern Arctic, especially in Canada's Hudson Bay lowlands and the Mackenzie River Valley as well as across Siberia, becoming less common further north where lower precipitation, low temperatures, and short growing seasons limit their formation. Peatlands play a crucial role in regulating the Earth's climate, absorbing and storing carbon dioxide from the atmosphere, helping to mitigate the effects of global warming.

Of the estimated 5 million km² of peatlands globally, most are in the subarctic and boreal zones. Northern peatlands cover close to 4 million km², almost half of which – about 1.7 million km² – are underlain by permafrost. Peatlands are significant carbon storehouses, containing almost one-third of the world's soil carbon, an estimated 450 to 650 billion tons. Most of this carbon, 415 billion tons, is stored in northern peatlands.


Peatlands play an important role in maintaining permafrost with the thick peat layer helping to protect it from temperature variations. In summer when peat is dry, it insulates the underlying permafrost from warm air. When peat is wet or frozen, it allows cold air to penetrate the soil, thus helping conserve permafrost where it might not otherwise exist. However, changes in temperature and precipitation are leading to a decrease in the extent of permafrost peatlands. The area of permafrost peatlands has already decreased from an estimated 2 million km² in pre-industrial times to 1.7 million km² today. It is projected that, under 2 °C of global warming, this area will decrease to 1 million km², and at 6 °C of warming, it is thought that almost all permafrost in these peatlands will have disappeared.

Peatlands in high northern latitudes, including permafrost peatlands, are vulnerable to changes in temperature and precipitation. As temperatures increase, it can dry out the overlying peat, making it more susceptible to fires. Not only do fires release vast amounts of carbon from the burning peat but they also destroy the insulating layers of vegetation, peat, and soil, increasing the rate of permafrost thaw. The combined effects of climate change and wildfires can turn permafrost peatlands from carbon sinks to carbon sources creating a positive feedback loop that exacerbates the effects of global warming. The impacts to permafrost are even more severe in the discontinuous zone.

Climate change is increasing the risk of catastrophic wildfire events in many regions of the world; even the Arctic is no exceptions. The summers of 2019 and 2020 saw unprecedented wildfires in the Arctic, the worst occurring in Siberia, an area with extensive permafrost peatlands. Higher winter and spring temperatures led to fires occurring as early as May and burning for longer periods of time, and further north than usual. In peat deposits, fires can also continue to smoulder undetected beneath the surface, bursting back into life weeks or even months later when the conditions are favourable. Almost 140,000 km² burned in the 2020 Siberian fires, most occurring in the permafrost zones. It is estimated that the fires in these carbon-rich peatlands released 244 million tons of carbon dioxide, 35 per cent more than the previous record set in the 2019 fires.

State of matter *Water, snow, and permafrost*

Whether as precipitation, in waterbodies, or as ice in the ground, water influences the state of permafrost, how quickly it thaws, and with what consequences. Snow, for example, can have either a cooling or a warming effect on the underlying permafrost depending on snow depth as well as timing of accumulation and duration over the cold



SOURCE: MCCRYSTALL ET AL. (2021). BY LEVI WESTERVELD | GRID-ARENDAL (2023)


season. Where snow is thicker, the ground surface temperature is also higher because cold winter air cannot fully penetrate into the ground to cool it down. Periods of permafrost warming in Alaska (late 1980s to early 1990s) and Norway (1999–2009), for example, have been linked to deeper snow cover as well as increased air temperature. A thin snow cover cools the ground due to the high albedo, which leads to reflection of long wave radiation from the white snow surface.

The timing of snow accumulation and melt is also important. When winter snow arrives late, the active layer may refreeze and the permafrost below may cool more quickly in response to decreasing air temperature, whereas earlier snow accumulation may delay these processes. In the spring, the effect is reversed: an earlier melt may lead to earlier thawing of the active layer, while thawing is delayed in areas where snow cover lingers longer. Snow-covered ground cannot thaw. The active layer can only start developing once the snow has melted.

All climate change scenarios indicate longer and wetter summers in the Arctic, with higher precipitation under the high emissions scenarios. Greater summer rainfall can lead to increased permafrost thaw because of the high heat capacity of water. After very wet summers, thaw depths may take longer to return to "normal" values. Higher rainfalls have an even greater effect in disturbed areas, for example, where the ground has subsided due to permafrost thaw. Surface water ponding is not unusual in these sites and may lead to further thawing. Increasing rainfall may also lead to increasing active layer depth in wetlands. Although the vegetation provides some protection, water tends to accumulate in these low-lying areas and, as it warms, can heat the permafrost beneath it.

Permafrost has a big impact on the hydrological functioning of the soil, which in turn impacts Arctic hydrology in general. Earlier snowmelt and increased river flows in the Arctic are leading to earlier and longer periods of inundation in floodplains resulting in greater warming of the underlying permafrost. Later freeze-up of rivers in permafrost-dominated catchments means that more and warmer water runs through permafrost terrain later into fall, which can also lead to a later build-up of sea ice along the Arctic coasts.

Thaw-induced changes affect all Arctic rivers, but smaller river systems are warming the fastest. The northernmost permafrost regions have thousands of small river reaches or segments, most of which are underlain by continuous permafrost and soils with high carbon content. In many areas, the landscape is strongly influenced by ice within the permafrost. Ice-wedge polygon terrain is an important example of this. In flat coastal systems, these ice-wedge polygons serve as small water retention basins. Warming will cause the ridges around the polygon to thaw and the ice wedges to melt, which will invert the system. This will lead to more efficient draining of the landscape and further thaw. This may, in turn, result in greater leaching and export of nutrients such as carbonate, nitrogen, and phosphorus into rivers, and the Arctic Ocean, and carbon into the atmosphere.




Fine sediments (bright colours) and dissolved organic matter (dark colours) are visible as cloudy water at the mouth of the delta. Increased dark organic matter absorbs more heat, warming the surface waters of the ocean. More suspended material increases turbidity leading to less light penetration, affecting primary production.

Newly formed ground comprises fertile wetlands and numerous small lakes. This mosaic of ecosystems supports rich flora and fauna, including migratory bird populations and grazing habitat for mammals.

5 kilometres

In a delta, the main river branches off into smaller channels rather than remaining in a single outlet. As the meandering channels approach their end, their flow velocity is reduced and the suspended material they carry sinks to the ground and settles. This builds new ground.

> High moisture content in sediments accounts for the ice-rich permafrost in Arctic riverbanks and deltas. Climate warming and associated permafrost thaw leads to the melting of ground ice, resulting in riverbank erosion and the formation of numerous lakes.





The rivers run through it Arctic rivers, deltas and hydrology

Rivers and river deltas play a vital role in Arctic ecosystems. While the Arctic Ocean accounts for about 1 per cent of the Earth's ocean volume, it receives more than 10 per cent of all global river discharge. In addition to its numerous small river systems, the Arctic is home to some of the world's longest rivers, including the Yenisei, Ob, and Kolyma, and three of the world's largest river deltas: the Lena, Mackenzie, and Yukon.

River deltas are fan-shaped landforms created by the deposition of sediment at the mouth of a river. Typically featuring shallow, interconnected channels, lakes, and wetlands, deltas support diverse ecosystems, flora, and fauna, including migratory bird populations. Deltas form a crucial link between the land and the ocean and influence the transport of fresh water, sediments, carbon, and other materials.

Permafrost underlies a substantial part of the catchment of all Arctic rivers, both big and small. The presence of permafrost largely controls the hydrology of the area by restricting the flow of water. The presence of ice in deltas and riverbanks affects erosion and sedimentation processes, as well as the seasonal flow of water into the Arctic Ocean. Frozen riverbanks hinder the transport of sediments because ice protects the ground from erosion. With increased permafrost thaw, the terrain may become unstable and lead to the erosion of riverbanks and formation of thermokarst lakes. In northwestern Canada, for example, increasing temperatures triggered riverbank erosion and an acceleration of retrogressive thaw slumps in the Mackenzie Delta, leading to rapid remobilisation of large masses of sediment and carbon from ice-rich permafrost terrain into adjacent rivers.

As permafrost thaw intensifies, it will become a greater source of organic carbon to the ocean through enhanced riverine discharge and the mobilisation of eroded materials. This will be augmented by increased precipitation in the Arctic, which will lead to more riverine transport in summer and early autumn when permafrost thaw is also most intense. These changes in hydrology will have a profound impact on the terrestrial ecosystems of deltas, as well as the freshwater balance, and biochemical fluxes in the ocean.

The Arctic Ocean is currently a large carbon sink, meaning it takes up carbon from the atmosphere and stores it in the water column or buries it in the sea floor sediments. Greater inputs of organic carbon to the ocean can change this, leading to an increased carbon flux or exchange from the ocean to the atmosphere.

When carbon dioxide reacts with water, carbonic acid is produced, which can lower the pH of the water, leading to ocean acidification. Not only does this affect the functioning of marine ecosystems, but it also reduces the ocean's ability to absorb more CO_2 . Since CO_2 dissolves better in cold water, the Arctic Ocean is more sensitive to ocean acidification. The increased influx of organic carbon from river run-off, resulting from permafrost thaw, will further reduce the Arctic Ocean's capacity to absorb carbon.

Along the edge of the world Arctic coastal classification

Arctic permafrost spans the coastlines of Alaska, Canada, Kalaallit Nunaat (Greenland), Norway, and Russia. Much of this permafrost coastline is characterised by the presence of ice in the ground and ice on the sea. As the Arctic warms, these coasts become vulnerable to erosion which changes coastal ecosystems, threatening human settlements and infrastructure. In recognition of this, a geomorphological classification scheme was developed for the more than 100,000 km of coastline bordering the Arctic Basin. It provides a baseline against which scientists can measure fluxes of sediments, carbon, nutrients, contaminants, and other parameters as these coastlines change over time.

Arctic coasts show considerable variation in their geomorphology, sediment properties, and rates of erosion. Approximately 35 per cent of the classified coasts (approximately 35,000 km) are lithified (rocky). These predominantly occur in the Canadian Archipelago, northern Kalaallit Nunaat, north-western Russia (Kola Peninsula), and the islands in the Barents Sea (northern Svalbard, Franz Josef Land, and Novaya Zemlya). Lithified coastlines are characterised by low-lying rocky shores, fjords, and cliffs, and tend to contain little to no ground ice. These rocky shorelines are relatively stable with erosion occurring mostly by weathering and wave action but are also affected by freeze-thaw fracturing and direct ice action. They were mostly covered by continental ice sheets during the Last Glacial Maximum (LGM), and some are recently uncovered by slow glacial and rapid ice shelf retreat. Extensive tidewater ice fronts remain, shedding icebergs.


The greater portion of Arctic Basin coasts – 65 per cent or approximately 66,000 km – are unlithified or sedimentary. Unlithified coasts dominate the shores of the north-western Canadian Arctic, Alaska, and most of Siberia: areas that were not covered by continental ice sheets during LGM. Some unlithified coasts are characterised by large volumes of ground ice, the amount of which is determined by the glacial history of an area. Unlithified coasts in Alaska and eastern Siberia, for example, are characterised by late Pleistocene-era, icerich Yedoma permafrost. Common features include icerich permafrost bluffs, beach ridges, barrier islands, spits, deltas, and retrogressive thaw slumps. Unlithified coasts in other parts of the Arctic, such as parts of the Canadian Archipelago, Kalaallit Nunaat, and Svalbard, contain less ground ice but may be affected by anchor ice, ice ride-up, and pile-up ridging. Boulder barricades and boulderflats demonstrate the importance of sea-ice rafting and tidal action but are largely unaffected by permafrost.

The amount and distribution of ground ice within permafrost affects thermal and mechanical erosion, as well as other geomorphological processes. As air temperature increases, soil is warmed from both the surface (top-down) and the coastal bluff or cliff face (inward from the coast) leading to rapid permafrost thaw and ground-ice melt. Unlithified coasts, especially those in ice-rich areas, are vulnerable to climate change. At some locations, yearly erosion rates can exceed 20 metres per year.

The presence of sea ice also has a great influence on permafrost coasts. As sea ice declines because of

climate warming, ice-free seasons become longer, leading to increased exposure of vulnerable shorelines to more wave action. Reduced sea ice means there is a longer fetch – the distance across open water – leading to higher wave energy. Since the beginning of satellite measurements in 1979, Arctic sea ice has been declining by 13 per cent per decade, with a loss of about 80,600 km² per year, with the greatest declines in the last 15 years. As the open-water season becomes longer, the coastlines will be exposed to more storms later into autumn, impacting both coastal ecosystems and coastal communities and infrastructure.

While erosion is a concern along many permafrost sedimentary coasts, less research has been conducted on emerging polar coasts -- areas where the land is rising or expanding, like deltas and other sedimentaccumulating landforms, leading to the development of new permafrost. Glacial isostatic adjustment is caused by the Earth's crust rebounding after being pressed down when covered by thick ice caps during the last ice age. The east coast of Canada's Hudson Bay in northern Quebec, for example, is rising at approximately 13 millimetres per year. Monitoring conducted here since 2005 shows that permafrost has been forming and deepening in land that is now exposed to a cold climate. Similar aggradation of permafrost has been documented at other sites around Hudson Bay and in the Canadian Arctic Archipelago.



Wear and tear Erosion of Arctic permafrost coasts

Like coastlines everywhere, the main drivers of coastal erosion in the Arctic are increasing air temperature, sea level rise, and wave action. Most coasts around the world experience physical erosion (through wave action). The Arctic coasts, however, are also affected by thermal erosion (through increasing air and water temperatures). The presence of ground ice is the major difference between Arctic coasts and all other coastlines and makes them especially vulnerable to erosion.

Unlike most other coasts in the world, **air temperature** plays a significant role in coastal change in the Arctic. Increasing temperatures affect not only the ground temperature and active layer thickness, but also sea temperature, sea ice extent, and the duration of sea ice cover. Ice-rich permafrost bluffs are particularly


impacted by **thermal denudation**, the process whereby increasing air temperature and thawing permafrost cause parts of the bluff face to loosen and collapse. As coastal permafrost degrades, it may also be subject to saltwater intrusion from wave action, further increasing erosion.

Changing **sea levels**, both rising and falling, affect shorelines. Rising sea levels can exacerbate thermal abrasion as waves reach higher on the bluff face. Even temporary changes in sea levels, such as during storms, can cause changes in the shoreline and increase thermal abrasion.

Wave action depends on wind speed and direction, fetch (the distance of open water over which wind blows), and nearshore bathymetry (the measurement

of underwater depth). In the Arctic, the mean annual fetch is strongly affected by the duration of the openwater season: the part of the year without sea ice. The longer the open-water season, the more erosion will occur. Some of the strongest storms occur in September and October, a time when sea ice is at its minimum extent and both the permafrost active layer and surface temperatures are at a maximum, making permafrost coasts more vulnerable to erosion.

Thermal abrasion is a mechanism whereby the water temperature together with wave energy erode ice-rich permafrost coasts. The impact of thermal abrasion on ice-rich coasts is three to four times stronger than wave action alone. In some areas, thermal denudation and thermal abrasion act together to attack the bluff top and base, respectively.



Another form of erosion known as **block failure** occurs when waves undercut sections of a cliff. The niches created by thermal abrasion can extend several metres into the cliff's base, particularly in ice-rich horizons, causing large blocks of ice-bonded sediment to collapse. In a matter of days to months, these blocks are eroded by wave action. Some of the highest erosion rates in the Arctic are the result of block failure, reaching 48.8 metres on the Beaufort Sea coast in Alaska in 2008.


Eroding coastlines can have numerous impacts on the natural environment and human settlements. As permafrost degrades, it can release large amounts of organic carbon, nutrients, and contaminants into both the aquatic environment and the atmosphere. It is estimated that coastal erosion contributes 14 million tons of particulate organic carbon annually to the Arctic Ocean, more than the input from rivers. Coastal erosion is also responsible for 1.6 million tons of nitrogen entering the Arctic Ocean annually. Although this large influx of nutrients may increase primary production in the nearshore, the overall impact on coastal ecosystems remains poorly understood.

While some carbon from eroding coastlines will be taken up in organisms and marine sediments, it will also be released to the atmosphere. As organic carbon currently stored in permafrost is exposed by erosion, it starts to thaw and decompose, releasing carbon dioxide and methane into the atmosphere. Studies show that the emission of greenhouse gases from eroding coastal bluffs begins as soon as the carbon is exposed to seawater.

Coastal erosion can directly threaten human settlements. The community of Tuktoyaktuk on the Canadian Beaufort Sea coast, for example, has suffered extensive damage because of coastal erosion. Although many efforts have been made to protect the community from the sea, the coast has eroded to the point that numerous buildings need to be relocated further inland or they will be washed away. Coastal erosion is so severe in some locations that whole communities need relocating (e.g., Shishmaref and Kivalina in Alaska). This is not an easy proposition since the costs of relocation are high, and the cultural identities of these communities are intimately tied to their current location.

Coastal erosion is also affecting cultural heritage in the Arctic. Numerous archaeological and cultural sites across Alaska and Canada have been destroyed. Even in Kalaallit Nunaat (Greenland) and Svalbard, which are less susceptible to erosion because of their generally rocky shoreline, cultural artefacts along their coasts are at risk as the climate warms.

Drew Point, Alaska (image from 23 July 2019). In the span of 10 days, coastal erosion led to the coastline retreating by several metres.







Eating into the landscape Retrogressive thaw slumps

One of the features of thawing permafrost is the appearance of retrogressive thaw slumps. These are large, curved depressions along coastlines, riverbanks, lakeshores, and hill slopes. These slumps occur when massive ice bodies in permafrost are exposed to warm air, making them vulnerable to rapid thaw and erosion. As the ice melts, a slurry of water and sediment moves downslope and the headwall retreats – or retrogresses – inland. The thawing permafrost has the potential to trigger the release of greenhouse gases and contaminants that have been stored in the permafrost, sometimes for thousands of years. Retrogressive thaw slumps are common in the icerich permafrost of the western Canadian Arctic, the Yedoma of eastern Siberia, and the central Yakutia region of Russia where they are referred to as thermo-terraces or thermocirgues. They can be several hundreds of metres wide and can retrogress several hundreds of metres inland.

As the climate warms, retrogressive thaw slump activity is increasing. On the Taimyr Peninsula, located between the Kara and Laptev Seas in Siberia, retrogressive thaw slumping has increased 43-fold between 2010 and 2021. In this same period, carbon mobilisation increased 28-fold. This increase coincides with an extreme heat wave in northern Siberia in the summer of 2020, in which the region experienced record-breaking temperatures (38 °C). As the climate continues to warm, heatwaves in the Arctic can be expected to occur more often, exacerbating the impacts on permafrost.

Despite the considerable risks associated with increasing retrogressive thaw slump activity, it is still poorly accounted for in the Arctic carbon cycle.





PORTRAIT INUVIALUIT SETTLEMENT REGION COMMUNITY-BASED MONITORING PROGRAM, INUVIK, NWT Angus Alunik

My name is Angus Alunik, and I was born by a small creek they call "Awa" in the Mackenzie Delta in 1952. Soon after my birth, my mother passed away, so I was adopted a few days later by my adoptive parents, Ishmael and Ruth. Luckily, they maintained a good relationship with my biological father, so I was able to learn about my roots.

We lived in the bush most of the time, until 1969 or 1970 when we moved to Stringer Hall. In May, before the ice would melt, my father would take me to the bushcamp for the spring. I'd spend a lot of time hunting with him because he wanted me to learn the ropes. My brother mostly stayed in school. While the ice was still there, we would often leave the cabin and set up a tent for the three of us. We'd walk around, leaving the dogs behind, setting up muskrat traps and sometimes even catch beavers. Years ago, the ice would stay there until June.

I remember, years ago, my dad would make a small pit to keep stuff frozen. You dig a bit until you hit permafrost and you put fish and meat in there. A lot of people built ice houses. They would put logs and then mud around it and then make a little door. It used to be good then, but years later I noticed that you had to dig deeper and deeper to hit permafrost. It's really thawed.

I also see a lot of erosion, not only at home where my camp is, but all along the foothills from Inuvik right down to Reindeer Station. After I moved from Edmonton, I built a house not too far from where my dad's log cabin was. When I first built my house, there was a lake behind me, with another lake towards the Mackenzie River on the left side. At the time, it seemed okay as we didn't see any signs of erosion. But after a decade or so, I noticed that we started to lose riverbank. We had gone to bed for the night, I had the generator running about 100 metres from the bank. This was in the summer of 2000. The next morning, while we were having breakfast, my wife said, "I thought your generator? I walked over and saw we'd lost a part of the riverbank the size of the house. It just disappeared.

Ever since then we've been losing riverbank all the time. Every few years, I'd have to borrow a winch to pull the house back because the riverbank was so close. Then, suddenly, the lake behind us was gone, drained into the channel. From there, it got worse. About five years ago, I lost my whole house. The cabin that I built went over the bank and I lost everything in it.


Some people were already having meetings about this, but I wasn't really involved in any of them. I just started going to meetings recently, but how can you fix Mother Nature? I don't think you can. I think it's going to keep coming. So, for me, I just live with it. I mean, why get mad about it? You can have all the meetings you want and everything and try to fix it. But I don't think we can. I think, now that it's started, we won't ever stop it.

Losing ground Projected rates of Arctic coastal erosion

Arctic coasts are subject to erosion by mechanical forces (e.g., wave action) and thermal forces (e.g., increasing air temperatures). These forces work together to thaw permafrost, melt ground ice, and wear away material at the coast. As the Arctic continues to warm, permafrost coasts thaw, potentially releasing greenhouse gases into the atmosphere and further exacerbating the drivers of coastal erosion.

The rate of coastal erosion has been increasing in recent decades. Between 1850 and 1950, the mean rate of coastal erosion was estimated at 0.9 metres per year, considering only the erosive segments, or 0.5 metres per year of the 400,000 km of coasts facing the Arctic Basin. By the end of this century, it is estimated

Modelled Arctic coastal erosion rates Mean values in metres per year. 4 m v




SOURCE: NIELSEN ET AL. (2022). ADAPTED BY LEVI WESTERVELD | GRID-ARENDAL (2023)


that the mean rate of coastal erosion for both rocky and sedimentary coasts will increase to between 2 metres per year under an intermediate climate change scenario, and 2.6 metres per year under a high climate change scenario. Current modelling shows that coastal erosion in the Arctic will likely exceed its historical range by around 2050, and possibly even earlier.

Coastal erosion releases large amounts of organic carbon into the ocean. The greatest current and projected losses of organic carbon – about 75 per cent – are in the Laptev and East Siberian Seas of Russia. Coastal erosion in this region is very high, up to 20 metres per year. This region is also dominated by icerich Yedoma deposits and high cliffs with high organic carbon content. The next highest losses are along the Beaufort Sea coast of Canada and Alaska, accounting for about 12 per cent of the Arctic total. The remainder of organic carbon loss in the Arctic comes from the other marginal Arctic seas.

There are numerous uncertainties surrounding future coastal erosion but even in low emissions scenarios, erosion rates are projected to increase until at least the middle of the century. In the higher greenhouse gas emissions scenarios, however, coastal erosion would continue to increase beyond 2050, releasing between 2.3 million tons (intermediate emissions scenario) and 4.2 million tons (high emissions scenario) of organic carbon annually. Starting from the year 1850 up to 2100, it has been estimated that about 2 trillion tons of carbon will be lost from permafrost as a result of coastal erosion. This amounts to 17–26 per cent of the overall carbon losses from thawing permafrost across the Arctic. But there are considerable uncertainties in these estimates. The actual losses will depend on the degree of warming that eventually occurs as well as the impacts of that warming on the ecological and geomorphological processes associated with coastal erosion.

Coastal erosion is important to marine primary production in the Arctic, as it releases and transports organic carbon into the aquatic environment. The fate of the increased organic carbon in the Arctic is still relatively unknown. It may settle into marine sediments where decomposition will be very slow (over millennia), or it may be partially decomposed before entering the ocean, releasing greenhouse gases into the atmosphere. It is also possible that it will degrade in the water column or be transported further away. Degradation in the water column happens through a process of mineralisation, which changes the ocean alkalinity and pH and increases dissolved inorganic carbon and nutrient supplies, all of which impacts primary production and air-sea carbon fluxes.






Beneath the waves *Changes in subsea permafrost*

Permafrost not only occurs on land but under the ocean floor too. Subsea permafrost is relict terrestrial permafrost that formed during the Last Glacial Maximum (LGM) when the global sea level was more than 100 metres lower than present day, exposing the present shelf areas. As the ice sheets retreated and the sea level rose, the permafrost that had formed on land was submerged under the ocean. Like terrestrial permafrost, subsea permafrost is composed of frozen sediment and rock, and may or may not contain ice. There is only about a fifth as much permafrost under the ocean as there is on land in the northern hemisphere. Most subsea permafrost occurs in northern Siberia, with some extending to Alaska and north-western Canada, land areas that were exposed during much of the last glacial period. The thickness

Bathymetric surveys show that seafloor depth on the Canadian Beaufort Sea Shelf increased up to 28 metres between 2010 and 2019.



SOURCE: PAULL ET AL. (2022). BY LEVI WESTERVELD | GRID-ARENDAL (2023)

of subsea permafrost varies across the Arctic. In most locations, it is less than 200 metres thick, but it can reach greater depths. Modelling suggests that permafrost in the Laptev, Kara, and East Siberian Seas may be over 700 metres thick in places.

There are numerous landforms that may be associated with subsea permafrost. **Pingo-like features** are found in some shelf areas of the Arctic Ocean, are similar in size and form to terrestrial pingos, and can contain ground ice. They are found in some shelf areas of the Arctic Ocean. **Mud volcanoes** are formed by the movement of sediments, fluids, and gases up to the sea floor. They can reach 40 metres in height and more than 500 metres in diameter. The unique environment created by mud volcanoes supports a rare community of tube worms in the Beaufort Sea. **Pockmarks** may be observed on the sea floor in areas where gases were recently released.

High-resolution bathymetric surveys in the Canadian Beaufort Sea have found large, newly created depressions on the sea floor, about 180 kilometres offshore and in water 120–150 metres deep. The largest depression is 29 metres deep, 225 metres long, and 95 metres wide. It is believed that these depressions are the result of permafrost thawing – similar to slumps and craters forming on land. While climate change is a driver of permafrost thawing on land, the limited data on sea floor temperature in this region do not show warming. Instead, these changes are thought to be the result of heat found in slow moving groundwater below the permafrost.

Current estimates suggest that approximately 2,800 billion tons of organic carbon has accumulated in

subsea permafrost over the last 450,000 years. This is more than twice the amount of organic carbon on land (estimated at 1,460–1,600 billion tons) and more than four times the carbon in our atmosphere. It is not known, however, whether it will be decomposed by microbes easily or quickly enough to have a correspondingly large climate effect.

Within and below subsea permafrost are gas hydrates, a solid form of gas and water. Gas hydrate molecules are made up of gas molecules, usually methane, within a cage of water molecules. The methane may originate from either deep deposits of oil and other hydrocarbons or from ancient microbial activity. They are only stable under high pressure, such as below the ocean floor and at very low temperatures.

Although there is interest in gas hydrates as a future energy source, they present a potentially significant source of greenhouse gases. If methane hydrates destabilise, they will release more than 160 times their volume as methane, one of the strongest greenhouse gases. It is not clear how much gas hydrate there is in subsea permafrost, but a conservative estimate is 20 billion tons of carbon. Better estimates are also needed of how much methane occurs as free gas in sediments.

Permafrost acts as a barrier to prevent gases from reaching the surface. If gases are released from, for example, oil deposits deep below the ocean floor, they become trapped by the subsea permafrost as they migrate upward through the sediment. But as subsea permafrost thaws, it starts to release the trapped carbon dioxide and methane which may reach the atmosphere, contributing to global warming.



Arctic Ripples

Impacts of Permafrost

Feeling the heat Permafrost thaw impacts on infrastructure

Permafrost has always presented an engineering challenge for infrastructure. Human settlements, industrial facilities, and networks for transportation, communication, and energy all need a stable surface on which to function. When infrastructure lies on top of permafrost, construction must accommodate the annual thawing of the active layer to keep structures secure while not disturbing the thermal balance of the frozen ground. Even just erecting a structure directly on permafrost can generate localised heat, increasing the thaw while decreasing the ground's bearing capacity. As the climate warms and permafrost softens, the impacts on infrastructure may pose significant risks to the safety and security of people who live and work in the Arctic.


The scale of this problem is immense: 70 per cent of all Arctic residential, industrial, and transport infrastructure is in areas where near-surface permafrost thaw is likely by 2060. Currently, almost 5 million people live in 1,150 settlements located on permafrost, the majority concentrated in just over 500 settlements located in zones of sporadic permafrost. By 2050, 42 per cent of the 1,150 permafrost settlements are projected to become permafrost-free due to thawing, affecting an estimated 3.3 million people.

Infrastructure is most prone to damage in areas with high ground-ice content and thick deposits of frost-susceptible sediments as well as in areas of "warm permafrost" where permafrost thaw is most pronounced and where most of the infrastructure is located. As permafrost thaws, it can become unstable, shift, and resettle unevenly. This can lead to both ground subsidence and decreased bearing capacity, which causes cracks, deformations, or the collapse of buildings, pipelines, and other infrastructure. To address this, most of the infrastructure on permafrost has been built with the goal of maintaining the frozen state of the ground to give it "freezing strength or bearing capacity". Other methods of building on permafrost include active cooling or modifying the ground properties prior to construction, such as by excavating thawing soil. Another option to actively cool the ground is the insertion of thermosyphons underground. Thermosyphons, like refrigerators, are closed heat exchangers based on gas-liquid flow circulation. This method, however, is expensive and thus not frequently used.

Some of the most striking examples of infrastructure damage from permafrost thaw have been to buildings in the Russian Arctic, where most of the infrastructure was built with the goal of keeping the permafrost beneath it in a frozen state. Housing blocks were typically built with a ventilated airspace (crawl space) beneath elevated buildings supported by concrete piles to prevent heat from the buildings reaching the ground. Unlike open crawl spaces where permafrost temperature is close to the mean annual air temperature, the air in poorly ventilated crawl spaces is warmer and heats the permafrost below. As permafrost warms across Arctic settlements, the impacts of poor design will worsen. In some Russian cities, such as Vorkuta in the Komi Republic, 80 per cent of all buildings now experience deformations in their building structure due to thawing permafrost, along with other factors such as poor construction

quality and poor maintenance. Many buildings have been abandoned or demolished because they are unsafe to live in or because the structural failures make it impossible to retain interior heat during the cold winters.

Linear infrastructure is also subject to damage from changes in permafrost. In summer, for example, roads can experience subsidence through the thawing of permafrost and loss of bearing strength on fine-grained sediments. In the winter, frost heaving may cause substantial dips, waves, cracks, and potholes in roadways, requiring constant maintenance and repaving. Groundwater can also seep into the embankments underneath roads. increasing the temperature of the soil and leading to further permafrost degradation. Pipelines, railways, and airport runways experience similar challenges. Railway lines often require costly upkeep. The necessity to maintain low gradients means that they are usually located in lowland areas where ground-ice content is often high.



Risky business I North American Arctic and Kalaallit Nunaat (Greenland)

Sparsely populated, the permafrost regions of North America and Kalaallit Nunaat (Greenland) account for just under 8 per cent of the entire Arctic permafrost population. These regions feature small individual houses and relatively lightweight administrative and industrial facilities. Buildings represent by far the most important infrastructure asset in these regions. Canada's road network represents a significant proportion of infrastructure value, but apart from the highways in the southern Northwest Territories and Yukon, and the road from Dawson to Inuvik and Tuktoyaktuk, the network is restricted to local roads in individual communities. Because of these restrictions, remote communities in Alaska, northern Canada, and Kalaallit Nunaat depend on ports and airports.


In Alaska, permafrost constitutes 80 per cent of the land surface. The greatest short-term risks to infrastructure over the coming decades are found in the discontinuous permafrost zone in the interior (central and western) parts of the state. The mean annual temperatures of Alaska's interior are currently around -2 °C. By 2050, they are projected to increase to +2 °C, indicating that the permafrost in this area will be actively degrading.

In Canada, permafrost underlies about 50 per cent of the ground surface, most of it in the Arctic. Many communities in the Canadian Arctic are located on emergent coasts. These uplifted former seabed deposits contain saline permafrost, which has reduced bearing capacity. Across the country, much of the residential, institutional, industrial, and transport infrastructure has been built on sensitive permafrost, i.e., areas with warmer, ice-rich permafrost. Amid climate change, the Northwest Territories, Nunatsiavut, Nunavik, and Nunavut face the widespread challenges of infrastructure degradation, increased maintenance requirements, and the dire need for adaptation.


In Kalaallit Nunaat, most of the ground that is not bedrock or covered by inland ice is exposed to permafrost thaw. While most houses and buildings in the country are built on bedrock, the roads and airports lie over permafrost, putting them more at risk than any other part of Kalaallit Nunaat's infrastructure.


Across these three regions where permafrost is already thawing, infrastructure needs constant maintenance and repair. This need is increasing, whether for airstrips or for highways such as the Alaska Highway in the Yukon, the Dempster Highway (Yukon, Northwest Territories), and the newly constructed Inuvik-to-Tuktoyaktuk Highway (Northwest Territories).


Permafrost degradation has a heavy cost: the estimated economic impacts of infrastructure damage across the circumpolar Arctic in these regions are projected to exceed USD 276 billion over the next 40 years under the higher-emission scenario. In the future, infrastructure is expected either to have a shorter lifespan or cost more to maintain. By 2060, the overall costs are estimated to increase by onethird for both Alaska and Canada (data unavailable for Kalaallit Nunaat), with pipelines having the highest estimated life cycle maintenance costs (60 per cent increase in cost by 2060), followed by roads, railways, airports (approximately 40 per cent each), and buildings (12 per cent).



NOTE: not all permafrost areas are covered in the model, including areas with substantial hazard potential. SOURCES: HJORT ET AL. (2018); KARJALAINEN ET AL. (2019); RAMAGE ET AL. (2021). BY LEVI WESTERVELD | GRID-ARENDAL (202







Risky business II The Russian Federation and Scandinavian Arctic

Russia will likely experience the most severe impacts of permafrost thaw. Not only does the nation have the highest number of settlements in the Arctic, but it also has the highest percentage located on permafrost (60 per cent of all Arctic settlements and 90 per cent of all Arctic inhabitants, respectively).


The Russian Arctic is also more urbanised and industrialised, encompassing the large cities of Norilsk, Vorkuta, and Yakutsk, which are characterised by large apartment blocks, centralised networks of heating and utilities, and large industrial facilities. High-risk areas in Russia include the north-western parts of the Ural Mountains as well as north-western and central Siberia.

Infrastructure failure is already very common across the Russian Arctic: by the early 2000s, an estimated 10 per cent of structures in the cities of Norilsk and Yakutsk and 80 per cent of structures in Vorkuta had suffered some form of deformation. Not all infrastructure damage can be directly blamed on permafrost thaw though. The collapse of the Soviet Union and worsening socioeconomic conditions in the years after led to the dismantling of permafrost monitoring infrastructure. Other factors include poor building design and construction, a lack of maintenance, and ageing infrastructure, as many buildings are being used beyond their anticipated lifespan. Consequences include the infiltration of hot water into permafrost from broken heating pipes, poor drainage, and ponding, as well as the low freeze-thaw resistance of the concrete foundation piles.

Urban infrastructure on permafrost presents numerous challenges because of the complexity of both the underlying soil conditions as well as the infrastructure above it (buildings, roads, powerlines, underground utilities, etc.). Russian city planners used one of two principles, either the passive method, i.e., maintaining permafrost in its frozen state, or accommodating thaw into the design. In some cities, however, violations of building codes took place and both principles were used in the same district. In Vorkuta, for example, plumbing and heating networks were laid underneath or close to buildings built using the passive method. These areas have experienced significant permafrost thaw and deformation of buildings.

Regardless of history, continuing permafrost thaw is detrimental to the bearing capacity of the underlying soil, which will keep causing serious problems for buildings and other infrastructure in the future. Already in north-western Siberia, permafrost thaw has reduced bearing capacity by an average of 17 per cent. By 2050, the infrastructure in the cities of Anadyr, Norilsk, Salekhard, and Yakutsk is expected to lose between 20 and 30 per cent of its bearing capacity under a high greenhouse gas emissions scenario.

In the Scandinavian countries of Finland, Norway, and Sweden, there has been far less damage to infrastructure lying on permafrost than in North America and the Russian Arctic, partly due to lower ground-ice content and less infrastructure being located on permafrost, but also as a result of higher investment and maintenance.





Coastal infrastructure in the Arctic is exposed to the impacts of both permafrost thaw and increased coastal erosion that accompanies sea ice loss. Mapping of Arctic coastal infrastructure and the type of permafrost it is built on is therefore important to understand potential future impacts and provide better risk assessments.

In 2021, satellite data was used for the first pan-Arctic assessment of coastal infrastructure located on continuous or discontinuous permafrost. The assessment defines coastal areas as being within 100 kilometres (km) of the coastline. High resolution data from the Copernicus Sentinel 1 and Sentinel 2 missions was combined with normalised difference vegetation index measures from Landsat imagery to identify how the human footprints in the Arctic have changed over the last two decades.


Of the approximately 6.2 million km² of permafrostaffected coasts in the Arctic (as defined above), about 0.02 per cent, or 1,243 km², are impacted by human infrastructure. Russia has by far the largest coastal infrastructure footprint on permafrost, followed by Canada and the United States. Most of this infrastructure is related to oil and gas development and mining activities. The infrastructure footprint has increased by 15 per cent since 2000, of which the majority has been in oil and gas development, highlighting the continued industrial development in coastal regions.

One of the continuously developing coastal industrial regions of the Russian Arctic is the Yamalo-Nenets Autonomous Region. As the largest natural gas province in the world, the region accounts for over 80 per cent of Russia's natural gas production and about 10 per cent of its oil. It is a region of huge importance to the Russian economy and, despite the challenges posed by building on permafrost, is home to several recently completed or ongoing industrial megaprojects. This includes the construction and expansion of the Bovanenkovo gas field and accompanying pipeline and rail infrastructure, as well as the Northern Latitudinal Railway project.


The Bovanenkovo gas field is one of the oldest gas fields in the Arctic, similar in age to Prudhoe Bay, Alaska, but where large-scale production only started in 2012. It is built on hilly terrain underlain by continuous ice-rich permafrost 140–230 metres thick and composed of wet silt and clay silt. Challenges due to permafrost thaw are well documented in this region. Included among these are landslides, which happen through a combination of rain and underlying permafrost thaw, and the melting of ground ice, resulting in water saturation and the detachment of the active layer downslope.

The Bovanenkovo gas field is also one of the most wellstudied regions in the Russian Arctic for greenhouse gas emissions from permafrost. The region is known to hold intra-permafrost, metastable (relict) gas hydrates (otherwise known as hydrate-bearing permafrost). Modelling suggests that emissions of methane (a powerful greenhouse gas) from this gas field could reach between 400 and 500 thousand cubic metres over a 30-year period in the absence of heat insulation in production wells.

Infrastructure



SPA



PORTRAIT PHD STUDENT, UNIVERSITY OF VIENNA, AUSTRIA Susanna Gartler

The reason why I, a central European, ended up living and working in the Arctic is because I focused on the term "subsistence" during my master studies. This led to my interest in Alaska and the Arctic in general, where subsistence is an important concept. I am living in Inuvik at the moment, at the intersection of the Inuvialuit Settlement Region and the Gwich'in First Nation traditional territories, where I've been conducting interviews with Indigenous land users and knowledge holders. I love the way it looks here in the beautiful Mackenzie River Delta, with all those lakes and rivers, and along the Arctic coast. Sometimes, when I go for a walk and I look around me during a sunset, I think: "Wow, this is the most beautiful thing I've ever seen."

And it's not just the land, it's the plants and animals too. But most importantly it's the people. I'm fascinated by the cultures, particularly the First Nation and Inuit cultures, and the way people are upholding and cherishing them. They're really beautiful cultures and societies to me: the philosophies, the arts, the handicrafts, the governance structures, the storytelling. It's also very interesting how the environment is storied here. I'm trying to see what the Indigenous stories and philosophies can tell us about how to live together well with all the other beings that surround us.

If you're only looking at the land in terms of its physical properties, you're missing a large part of why permafrost is important. Permafrost is literally the foundation of life here. It surprised me how diverse permafrost is. It is sort of a living, breathing, moving creature. During summertime when the ground broke up on the tundra, and you could see that not even half-a-metre underneath the earth there was a layer of pure ice. It blew my mind because I realised how vulnerable that is. When you realise that you and all the people around you are living on water, then you appreciate how important it is for that environment to stay cold enough.

I think part of what social science can contribute is to convey this message: that permafrost should not just be of interest because there is a lot of methane and carbon stored in it, but also because it's the home of people who've lived here for millenia. As an anthropologist, I'm trained to talk to people who come from different knowledge systems, and we have an ability to bridge different social and cultural worlds. The Nunataryuk project is very interdisciplinary. We've had workshops where community members were involved and provided input, while also working with natural, health, and engineering scientists, who have a lot of knowledge of our different field sites.

I'm very sceptical about bringing my own agenda and imposing that on a place that I, compared with the people who lived here all their lives, barely know anything about, no matter how much I study beforehand. However, I feel like as long as my research is guided primarily by local needs, it's a good thing. We are collaborating closely with different organisations and governments here, and we're making sure that our findings are not just taken away and published in scientific papers, but also made available to the people that we actually develop this knowledge with. Something I've heard a lot of times from Indigenous elders was that we have to tackle the big challenges of our days together, a notion that really motivates me.

Terra infirma II Reinforcing runways in Paulatuk

Paulatuk, a small Inuvialuit coastal settlement of about 265 people, is located on the shores of Darnley Bay in the Amundsen Gulf of the Beaufort Sea, Northwest Territories, Canada. Like many Arctic coastal communities, Paulatuk is facing impacts from coastal erosion and permafrost thaw. Located on a low-lying peninsula with an average elevation of just 3.2 metres above sea level, its western shore is experiencing rapid erosion. Between 1993 and 2020, the shoreline retreated at a rates of between 40 and 80 centimetres per year. The town is also within an area of continuous permafrost dominated by icewedge polygons, which are particularly vulnerable to permafrost thaw. This means that all its infrastructure, including people's homes, is at risk of damage or collapse as the warming climate causes increased erosion of the permafrost underlying the settlement.

Like so many communities in northern Canada and across the Arctic, Paulatuk is not accessible by road. Instead, it relies on air transportation and marine shipping to link both people and cargo to the rest of the country and beyond. The summer sealift, the only way to transport very large or heavy items to the community, is only available during the icefree season in the Amundsen Gulf. This makes air transportation, which is available year-round, vital to communities like Paulatuk. Airports are some of the Arctic's most valuable assets located on permafrost and are considered critical infrastructure for communities. Canada alone operates over 85 airports in the north, most of which have unpaved gravel runways which can become significantly weaker after periods of heavy precipitation or spring thaw.

Paulatuk's airport is situated less than 4 metres above sea level, built on ice-wedge polygons, and located 100 metres from the eroding shoreline. Subsidence caused by thawing permafrost threatens to breach the berm built to protect the runway. This could lead to flooding in an adjoining depression located at the southwestern corner of the airstrip, potentially creating problems with erosion. The airport and other infrastructure may also prevent meltwater from ice-wedge polygons from draining in the spring. The resulting ponding may damage the runway and embankments and potentially delay the refreezing of the active layer in the autumn. Furthermore, the low elevation and location of the airstrip subjects it to impacts from more intense storms, sea level rise and coastal erosion. Remote Arctic communities such as Paulatuk cannot afford to sever this critical link and need solutions to address the threats posed by climate change.



Paulatuk's airport is built on permafrost, with sand and gravel elevating it about 2 metres above the ground. The runway is located less than 100 metres from the shoreline facing the Amundsen gulf. It has experienced rapid erosion up to 80 centimetres per year.

ACT ANT ANT ANT

The airport's runway generates disturbances in the permafrost and prevents drainage of meltwater from ice wedge polygons in the spring, resulting in ponds.


> Ice-wedge polygons are a common feature of this northern landscape. Different polygon densities reveal varying ground-ice concentrations.

Paulatuk's infrastructure, including houses, are built over ice-wedge polygons. They are particularly vulnerable to permafrost thaw.

100 m

SOURCE: MODIFIED FROM VIEIRA ET AL. (2019). BY LEVI WESTERVELD | GRID-ARENDAL (2023)


Terra infirma III Keeping cold food cold in Alaska



Ice cellars are a form of Arctic Indigenous technology that take advantage of the permafrost to provide natural year-round refrigeration for food. They are found in Alaska's far-north Iñupiat communities and in parts of the Canadian and Russian Arctic. As well as being a cost-effective solution to storing meat, they play an important role in Indigenous cultural identity, representing the transfer of knowledge of food preservation from one generation to the next. Ice cellars on Alaska's North Slope come in a variety of shapes and sizes, but on average provide 10 cubic metres of space. Typically, they have a shaft 1 to 6 metres deep, ensuring that the ceiling of the cellar is well below the active layer of permafrost, are accessed through a wooden hatch or shed built over the entrance. The large size of the cellars means that not only can waterfowl, seals, and caribou be stored, but also whale meat, which requires more room for longer storage periods. While temperatures in ice cellars do not reach -18 °C as recommended by the United States Department of Agriculture for meat storage, they do allow for meat to safely age and create a particular taste that is culturally preferred.

Concerns have been raised about the future of ice cellars in Tikiġaq (Point Hope) and Utqiaġvik (formerly known as Barrow) in Alaska due to their recent degradation. At least 50 ice cellars used to exist in Tikiġaq, while today only 20 are thought to remain. A similar trend is occurring in Utqiaġvik where many ice cellars are being abandoned or considered unreliable. This decline has been attributed to various reasons, including: not being able to access the cellars safely, the presence of mould in the cellar, splintering or bowing of wood lining the cellars, water pools, seepage of briny water, the loss of sediment cohesion (sloughing), and the so-called catastrophic failure where cellars flood or collapse in on themselves.


Utgiagvik's ice cellars were the subject of a decadelong research study between 2005 and 2015. This study revealed that temperatures stayed fairly stable over the study period, with mean annual temperatures ranging from -7.8 to -6.4 °C. While warmer than the constant temperature of -11 °C reported about a century earlier, the researchers pointed to other possible contributing factors alongside permafrost thaw that could be negatively impacting ice cellars. These include local permafrost conditions, hydrological changes, irregular or poor maintenance, factors associated with urban development, and snow redistribution. Salinity is an important factor, for example, because higher salt content can lower the freezing point of water, which can decrease the ground's bearing capacity. Utgiaġvik's sediments are thought to have been deposited under marine or brackish conditions. The use of snowploughs may also affect the mechanical properties of cellars: snow is often piled on top of cellars insulating the ground and warming underground cellars, while snowbanks left over in spring can generate large amounts of meltwater that then flood cellars.




Seasonal surface deformation 2015–2019 average (cm per year)

NOTE: downward deformation in summer when the active layer thaws, and upward deformation in winter when it freezes.

8 cm




Development areas Bedrock (stable) Sediments (unstable)



Northern Isthmus is a new Northern Isthmus is a new development area north of Ilulissat, including commerce and tourism, housing, and institutional buildings. Parts of different zoning areas overlap permafrost sediments with significant annual deformation rates.

Boreholes show very high ice content, indicating a high risk of large permanent deformations as the permafrost degrades.



Terra infirma IV Urban planning in Ilulissat

Located on the country's west coast, Ilulissat, is the third largest town in Kalaallit Nunaat (Greenland) and a UNESCO World Heritage Site. As the top tourist destination in the country, the town is undergoing a period of expansion and economic growth. A new international airport is planned and, along with an influx of tourists, is expected to bring new business opportunities to the region. A new access road connecting the town to the airport will also encourage the development of new urban areas, including business parks and residential areas. But this planned expansion is facing some significant technical challenges related to permafrost. If not adequately accounted for, these could compromise both the new infrastructure and Ilulissat's drinking water supply.

Most communities in Kalaallit Nunaat are located on bedrock outcrops, so the main concern around permafrost relates to linear infrastructure, such as the roads and pipelines which cross sedimentary basins. Ilulissat's planned expansion will also be on bedrock, but both the airport runway (which will be 2 km long) and the roads leading to it lie on sedimentary permafrost, which is extremely ice rich in places. For the new airport, it was decided to completely excavate the permafrost sediments underneath the planned runway and replace it with other stable rock-fill material. For other linear infrastructure, however, this option is too expensive as the ice-rich permafrost can be many metres deep in places. Permafrost sediments around Ilulissat have an extremely high ice content in the upper parts. Meanwhile, the deeper sediments, having been deposited at a time when the relative sea level was 50 metres higher than today, have a high salt content. The characteristics of Ilulissat's permafrost are such that, as it thaws, its high ice content causes the ground to subside, while the residual saline pore water brings these sediments to temperatures as low as -2 °C.

With these challenges in mind, scientists are assisting the municipality of Ilulissat in their urban planning efforts, focusing on understanding the properties of the permafrost in and around the town's sedimentary basins, particularly in terms of salinity and ice content. Several sediment cores, each with varying ice and salt contents, have been collected around Ilulissat. The sediment properties and the ice content are analysed in the laboratory and then different loads are applied to each sample to determine how thawing affects surface deformation. This allows the scientists to study how salinity, ice content, and thawing influence the mechanical characteristics of the sediment and the risk this might pose to roads and other infrastructure lying on top. Ultimately, scientists aim to develop a map and evaluation framework which can inform the municipality about the risks of construction in different areas.

Nothing in isolation *Health and wellness and permafrost*

The lives of many people living in the Arctic are intimately linked with their natural environment. As climate change intensifies, a major concern becomes how the rapid environmental changes experienced in the Arctic affect both the physical and mental health and well-being of its inhabitants. Coastal erosion, decreased sea ice, weather changes, permafrost thawing on land and beneath the sea, and shifting ranges and distribution of wildlife are all happening at an unprecedented pace. This is creating new risks in terms of health and safety from bio- and geohazards such as slumps and wildfires, food and water security, and the emergence or re-emergence of contaminants and infectious diseases, including zoonoses (diseases that spread from animals to humans).

Climate change, among other challenges, is also recognised as taking a toll on the mental health, wellbeing, and quality of life of people. Research in the community of Rigolet, Nunatsiavut, Canada, revealed that changes in weather, snow and ice stability, and wildlife and vegetation patterns are disrupting landbased activities, damaging cultural identity and placebased solace. This has been contributing to decreased mental health and well-being, along with feelings of powerlessness and loss of control. These factors can lead to increased family stress, a greater likelihood of heightened drug and alcohol consumption, and suicidal thoughts.

Understanding the health implications of rapid environmental change in the Arctic requires a holistic approach. "One Health" recognises that human health is interconnected with environmental and animal health. It requires collaboration across various disciplines, including both natural and social sciences and knowledge systems to identify risks and to prevent and manage health risks in humans, animals, and their shared environment.

The One Health approach also recognises and seeks to include traditional and Indigenous knowledge as one form of knowledge. Indigenous communities have their own worldviews of health and well-being, which are very often grounded in the strong connection to the land and the community. In Nunavut, for example, the Inuktitut language conveys a diversity of perspectives on health, including Inuusiqatiginniq (the way of being for a person). This generally includes a holistic view of wellness that considers the surrounding environment, individual behaviours, and relationship to others.

One Health and similar approaches are now being used to understand how permafrost thaw affects human health and well-being. Permafrost degradation has several possible direct and indirect impacts on humans, animals, the environment, and the interactions between them. A study from north-western Alaska found that permafrost thaw is decreasing water and food security, leading to higher mental stress. The community of Point Hope/ Tikigaq, for example, gets its water from a small tundra lake situated on top of permafrost. Higher summer temperatures, however, have led to increased algae and other biological growth, including mosquitoes, requiring water filters to be changed up to 50 times a day as opposed to the normal four in order to maintain drinking water quality. Likewise, in both Tikigaq and Kivalina, many traditional ice cellars now thaw in summer, rendering harvested bowhead and beluga meat and blubber unsafe to eat and impacting food security.

The perception of health challenges resulting from permafrost thaw vary across the Arctic. A case study from northern Canada showed that permafrost thaw creates challenges for pursuing traditional livelihoods, travelling on the land, and maintaining infrastructure. This may further impact people's self-rated health, work-life balance, and feeling of empowerment. In Kalaallit Nunaat's (Greenland) Disko Bay and on Svalbard, studies found that permafrost thaw is not perceived as a direct risk to health and mental wellbeing, despite the recognised impacts on culture and the built environment, and from natural hazards. People recognised the problems associated with permafrost thaw in all these areas, but still reported good well-being and quality of life. These results suggest that an awareness of challenges related to permafrost thaw supports the overall well-being. While these results were derived from a relatively limited number of participants, they could be expanded to help communities build adaptation capacity.

One Health

HUMAN

5

NIMA!

An approach that recognises that human health is closely related to the health of animals and the environment. This is particularly true in the Arctic where a community's well-being relies on healthy animals and the environment.

Human health is more than the absence of disease; it encompasses physical, mental, behavioural, cultural, and spiritual health.

This graphic illustrates some of the ways that permafrost thaw affects the health of the environment, animals, and humans.

ENVIRONMENT:

- Processes such as coastal erosion, heave, and subsidence are driven by permafrost thaw. Changes to the physical environment can be a risk to the health of both humans and animals.
- 2 Water and air quality are affected by permafrost thaw. The release of sediments and contaminants in the water due to erosion and thermokarst, as well as an increase in Arctic fires, are detrimental to the health of the environment.

ANIMALS:

(1)

ENVIRONMENT

.

- As permafrost thaws, contaminants (A) such as mercury are released and transported by rivers to the Arctic Ocean. They travel up the food chain, affecting the health of both small and large animals.
- 4 Location and migration routes of animals are changing as they adapt to changes taking place in the environment, this affects food accessibility for humans.
- 5 As temperatures increase, old pathogens may re-emerge, potentially affecting the health of both animals and humans

HUMANS:

- 6 Limited access to healthy animals, damaged transportation infrastructure and food storage facilities such as ice cellars, can affect food security and human well-being.
- Permafrost thaw changes the cultural landscape of the Arctic. On riverbanks and coasts, important sites such as graveyards can be eroded and damaged due to flooding, affecting human well-being.
- (8) Infrastructure damage due to permafrost thaw can limit access to safe housing, water and sanitation facilities, and other institutional and human-built infrastructure.

BY LEVI WESTERVELD | GRID-ARENDAL (2023)



© Olga Borjon-Privé (Oluko)

FORTRAIT EXECUTIVE DIRECTOR, QAUJIGIARTIIT HEALTH RESEARCH CENTRE, IQALUIT, NUNAVUT CANADA Gwen Healey Akearok

I was born and raised in Iqaluit, the capital of Nunavut on the southern coast of Baffin Island, just south of the Arctic Circle. I am an epidemiologist, community health researcher, and founder of the Qaujigiartiit Health Research Centre. Through my work, I seek to answer pressing health and well-being concerns of my community.

I would describe my work at the research centre as heart-centred and relational. We work very holistically, for example by ensuring access to food that keeps us healthy, access to shelters that keep us safe from the weather and cold, and access to healthy animals that we harvest. These are all interconnected. We nurture these relationships in a way that promotes health in all these areas. I work jointly with all members of the community including elders, students, and government officials.


People outside the community, including the media, often think that mental health and high suicide rates are some of the main issues of northern communities. But they rarely look into the determinants of the mental health challenges, such as trauma history and the generation of parents, grandparents, and great-grandparents being sent to residential schools. Building on the relationship between community members and the land and reconnecting people with language and culture has helped community members heal and move down a path to overcoming traumatic events. Now, changes in the landscape due to climate change, especially changes in the migration of different animals, for example, is a source of additional stress. People talk about those concerns, the health of animals, the ability to harvest. If you can't go out and have a relationship with the land and water, that can have a significant impact on the mental well-being of our communities.

On permafrost, damage to infrastructure is one of the most important public health risks associated with thawing ground. Unpredictable swings in weather that are undoubtedly climate change-related cannot be handled by our infrastructure. Our buildings are built on steel piles right into the bedrock and permafrost. When there's heavy rain, the water and thawing permafrost, for example, causes the buildings to shift, which can lead to the collapse of structures. If our houses, our grocery stores, and all these things we rely on for shelter and food won't stand up, it's a public safety concern – which people don't usually consider.

I am also a mother, and when thinking about future generations it is hard not to think about my own childhood. We could crawl around caribou in the tundra, you could go on a walk and run into one. But now my children have never seen a caribou. It's hard not to think about all the other things that will change by the time they become adults. As a parent, you want to protect them from everything, but everything in our communities is changing so quickly, that it's hard to imagine what they might encounter. I remain positive about the future, however, thanks in part to the innovative energy of my community, but also to the increased access to knowledge and modes of communication. All of our communities have access to the Internet and cell phone coverage, so accessing knowledge from other people and information is a huge asset. I have no doubt the future generation will be even more creative when they are adults, nurturing connections across the Arctic that will help to move things forward for the good of everyone.

Toxic grounds *Contaminants and environmental health*

Environmental contaminants are chemical substances released by human activities that may adversely affect living organisms, including humans. Environmental contaminants found in the Arctic include heavy metals, radioactive compounds, black carbon (soot



SOURCE: SCHUSTER ET AL. (2018). BY LEVI WESTERVELD | GRID-ARENDAL (2023)


from fires, shipping, or industrial activity), plastics (including microplastics), and other synthetic chemicals (including persistent organic pollutants such as dichlorodiphenyltrichloroethane (DDT), polychlorinated biphenyls (PCBs), and dioxins). These chemicals remain stable in the environment for a long time and can bioaccumulate in the food chain, posing a danger to wildlife and human health. Many contaminants originate from more populated areas and industrial activities further south. The contaminants are carried north via air, rivers, and ocean currents and are deposited across the Arctic where they are locked in surface sediments, fresh water, seawater, sea ice, glaciers, and permafrost.

Localised sources of contaminants are also important in some areas of the Arctic, especially in and around industrial sites such as mines and other heavy industries. Environmental contaminants from these operations may include drilling and mining remains, drilling muds and fluids, mining heaps, heavy metals, spilled fuels, and radioactive waste. As the human footprint in the Arctic grows, both local and long-distance sources of contaminants are expected to increase, including chemicals of emerging Arctic concern, such as fluorinated surface protectants (PFAS) and current use pesticides.

The changes occurring in Arctic snow, ice, and permafrost raise a serious question about whether these contaminants, deposited over decades, may be released again into the environment. One of the main issues relating to thawing permafrost is the remobilisation of previously deposited contaminants. Contaminants of special concern include persistent organic pollutants such as pesticides (e.g., DDT) and industrial chemicals such as PCBs, flame retardants, or fluorinated surface protectants (PFAS). Mercury, a naturally-occurring toxic compound, is another source of concern. Human activities have altered the global mercury cycle and released large quantities into the air, land, and water. Research suggests that permafrost is a very large reservoir for mercury compared with other soils, the ocean, and the atmosphere. However, there is a great deal of uncertainty over how much it actually contains. Regardless, it remains widely accepted that permafrost degradation could have major implications for mercury redistribution worldwide.

As permafrost thaw puts Arctic infrastructure at risk, there is also a concern about increasing environmental contamination from industrial accidents. A recent estimate suggests that more than 10.000 contaminated sites exist on permafrost in the Arctic. Future projections of permafrost dynamics indicate that up to one-third of those sites may be exposed to permafrost degradation between 2050 and 2100, which can both affect the stability of industrial infrastructure and remobilise contaminants. How changing permafrost conditions will affect each individual site, however, depends on the conditions of where it is situated. For example, waste disposal sites in some locations are now selected with greater care and the ground materials at the disposal site may be less vulnerable to the impacts of permafrost thaw.

Most remote Arctic communities have limited waste management facilities and depend on dumpsites with open burning or on sewage lagoons. If the permafrost beneath these sites thaws, it may no longer provide a barrier between adjacent soils, groundwater aquifers, and water sources.



Coming back to life Re-emerging pathogens


Since the late nineteenth century, scientists have known that permafrost harbours pathogens, diseasecausing microorganisms that include viruses, bacteria, fungi, parasites, and potentially prions. As long as permafrost remains frozen, these organisms are dormant, sometimes for thousands of years. As permafrost thaws, they can come in contact with air and water as well as animals and humans, creating conditions the pathogens need to grow and spread. The potential release of pathogens from thawing permafrost is of growing concern for human, animal, and plant health.


Pathogens that form endospores (dormant cells) have a high chance of survival under harsh environmental conditions and may pose the greatest risk from thawing permafrost. They include fungal pathogens, species of *Clostridium* bacteria that cause tetanus and botulism. and the anthraxcausing bacterium Bacillus anthracis. Anthrax, a disease which primarily affects domestic and wild-hoofed animals, was once widespread across the Russian Arctic until animal vaccines became available. Between 1897 and 1925, over 1.5 million reindeer died from anthrax, their carcasses buried in shallow graves. There are more than 13,000 of these burial grounds in Russia, over half of which are located in permafrost. A recent outbreak of anthrax on the Yamal Peninsula in 2016, which led to the death of one child, the hospitalisation of over 20 adults, and the death of several thousand reindeer, was traced back to the thawing of a

75-year-old reindeer carcass under unusually dry and warm conditions. This outbreak highlights the very serious, localised risk of anthrax to humans and animals in certain parts of the Arctic, as well as the risk to traditional livelihoods.

Permafrost thaw is also leading to rapid changes in boreal ecosystems, including changes in the composition of fungal communities, the diverse group of organisms which includes yeasts, moulds, and fungi. Pathogenic fungi originating from thawing permafrost could be potentially dangerous to human health, as well as to forestry and agriculture. For example, *Hymenoscyphus fraxineus*, a fungus originating in Asia that has devastated ash trees across Europe over a 30-year period highlights the ease with which fungal pathogens and other invasive species hitchhike on globalised trade.


As demonstrated by the COVID-19 pandemic (caused by the severe acute respiratory syndrome coronavirus 2) and the re-emergence of the highly pathogenic avian influenza A(H5N1), RNA viruses have high mutation rates and can readily adapt to new environments, making them an increasingly important group of zoonoses (diseases that transmit between animals and humans). Within the Arctic, fragments of RNA from the 1918 Spanish influenza virus have been discovered from mass graves in the Alaskan tundra. In Siberia, fragments of DNA from the smallpox virus have been discovered in 300-year-old mummified humans. Permafrost is






SOURCE: CAFF (2001). BY LEVI WESTERVELD | GRID-ARENDAL (2023)

home to ancient "giant" DNA viruses, which could be revived under conditions of permafrost thaw, though many of these "zombie viruses" pose no threat to humans. Many scientists are now calling for more attention on and research into paleo-microbiology to better understand, predict, and plan for the resurgence of ancient infections.





Frozen assets l *The formal economy*

A common feature of Arctic regions is their mix of formal and subsistence economies. The formal economy includes large-scale resource extraction, such as oil and gas, timber, fishing, and mining, all of which serve national and international markets. Local economies, on the other hand, provide goods and services to the local population, including public sector jobs. Traditional, subsistence activities, which include hunting, fishing, herding, and gathering, can also form a significant part of the local economies.

Calculating the potential economic risk and cost of permafrost thaw can be done in several ways. Standard economic indicators like GRP* can be combined with the extent and types of permafrost to understand how much economic activity is potentially at risk from permafrost thawing. The entire Arctic permafrost region is, in fact, one of the most productive economic regions on Earth. With a GRP of USD 458 billion (2017), this is equivalent to a GRP per capita of over USD 71,000, a figure that is higher than for Germany and the United States and just below the GRP per capita of Switzerland. However, the financial benefits of production are not necessarily re-invested in the region but are often transferred out so these high numbers may hide some of the poverty observed among local communities.


GRP per capita can also indicate the availability of economic resources to adapt to the adverse impacts of climate change. Permafrost regions with low GRP per capita possess limited resources for adaptation and are thus considered more vulnerable. The type of permafrost is also a contributing factor: economic activities situated in areas with discontinuous and continuous permafrost will experience greater change than those in sporadic and isolated patches, so that regions such as Sakha Republic and Yamalo-Nenets are considered to be the most vulnerable. GRP per capita has also been combined with other social and physical indicators to create the Arctic Permafrost Population Vulnerability Index.

Permafrost thaw has already resulted in widespread impacts and costs on all parts of society. Settlements may have to be relocated, new housing may need to be built, existing buildings will need to be repaired to avoid collapse, while roads, railways, airports, and pipelines will need to be fixed. While calculating the economic cost of permafrost thaw in the Arctic is an emerging field of research, several estimates are available, many of which focus on the cost of maintaining, replacing, or upgrading existing infrastructure. Typically, these will explore several scenarios, including a lower or higher emissions scenario, depending on how carbon intensive the future direction of our global society will be. In certain cases, the extent to which timely adaptation measures can reduce future damages and thus costs is also calculated.

These types of estimates help to communicate the urgent need to reduce emissions, while simultaneously pointing to the need to adopt smarter adaptation choices in the near term to reduce costs in the long term. For example, the cost of maintaining Alaska's public infrastructure under a business-as-usual scenario is estimated at USD 5.5 billion by 2099 (under a high-emission scenario) or USD 4.5 billion (under a lower-emission scenario). Meanwhile, the adoption of proactive measures, mainly related to flood measures on Alaska's roads, was estimated to bring those costs down to USD 2.9 billion and USD 2.3 billion for highand low-emission scenarios, respectively.

Moreover, the contribution of carbon and methane into the atmosphere can be calculated in terms of economic costs to global society. The global costs associated with continued permafrost thaw through the release of carbon and methane are projected to far outweigh any benefits, such as the spread of agriculture northward, underscoring the need for urgent action to stabilise emissions. In 2015, researchers estimated that continued permafrost thawing could cost the global economy USD 43 trillion by 2100 - roughly equivalent to half the annual global economic output today. Even if the world manages to stabilise temperatures well below 2 °C above pre-industrial levels, permafrost thaw, through the release of greenhouse gases is calculated to cost the global economy almost USD 25 trillion, or about 4 per cent of the total cost of climate change to the global economy.

^{*}Gross Regional Product is a measure of the total value of goods and services produced in a region. When GRP is divided by the total population, a per capita value is obtained, which indicates the average income per inhabitant.



Frozen assets II *Traditional and subsistence activities*


Traditional and subsistence activities are mostly unaccounted for in official economic statistics across the Arctic (excluding Alaska), which makes such activities largely invisible in measures of income or wealth, including GRP. However, many studies of individual communities highlight the crucial importance of subsistence activities, particularly in terms of the food obtained from fishing, hunting, and foraging, also known as "country food". It is critical to understand how permafrost thaw may impact the type and abundance of country food as well as access to it.

Market pricing techniques* have been used to put a monetary value on subsistence activities happening on permafrost landscapes of three coastal communities: Aklavik (Canada), Qeqertarsuaq (Kalaallit Nunaat/ Greenland) and Longyearbyen (Svalbard, Norway). Each community differs in the types of subsistence activities undertaken by its inhabitants, including the proportion of country food in their diets.

Almost 50 per cent of the food eaten in Aklavik and 60 per cent eaten in Qeqertarsuaq are obtained through hunting, gathering, trapping, and whaling, while in Longyearbyen it is only 10 per cent. Should the permafrost landscapes and the food they supply disappear, inhabitants of Aklavik and Qeqertarsuaq would have to pay EUR 240 and EUR 460, respectively, to purchase this food, which represents 43 per cent and 17 per cent of their monthly incomes per capita. While these numbers are theoretical, they highlight the real economic importance of subsistence activities and a potential cost-of-living crisis, should country foods become unavailable in the future. Store-bought foods are very expensive given the high commodity prices across the Arctic. Such high costs would severely impact purchasing power and reduce the well-being and ability of inhabitants to satisfy their basic needs.

The way in which Arctic ecosystems and plant and animal communities will respond to thawing permafrost is an emerging field of research and it is relatively unknown how these changes will affect subsistence activities. Changes in soil temperature. moisture, and nutrients are likely to alter plant composition. Where ground-ice content is high, subsidence of the landscape may lead to the formation of ponds and lakes, shifting a terrestrial habitat to an aquatic one. Changes in the vegetation and landscape may lead to altered animal migration routes or changes in animal species composition, and influence humans' access to country foods. For example, hunting may take longer if hunters need to go around thaw slumps, or because moose hide better behind taller willows. As the climate warms, some Arctic species may be displaced by others better adapted to the new conditions. On land, changes in ground conditions could lead to a change in hunting routes or reduced snowmobile access. It is during this "transition phase" when human societies are adapting to the changes that the costs may be greatest.

Proportion (%) of the most-consumed species in Aklavik, CA.



SOURCE: RAMAGE ET AL. (2022). BY GRID-ARENDAL (2023)

^{*}Market pricing techniques use market prices from a surrogate market to assess the monetary value of non-market goods and services. For example, for assessing the value of country foods, the equivalent price that it would cost to obtain that type of food from a conventional market (shops, stores, or supermarkets) is used.

Cultural homeland Alaas landscapes in Yakutia

Situated in the Russian Far East, the Republic of Sakha (Yakutia) is a unique permafrost region. The Central Yakutia lowlands, which extend along the middle basin of the Lena River, are characterised by ice-rich permafrost which can be several hundred metres deep. A key feature of this region are the alaas permafrost landscapes, which consist of thermokarst depressions or drained thaw lakes with large areas of subsided ground surface. Over 16,000 alaas exist in Central Yakutia covering an area of 4,400 km².

In the Sakha language, "alaas" means "a field or meadow, surrounded by forest; a piedmont valley".

Sakha people depend upon alaas and consider it their homeland, imbued with spiritual qualities in accordance with their animistic beliefs. Later, as land development demanded the scientific study of alaas ("alas" in English and Russian) to build and exploit resources, the definition became highly technical: a landform distinguished by large areas of subsided ground surface resulting from the thawing of ice-rich permafrost.

The formation of alaas can take anywhere from decades to hundreds of years. The process starts when the area becomes naturally deforested, followed by deepening subsidence and ice-wedge degradation. As the ice wedges thaw, the depression is filled with meltwater and becomes a lake. In the final stage, the landscape dries out and grassland is formed on the middle and outer edges of the depression.

Alaas grassland areas have attracted and sustained animal husbandry for at least 500 years, forming the basis for the thriving communities and culture of the Sakha people. Between the twelfth and fifteenth centuries, the region's Turkic ancestors migrated north from the Lake Baikal region and settled around the alaas within the Lena and Viliui River basins. Using practices carried over from their southern ancestors,




Sakha successfully manipulated the alaas landscapes to maximise hay production for their cattle and horses. They drained waterlogged fields during wet years and created a system of canals and dams to hold water in times of drought. They also created more pasture and haylands by draining lakes and removing forests by burning. Alaas are considered sentient beings according to Sakha's historic beliefs. Sacrificial foods as well as rituals and blessings are offered to the alaas. Extended clans consisting of groups of five to 15 households would live on the alaas, migrating between summer and winter camps.

Starting from the seventeenth century and coinciding with the introduction of Russia's land tenure taxation system, cattle surpassed horses in importance, and the alaas were primarily managed to produce hay as fodder for cattle. The Soviet period, starting in the 1920s, brought the collectivisation of farms and the formation of compact villages. The alaas began to be used for large-scale crop production, pasture, or fodder. From the 1960s onwards, there was a shift from using native Sakha cattle to European breeds. Not able to withstand the harsh winters, the European breeds needed to remain indoors for 9 months of the year and fed fodder. The result of this was an intensification of hay farming and mass drainage of the alaas to create more grassland. The collapse of the Soviet Union brought further changes: state farms were disbanded and largescale cattle breeding was no longer profitable. Most households adapted, however, by holding a few cattle for their subsistence. Many of the grasslands, which were converted to croplands during the Soviet period, were converted back to natural hayfields.

Like other landscape features, alaas are not immune to the ongoing changes in permafrost. Many thermokarst lakes are expanding in size and volume with increased summer precipitation, above average snow cover, and increased permafrost ice meltwater as the principal causes. Alaas have also seen changes in their flood regimes. Typically, they only flood during the spring due to melting ice and snow, but now summer floods have become more common. While spring floods bring nutrients and are considered beneficial, summer floods make hay production impossible. Waterlogging of alaas not only limits hay production, but limits access to forests and other resources. For some households, relocation is needed due to the increasingly damp conditions.

The ongoing use of alaas is dictated not only by environmental conditions, but also by social and demographic trends and consumption patterns. Due to a combination of factors including a lack of fodder from the land, youth outmigration to Yakutsk, and a shortage of workers, some inhabitants have abandoned cattle breeding. Instead, many are now opting for horse breeding, which generally involves less work. Horses also carry more prestige in post-Soviet Sakha culture.





Holding Tight

Adaptation to Permafrost Thaw

Bumpy road ahead *Transportation infrastructure and permafrost*

A wide array of transportation technologies are used to move people and goods into, out of, and around the Arctic. These range from raised wooden paths over muddy ground to elevated utilidors* above frozen ground, from year-round highways to seasonal ice roads, and from marine ports to airports which provide some communities their only link to the rest of the world. To some degree, permafrost degradation will affect most of these transportation technologies, if not all.

There are two primary drivers of ground thermal changes and thawing around transportation infrastructure: increasing temperatures caused by climate change and direct impacts from infrastructure itself (e.g., localised heating of the ground and disruption of insulating vegetation). Combined, these effects double the pressure on permafrost, stressing transportation infrastructure as well as Arctic lives and livelihoods.


*A utilidor is a utility corridor or tunnel used to transport electricity, water, and sewage. In permafrost regions, insulated utilidors are placed above ground to prevent permafrost thawing and decrease the risk of infrastructural damage.

Changes in adjacent permafrost terrain is also a threat in many places, particularly thermokarst features. When permafrost thaws, ground ice melts leaving behind depressions in the landscape. These fill with meltwater to form ponds or larger thermokarst lakes. These depressions expand during summer when air temperatures are higher, which fragments the landscape. Expanding thermokarst features threatens foundations of transportation infrastructure as it weakens the sediment strength, and, in some cases, even leads to total collapse.

Both point infrastructure (e.g., ports and airports) and linear infrastructure (e.g., roads and pipelines) are vulnerable to permafrost thaw. Infrastructure development in coastal areas has been growing rapidly in recent decades, and much is built on ground which will be affected by thawing permafrost. This infrastructure is often used for the transportation and export of natural resources. Damage caused by a reduction in the ground's bearing capacity as the permafrost deteriorates poses a risk for current and future economic development. At airports, distortions created by permafrost thaw are negatively impacting the integrity of runways, causing safety concerns and potentially paralysing this type of transportation. Linear infrastructure is vulnerable to permafrost thaw as changes in the heterogeneous ground cause differential settlement and distort the structure. One weak point along linear infrastructure can make the whole network unusable (for example, a break in a pipeline). The length of infrastructure and the weak points along its route are therefore important factors to consider when assessing overall vulnerability.

Transportation in the Arctic is multimodal, meaning that different transportation technologies are interlinked in a network to move people and goods between places. For a multimodal network to function, it requires the integrity of all the different infrastructural components. A defective runway or road may have negative impacts across the network. Almost 30% of roads located in Arctic permafrost regions are considered at risk of ground subsidence under moderate greenhouse gas emissions scenarios, rising to 44% under high emissions scenarios. Among these, the largest number of roads at risk is in Russia, where the estimated average costs for replacement range from about USD 50 to USD 70 million, depending on the scenario.

Multimodal transportation infrastructure and impacts of permafrost degradation




Mitigation methods for transportation infrastructure


Since the 1960s, greater efforts have been made to understand the physical impacts of climate change on transportation infrastructure and to adapt construction and maintenance to those changes. Improved risk assessments of existing and planned infrastructure projects can help better manage impacts from permafrost thaw. For example, integrating high-level details from geotechnical models with large-scale, long-term land surface models may better project the combined impacts from climate warming, infrastructure development, and permafrost degradation in order to adapt to both slow and rapid changes.

In addition to better planning, there are three broad approaches to protecting infrastructure from the impacts of thawing permafrost: heat extraction, heat intake reduction, and reinforcement. Heat extracting methods emulate winter conditions to cool embankments. The aim is to remove accumulated heat in the embankment thereby reducing the impact of infrastructure on the thermal structure of the ground. Techniques include air convection embankments that reduce summer warming and enhance winter cooling, air ducts to capitalise on local wind conditions, heat drains, thermosyphons, and sloped embankments to limit snow accumulation.


The sun is a major source of heat during the Arctic summer. Methods to reduce or prevent heat intake focus on limiting the amount of energy that gets transferred to permafrost. This may be achieved through insulation, building on piles, using materials with higher albedo (reflection), or using sun-sheds to deflect sunlight away from vulnerable infrastructure.

Reinforcement methods used to limit permafrost degradation around infrastructure include strengthening embankments and using controlled thawing prior to the construction of embankments. Sections of the Alaska Highway, for example, use pillow embankments while the Inuvik-Tuktoyaktuk Highway in the Northwest Territories, Canada is testing the use of woven geotextile to mitigate slope movements.






Heat intake reduction techniques



SOURCE: HJORT ET AL. (2022). BY LEVI WESTERVELD | GRID-ARENDAL (2023)




Undermined *Mining infrastructure and permafrost*

While all Arctic nations have mining operations, only Canada, Kalaallit Nunaat (Greenland), Russia, and the state of Alaska have operations in permafrost landscapes. In Alaska, gold and zinc are the main minerals being extracted (except for oil and gas), and the industry is a significant contributor to the region's economy. Diamond mining also plays an important role in the economy of northern Canada, especially in the Northwest Territories. In Kalaallit Nunaat, mining only accounted for 0.4 per cent of the gross regional product (GRP) in 2018, but this sector is likely to increase in the coming decades as Kalaallit Nunaat is home to some of the largest mineral deposits in the world, including rare earth minerals. A large share of the world's mineral extraction occurs in Russia's vast territories, including iron and ferro-alloy minerals, non-ferrous minerals, precious metal ores, and industrial minerals. Combined with petroleum extraction, these sectors constitute 60 per cent of the Russian Arctic's GRP.

Many mines in the Arctic are found in areas impacted by permafrost thaw, and subsequent landscape changes are creating challenges and risks for these operations. Mining infrastructure can impact the energy exchange between the surface and the atmosphere, known as the surface energy balance, which may cause local thawing. The accelerating permafrost thaw caused by global warming, however, has the potential to change the landscape even more drastically. While permafrost thaw will not make mining in the Arctic impossible, it will have consequences in terms of safe operations, particularly when it comes to impacts on freshwater systems and the secure storage of mine tailings. Tailings are a byproduct of mining, comprising a mixture of solids and fluids used in the extraction process. In the Arctic, permafrost has been an effective barrier in tailing deposits, preventing contaminants from seeping into the environment. However, as permafrost thaws and the ground ice within it melts, the hydraulic conductivity increases. This means that water, along with any dissolved elements, can move more easily and potentially infiltrate groundwater or local water bodies. As a result, relying on permafrost for tailings containment is no longer a viable option and other solutions are needed.

One way to ensure containment of tailings is to reinforce the physical strength of deposits through thick gravel covers or dry stacking. The latter approach aims at reducing the moisture levels in tailings to lower the risk of seepage and minimise the risk of catastrophic failure. Another option to secure tailings is using thermosyphons, a device designed to cool the ground by extracting heat from the soil to impede thawing below and around tailings dams.

Mining projects should build on collective knowledge through the sharing of experiences across the Arctic as a way of contributing to best practices for safe and sound operation in the region. For example, the International Council on Mining and Metals, a membership organisation consisting of mining companies operating across the globe, has recognised permafrost as a potential risk factor in the years to come, particularly for mines that have long life spans. The design of mines should be holistic in the sense of meeting the diverse challenges linked with Arctic mining. These challenges include remoteness, limited infrastructure, harsh weather conditions, water availability, climate change, and the long-term stability of tailings. Planning for a future where permafrost conditions are changing will be fundamental to ensuring a holistic design. Such designs should always include scenario modelling, for example, by studying how ground and hydrology will change as the ground warms.



Keeping the light on *Energy infrastructure and permafrost*


With great distances and long cold and dark seasons, access to energy is essential in the Arctic. These same factors, however, also put pressure on energy security. Fossil fuels remain the primary source of energy throughout the Arctic region. To cope with the increasing costs of fossil fuels and help reduce greenhouse gas emissions, some communities are considering alternative sources of energy. For example, in Kalaallit Nunaat (Greenland) 70 per cent of energy comes from renewable sources, mainly hydropower. Two major hydropower plants have been constructed in permafrost areas. The difficulty of building on permafrost, coupled with increasing permafrost thaw, pose challenges to all types of energy development.

Many communities, especially those in remote and isolated areas, rely on fossil fuels that must be transported long distances, often from outside the Arctic. As thawing permafrost impacts transportation infrastructure like roads and runways, it may also affect the energy security of many communities. In Russia, for example, changes in permafrost are one of several factors that have left the infrastructure supplying Russian Arctic communities with energy at risk. Because of unstable deliveries, fuel costs have risen. With an increase in Arctic oil production, a shortterm improvement can occur. Eventually, however, the energy mix will need to be diversified and the benefits may not weigh up against potential impacts on the environment.

Together with communities, government, and private partners in Nunavut, Canada, WWF is developing small-scale energy infrastructure. The project includes constructing solar panels to provide energy for community freezers and other needs. Freezers are important for communities where hunting and fishing are widely practised, and the need for them is greatest during summer when the temperatures are high and daylight hours are long. Ice cellars in permafrost have long been used for storing food, but permafrost warming means that temperatures in the ice cellars are too high to store food safely and, in some cases, the cellars thaw out. Freezers running on energy generated from solar panels can be a potential sustainable alternative for these communities.

2 - Energy produced from the windmill is converted to electricity and hydrogen which allows for short- and long-term storage. The energy is delivered to a local mining industry when it needs to reduce its diesel consumption.

ACTIVE LAYER




Wind farms in permafrost regions face unique challenges. Constructing wind turbines requires heavy equipment that may damage the underlying permafrost. The turbines also require strong and stable foundations to counter the stress exerted on the structure when in operation. Problems can emerge as thawing permafrost weakens the integrity of the ground.

Numerous techniques have been developed to adapt wind turbines to permafrost landscapes. In southwestern Alaska, helical piers were dug 11–12 metres below the surface to anchor the steel and concrete foundations in deeper layers of the ground. In Nunavik, Canada, a pilot windmill built in 2014 used a pile-mounted foundation that extends below the active layer in a "spider-like steel foundation". In addition to stabilising the foundation for future permafrost thawing, the project also reported a 90 per cent reduction in concrete needed for the foundation, highlighting its benefits for remote Arctic areas. Other adaptation methods being used or tested include the use of heavier foundations to ensure stability, controllers that adapt turbine operations to changing ground conditions, and mechanisms to reduce structural vibrations.

Many permafrost landscapes are suitable for hydropower and changing climate conditions can make new areas attractive for development. Hydropower has already been extensively developed in high altitude alpine landscapes to harvest the energy of flowing water on steep gradients. Also, as precipitation is predicted to increase in the Arctic, more hydroelectric power can be harvested in the north. Planning for changing climate with altered hydrological and ground conditions and accounting for the impacts of enlarged water reservoirs on permafrost thaw should be central in the future development of hydropower in the Arctic and in high mountain regions.

When permafrost thaws, water can percolate through the previously frozen ground and enter the subsurface layers. Depending on factors such as whether the permafrost is continuous or discontinuous, this process can change the run-off and impact the water availability for hydropower plants. To plan for future changes, there is a need to use modelling of both climate and local hydrological structures. The development of such models also includes sharing and communicating results to relevant stakeholders involved in hydropower development, as well as conducting proper environmental impact assessments.

3 - In Nunavut, Canada, a village installed a solar-powered community freezer. It provides space to store harvested meat and fish, especially in the warmer and longer summer months. Reducing the reliance on diesel also provides an alternative to ice cellars.



No time to waste *Waste management and permafrost*

Contrary to the image of the Arctic as a pristine landscape, the region does have waste management issues. Arctic cities, for example, produce more waste per capita, recycle less, and have lower rates of waste collection than the global average. Waste can also be an issue for many Indigenous communities whose traditional lifestyles produce little inorganic waste but who, in more recent times, have adopted a more urban lifestyle. Proper waste disposal is important not only for the storage localities themselves but also for the health of the wider environment. Thawing permafrost is affecting storage facilities in many places, making this an urgent problem in need of further research.

There are significant differences in the types of waste produced between large urban centres in the Arctic and small remote communities. A large city, such as Anchorage, with a large population and diversified economy produces a larger variety of waste, including hazardous waste, compared with a small settlement, such as Unalaska, where the king crab fishery is the biggest economic activity. A study conducted by the Arctic Council's Sustainable Development Working Group (SDWG) identified several factors that influence successful waste management. These include accessibility, population size, and access to government support and funding. Connectivity to regional waste facilities serving a larger population, for example, allows access to equipment and maintenance which are not available to smaller communities without road access. Having an engaged community with local champions is also important for developing appropriate waste management systems, as is the support of regional and national governments for local communities in this task.


Landfills on permafrost need to be treated differently than those in warmer climates. Permafrost is sensitive to disturbance, so digging holes for landfills or removing the insulating cover will affect the permafrost stability. As such, above-ground freezeback landfills are preferred in many Arctic communities as this technique does not require digging deep into the ground. In freezeback landfills, the top waste will completely or partially freeze because of the cold climate and can thus be contained. Permafrost is also used as an impermeable bottom layer, preventing leakages into the surrounding environment.


The design of freezeback landfills vary. In the Alaska Administrative Code, for example, some landfills in permafrost regions can be exempted from certain regulatory rules such as having a liner (an impermeable barrier between the landfill and surrounding areas used to avoid leakages) and monitoring of groundwater impacts. This exemption is based on the premise that a freezeback design is developed where permafrost is maintained and that the waste will remain frozen during and after the lifetime of the landfill. Some landfills, however, cannot accomplish freezeback and in such instances a frozen barrier can be developed around the landfill.

While relying on permafrost as an impermeable layer might be a functional design in today's climate, increasing temperatures will take away this option in many places across the Arctic. If the underlying permafrost layer thaws, contaminants can seep into and pollute the surrounding environment, rivers, and groundwater. In addition, the waste itself can cause local thawing in and around waste landfills. The actual amount of heating depends on the type of waste that is stored. Dumps of coal fragments, for example, can produce heat due to oxidation, while municipal solid waste sites produce heat through decomposition. When the permafrost around a landfill thaws, unfrozen areas in the permafrost known as talik zones are created under the waste, and leakages from the dumps into water and the ground can occur. If taliks are allowed to develop, the ground can eventually collapse, forming thermokarst depressions. While some adaptation options exist (e.g., strengthening landfill embankments, introducing air convection embankments, and thermosyphons), better technologies are needed in this area.

Dumps and landfills on permafrost have several potential concerns associated with them. It is therefore important to ensure that they are properly constructed and managed to both contain waste securely and prevent permafrost thaw. To be cost-effective, governments can also apply land-use planning and risk maps as practical tools to locate and target the most hazardous areas in need of new engineering solutions.

Laws and regulations play an important role in ensuring proper waste management, but not all Arctic states have paid attention to the special conditions of permafrost landscapes. For example, the Russian strategy for household waste adopted in 2013 does not specifically mention permafrost regions. Developing such strategies will be beneficial in the long-term considering the pressure on permafrost landscapes.





SOURCES: NORWEGIAN DIRECTORATE FOR CULTURAL HERITAGE (2022), OBU ET AL. (2019). BY LEVI WESTERVELD | GRID-ARENDAL (2023)

Modern history *Preserving Svalbard's cultural heritage*

The Arctic region has a rich and diverse cultural history which has left behind a myriad of cultural heritage sites. Arctic culture is conserved through people who continue to practice their traditional and Indigenous livelihoods. These livelihoods and cultural heritage, however, are under increasing pressure from the impacts of climate change, among other threats. The foundations of older buildings and infrastructure are of various quality, and as permafrost thaws, cultural heritage sites are at risk of damage and potential collapse. While technological solutions are available to protect cultural heritage sites (e.g., reinforcing foundations or building protective barriers), there are limitations. Chief among these is the issue of cost, a problem exacerbated in the Arctic due to the high cost of transporting tools, materials, and personnel.

Svalbard and its population represent a unique context in the Arctic. It has a highly transient population with an average residency time of 7 years. Also, in contrast

Coal cable car-trestle

ACTIVE LAYER

PERMAFROST

to many other Arctic communities, the island was not permanently inhabited until mining operations started in the early twentieth century. This has resulted in a sense of place which is different from other parts of the Arctic where culture can be traced back through ancestral history. Cultural heritage sites, therefore, play an important role on Svalbard as the built environment helps preserve a particular historical and cultural identity. Svalbard's heritage sites hold international importance as they represent unique stories of human settlements.

The Svalbard Environmental Act states that all structures and sites dated before 1946 are automatically protected. The island has 8,300 officially registered cultural heritage sites spanning from buildings and infrastructure to graves and protected cultural environments. Researchers found that out of 872 surveyed sites, 44 were considered vulnerable to the formation of thermo-erosion gullies and therefore under threat from future climatic warming. The situation is further aggravated by other environmental stressors such as coastal erosion. Action is required to ensure that the sites remain intact over time. The Polar Climate and Cultural Heritage project highlights three values that compete in the current cultural heritage system on Svalbard. The antiquarian value is related to a conservationist attitude where authenticity and research interests are most important. A more pragmatic user value argues for an emphasis on the practical value of buildings and property. Finally, the experiential value addresses the holistic experience that both visitors and residents have when interacting with cultural heritage sites. These value systems can be complementary, although controversies sometimes arise. For example, while the actors with a user value may be interested in applying modern and efficient building techniques to protect and restore heritage sites, doing so could decrease their antiquarian value.

1 - The deepening of the active layer exposes the wooden trestle legs to moisture and higher temperatures, causing them to rot and fail.

2 - Maintaining cultural infrastructure sites involves drilling new and deeper holes into the permafrost for new foundations next to the original ones.

3 - The original cable-car trestles are then lowered back in their original positions into the new and more stable foundations.



PORTRAIT ADVISOR AT KINGS BAY AS, NY-ÅLESUND, NORWAY

Part of why I ended up in Ny-Ålesund was that I've always been a bit restless. I became interested in cultural monuments and what you can learn from them and decided to become an archaeologist. Cultural heritage is exciting because people are exciting. Cultural monuments can tell us how humans at different times and in different locations have used their intelligence, skills, and creativity to survive and give life meaning.

At 79 degrees north, Ny-Ålesund is the northernmost community in the world, and the last company town in Norway, run by Kings Bay AS. Today we run a research station, and in summer up to 150 people live there. But in the winter, we are only around 35 permanently employed people here. Ny-Ålesund was originally a mining community, founded in 1916, but coal production ended in the 1960s. The buildings here give us a window into the past and can tell us a lot about the hopes, creativity, and perseverance of the people who lived here before us. Miners came here in the fall, knowing they would be stuck in total darkness and isolation for 6 months. Witnessing how they were able to live in rather pitiful conditions, compared with the standards we have today, is inspirational.

I first encountered permafrost when I started working in Svea, another old coal mine in Svalbard. I quickly learned that it's not easy to dig in the ground. We had to wait until it was the right season or wait for weeks until the sun heated up the ground sufficiently before we could start digging decimetre by decimetre. Even large 30-ton excavators could not get through. During that time, I gained a practical understanding of concepts like the active layer. I was familiar with frozen ground from archaeology on the Norwegian mainland, but the frost here does not come from cold air above ground, but rather from below.


In my earlier work in Svea, I also learned a lot from more experienced colleagues, miners, and people that have lived on Svalbard for a long time. They told me how things were changing, and how the permafrost was more stable and reliable. All the buildings in Ny-Ålesund are built on permafrost, so we are dependent on it. When the permafrost thaws deeper and deeper every year, it impacts the buildings. The posts that the buildings are standing on are frostjacked, but often unevenly which further affects the stability of buildings. It can create a lot of issues, and changes have accelerated. And it's not only the houses. Suddenly a stretch of road disappears. Everything is suddenly a bit less stable. When you live 1,000 kilometres away from the nearest major settlement, you have to think about what that means for community safety and emergency preparedness.

In Ny-Ålesund, we have Svalbard's largest collection of protected buildings. We have 29 buildings under the strictest form of preservation built in 1909–1945, but a total of 40 buildings are considered to have cultural value. It's a lot of responsibility, because this is a cultural heritage on a national level from the mining period. When it comes to fixing the foundation of old buildings, we have to consider how big the historical value is when measured against the usefulness and economic cost of maintaining a building. We have to weigh advantages and disadvantages. Gradually, it's getting more and more expensive to deal with the impact of climate change. So how much time, money, and material should we use to save historical monuments? At some point we have to prioritise more.



Going South

Permafrost in Other Areas





Rock glaciers in the Andes are underlain by an ice-rich permafrost layer and are thought to be important seasonal contributors to stream flow and thus, water security.

D A N T A R C T I C A

South Pole

Permafrost occurs within most of the ice-free areas of the Antarctic, less than o.2 per cent of the continent. It might exist under the massive Antarctic ice sheet but there is no direct evidence of it.

SouthernOcean

A planetary perspective Permafrost outside the Arctic

Permafrost also occurs outside of the Arctic, in the ice-free areas of the Antarctic, and at high elevations around the globe where the temperature remains below o °C for much of the year. Permafrost occurs extensively in mountainous areas across the Qinghai-Tibetan Plateau, in the Hindu Kush Himalayan region, Central Asia, and Mongolia, as well as throughout the South American Andes, the European Alps, the Rocky Mountains of North America, and even on Hawai'i (not shown on map).

While there are some similarities between polar and mountain permafrost, there are also significant differences. Compared with the Arctic and Antarctic, for example, mountainous areas with permafrost show an extreme range in precipitation, from the very dry conditions of the Andes to the very wet environments found in Norway. Permafrost occurrence in mountain areas is strongly controlled by local topography and is most likely to occur in places that commonly lie in the shade, as well as in valleys with steep sides located at higher latitudes. These factors lead to a much higher degree of spatial heterogeneity (patchiness) in mountain areas with permafrost than in polar environments where permafrost is more often spatially continuous across vast areas.

Frozen giants *Permafrost in the mountains*

Outside of the Arctic, permafrost most commonly occurs in high mountaintops. It is most pronounced on the north-facing slopes in the northern hemisphere and south-facing slopes in the southern hemisphere. With decreasing elevation, permafrost becomes less extensive and more discontinuous, isolated, or sporadic until it disappears all together. In addition to latitude and altitude, the presence of permafrost is also determined by precipitation, seasonal and daily temperature extremes, air humidity, soil moisture, and geology.

Direct observations of permafrost cannot be made using remote sensing technology, and so permafrost mapping depends on sparse ground temperature measurements that are augmented by extrapolation and interpretation. Observational data, in turn, are unevenly distributed among and within different regions, and sampling strategies vary locally. For example, in the European Alps, permafrost is often observed at high elevations or on the summits of mountains, whereas on the Tibetan plateau sampling is mainly done for practical applications, such as along roadsides to monitor thawing.


These different observational techniques can bias analyses of permafrost occurrence. When combined with the environmental factors mentioned above, this leads to a high degree of uncertainty in models of permafrost occurrence in mountains, as well as in its response to global climate change. Regardless, evidence shows that higher temperatures resulting from anthropogenic global climate change are causing permafrost to thaw. Although measurements are sparse and unevenly distributed among mountain regions, existing data indicate warming has occurred by 0.14–0.24 °C per decade on average for 28 mountain sites in Asia, Canada, the European Alps, and Scandinavia.

As in the Arctic, the impacts of thawing mountain permafrost can be quite devastating. This thawing affects water quality, increases the risk of landslides, and causes damage to infrastructure. Permafrost acts as a barrier for water so that both rain and meltwater flow over the permafrost on the land surface, feeding streams and rivers. Permafrost thaw may affect the surface and subsurface water flow when the soil permeability increases and the active layer deepens. The changes in hydrological pathways can disrupt local agriculture and drinking water supplies. Water quality is also affected by changing stream flow rates and by the release of heavy metals, sediment, and contaminants from the thawing permafrost.

The thinning and thawing of permafrost make mountain slopes unstable, which in turn increases the risk of landslides as well as rock avalanches. Both landslides and rock avalanches can lead to damaged infrastructure and loss of life, especially in densely populated areas or areas with high tourist activity. Outburst floods and catastrophic floods resulting from the sudden release of a large volume of water can be triggered by permafrost thaw in mountain lake dams. Roads, railways, and buildings that stand on permafrost are also at risk as the ground becomes less stable with permafrost thaw. In fact, the very presence of roads and buildings can cause permafrost to thaw faster through disturbance of the soil and localised heating of the ground.

There are major gaps in our understanding of the occurrence of permafrost in mountain areas and of the impacts that anthropogenic global climate change will have upon communities located in mountain permafrost regions. Thawing permafrost is impacting mountain communities. These impacts are expected to become more pronounced and more frequent with the anticipated increase in global warming.

Permafrost distribution by elevation and latitude



NOTE: the graphic includes all modelled sporadic, discontinuous, and continuous permafrost data from Obu et al. (2019). SOURCES: OBU ET AL. (2019a), (2019b), (2020); SRTM (2015). BY LEVI WESTERVELD | GRID-ARENDAL (2023)

The view from the top

R

U

Altay Mountains

Ürümai

Plateau of Tibet

-Z Kathmandu

NEPAL

The Qinghai-Tibetan Plateau, Hindu Kush Himalaya, and Andes

S

The largest permafrost area outside of the Arctic is found in the Qinghai-Tibetan Plateau extending to the slopes of the Hindu Kush Himalaya to the south. About 40 per cent of the plateau is characterised by

Nur-Sultan

Bishkek

Kashgar

KYRGYZSTAN

KAZAKHSTAN

New Delhi

INDIA

SOURCE: OBU ET AL. (2019) A, B. BY LEVI WESTERVELD | GRID-ARENDAL (2023)

Tashkent

TAJIKISTAN

Dushanbe

Islamabad

PAKISTAN

Kabul

Surgut

Tian-Shan

-Mountains

Hotan

the presence of permafrost, making it the largest store of mountain or high-elevation permafrost carbon on Earth. Atmospheric warming is already affecting the permafrost of the Qinghai-Tibetan Plateau, with recent computer models predicting that climate change may reduce it by as much as 70 per cent by the year 2100.

A

Mongolian Plateau

Ham

Tanggula

Thasa

Thimphy

BHUTAN

🚊 Qinghai-Tibet railway

Ulaanbaatar

Chengdu

MONGOLIA

Jiayuguan

The Hindu Kush range is home to the world's highest mountains, including Mount Everest which reaches 8,849 metres above sea level. Large variations in topography combined with regional variations in air temperature lead to a large range in the elevation at which permafrost occurs. Permafrost can be found as low as 3,500 metres above sea level in the western Hindu Kush Himalaya, while bedrock without permafrost is likely to exist in south-facing slopes well over 6,000 metres above sea level in the central and eastern parts. Although data are still scarce, permafrost degradation has been linked to many hydrogeological and ecological changes, Yakutsk such as a decline in water

Beijing

Zhengzhou

N

Chongqing

Modelled permafrost coverage

> Continuous (>90%) Discontinuous (50–90%) Sporadic (10–50%)

conservation capacity, the shrinking of wetlands and meadows, and reduced grassland productivity, diversity, and vegetation coverage.

Across the region, thousands of kilometres of roads and railways have been built in permafrost areas and are now exposed to potential damage from permafrost thaw. In addition to climate warming, the engineering activities themselves can create thermal changes that lead to permafrost degradation and result in very high infrastructure maintenance and repair costs. About 550 kilometres of the 2,000-kilometre-long Qinghai-Tibet railway from Lhasa to Xining is built on permafrost, reaching over 5,000 metres above sea level in places. To increase railbed stability, embankments of loosely piled rocks have been built to allow air to circulate freely. In some stretches, ventilation pipes buried into the ground allow the cold air to circulate underneath the railbed. The cost of maintenance and repairs is expected to increase as projected climate warming deepens the active layer above permafrost on the plateau.

The Andes are the longest mountain range in the world, extending from tropical Venezuela to the southern tip of Argentina. Conditions favourable to the formation of mountain permafrost can be found across the highaltitude areas. It is challenging to predict where this permafrost might occur, as the complex mountain topography results in extremely variable patterns of permafrost formation. Field surveys that confirm the presence of permafrost are expensive and logistically difficult. More commonly, permafrost distribution is estimated using probability models based on environmental variables, including mean annual air temperature, incoming solar radiation, topography, and elevation. Active rock glaciers are also frequently used as indicators of permafrost presence. These distinct landforms occur because of the downslope movement, or creeping, of frozen ground and can be identified from aerial photographs and satellite imagery. Rock glaciers consist of a surface layer (the active layer) underlain by an ice-rich permafrost layer. The amount of ice in the permafrost layer is variable (10–45 per cent). The surface layer of ice, rock, and soil insulates the underlying frozen ground, making rock glaciers more resistant than ice glaciers to the impact of climate warming. The seasonal contribution of rock glaciers to stream flow, and thus water security, is thought to be important but is not well understood.

While rock glaciers and other permafrost features are less obviously affected by atmospheric warming than ice glaciers, there are signs that they too are degrading. In the Bolivian Andes, projected warming could result in the loss of 95 per cent of current permafrost (including almost all rock glaciers) by 2050, and up to 99 per cent by 2099. Climate warming can increase the thickness of the active layer, releasing water frozen in the upper permafrost. However, this is a slow process and does not result in a significant increase in seasonal flow from the active layer. Climate-induced permafrost degradation is also implicated in the downslope acceleration of some rock glaciers and an increase in rockslides and rock avalanches.


Permafrost in the Andes is also affected by mining and other forms of development that can increase the ground temperature, leading to subsidence and the formation of thermokarst. This is of serious concern because mining in the Andes is increasing, causing major changes in the landscape including the disposal of mine tailings and waste rock onto the surface of rock glaciers.



Europe's frozen heart *Permafrost in the Alps*

The European Alps are a densely populated and relatively accessible region in comparison to the other mountain regions of the world, making it a very popular destination for mountain tourism. Permafrost is a prominent feature in the Alps where it occurs above 2500-3000 metres elevation above sea level and underlies about 3 per cent of the surface area, with the largest portion in Switzerland, followed by Italy, Austria, and France.

Mountain permafrost is degrading due to climate warming, destabilising rock walls on the mountain slopes. In recent years, summer heat waves have triggered rockfalls in the Alps. Permafrost degradation



has an increasing impact on mountaineering practices and mountain tourism in general as climbing seasons have shifted and many popular mountaineering and climbing routes have become more dangerous.

In contrast to the Andes or the Hindu Kush Himalaya, rockfalls in the Alps are small but frequent during summer. Although relatively rare, major rock avalanches also occur. The Drus (3,754 metres) mountain in the Mont Blanc massif of the French Alps has been the site of major rockfall events, which scientists attribute to a warming climate and permafrost degradation. In June 2005, the 700-metrehigh Bonatti Pillar collapsed from the Drus, bringing down almost 300,000 cubic metres of rock, the equivalent to 5,200 large shipping containers. Between October 2005 and September 2016, scientists used laser technology to scan the granitic rock face. During the study period, researchers mapped over 300 smaller individual rockfalls, ranging from 0.002 to 476 cubic metres. The detailed records of rockfall frequency and volume allow researchers to better understand the role of permafrost thaw on rockfall in mountainous areas.

The European Alps are home to much permafrost research that has produced many unique permafrost data sets. These include: the longest time series of borehole temperatures in Europe from the Murtèl-Corvatsch rock glacier in the Swiss Alps, which started in 1987; annual terrestrial laser scans showing in detail how a mountain wall in Mont Blanc massif has changed over 10 years; and a long-term, large permafrost data series from the Matterhorn, one of the highest peaks in Europe. Adaptation strategies for mountain guides have also been developed in response to the changing mountain environment. These kinds of diverse data are invaluable for understanding mountain permafrost degradation and its diverse impacts on the environment and mountain tourism.



The ends of the Earth I Permafrost in Antarctica

Covered by the largest ice mass on the planet, Antarctica is the coldest continent. The continental ice sheet, which on average is more than 2 kilometres thick, insulates the ground surface from low air temperatures and leaves less than 0.2 per cent of the surface area ice free, mostly in the coastal areas at the edges of the ice sheet and where mountains protrude above the ice.


There are fewer permafrost monitoring boreholes in Antarctica than in the Arctic, but field observations and computer modelling show that permafrost is present under almost all the ice-free areas. Permafrost might exist under the Antarctic ice sheet, especially in areas where the ice cover is thin, but as there is no direct evidence of it, very little is known about the subglacial permafrost.

Climate and surface topography vary greatly from place to place in Antarctic permafrost areas. There is a huge latitude difference, as well as elevation range in the areas with permafrost. Ice-free areas range from the sea level to the top of the highest mountain, Mount Vinson in West Antarctica at 4,892 metres above sea level. Air temperature and the distribution of snow on the ground, the two factors affecting Antarctic permafrost temperatures most strongly, vary greatly across the continent. Antarctica is not only the coldest but also the driest continent in the world, despite many areas experiencing increased precipitation under climate change. Even if mean annual temperatures remain well below o °C across Antarctica, some coastal areas, especially on the Antarctic Peninsula, are warming rapidly with a tendency to above freezing temperatures.

Ground temperature as an indicator of permafrost existence has only been systematically monitored for the last two decades in Antarctica, and direct permafrost data for the continent are scarce. Sparse measurements give little information about spatial variability, and one must rely on computer models to better understand the ground temperature variability at the regional and continental scales. Modelling indicates that the coldest near surface permafrost in the world can be found at Mount Markham in the Transantarctic Mountains where the lowest ground temperature is estimated to be -36 °C.

There is no permanent human population in Antarctica. However, there are more than 80 research bases there. The number of people living and working on the continent increases to about 5,000 in the austral summer (December-February), while about 1,000 people stay over during the dark and cold winters. Many of the research stations are built in coastal areas on permafrost, where its thaw can create a potential risk for the infrastructure. Because annual mean ground temperatures are still very low, most buildings and other infrastructures are safer from the impacts of thawing permafrost than those in the Arctic. However, in some locations on the Antarctic Peninsula, permafrost thaw is having an impact on infrastructure, a problem that may become much more common under a warmer climate.

Permafrost thaw has local impacts, not only on research infrastructure but also on local hydrology. As the ground thaws, meltwater or rain will partially infiltrate the ground rather than flowing across its surface. As water is the most important physical driver of Antarctic terrestrial communities, this bears consequences for the movement of nutrients in the environment and the potential impacts on ecosystems and biodiversity. Pollutants accumulated in the permafrost may also be remobilised into the environment as a result of permafrost thaw, although more research is needed to understand their movement and environmental impacts. Unlike the Arctic, there will be lower greenhouse gas emissions from Antarctic permafrost because it contains very little organic matter. The terrestrial Antarctic could even become a short term carbon sink in the future, should vegetation expand with increasing temperatures.







The ends of the Earth II Antarctic Peninsula

The highest modelled near-ground surface temperatures in Antarctica are found on the Antarctic Peninsula, the northernmost part of the continent. The climatic conditions vary strongly across the peninsula, where the east coast is influenced by a continental climate, while the central and western parts experience maritime conditions. Very narrow coastal areas on the Antarctic Peninsula have sporadic or discontinuous permafrost, and the highest ground surface temperatures are found on the north-western areas of the peninsula. As elsewhere, air temperature and thickness of the snow cover play an important role for the existence of permafrost in the ice free areas.

As in other locations, thawing permafrost in the Antarctic poses a threat to infrastructure. The Marambio airstrip, which serves an Argentinian research station on Seymour Island in the Antarctic Peninsula, is built on permafrost. In 2020, the air temperature at the Argentine research base Esperanza, not far from Seymour Island, reached 18.4 °C, the highest temperature ever recorded in Antarctica, confirmed by the World Meteorological Organization. During this exceptional heat wave, flights to Marambio were impacted because the permafrost had started to thaw, transforming the surface of the airstrip to a deep watery mud which was not safe to land on. Meteorological observations on the peninsula over recent decades show an overall increasing trend in the number of such extreme warm days. Knowledge and understanding of these kinds of extreme events is important for the maintenance and safety of infrastructure such as airstrips and research stations on the continent.






The ends of the Earth III Queen Maud Land, Victoria Land, and the McMurdo Dry Valleys


In Antarctica, cold and very dense air blows from the higher interior plateau down the steep slopes along the coast due to gravity. This process generates katabatic winds, which can be very strong at times. These winds blow down from the East Antarctic Plateau and keep large parts of the mountain ranges dry and free of snow in Queen Maud Land. The ice-free and snow-free areas of Antarctica are often referred to as oases. One of these, the Schirmacher Oasis, is an ice-free area that is 25 kilometres long and up to 3 kilometres wide, encompassing more than 100 freshwater lakes. In addition to the Indian research station Maitri located in the Schirmacher Oasis, there are 11 other research stations in Queen Maud Land, most of them built on rocky outcrops of the mountain chain where high winds keep the ground snow free.

Queen Maud Land is one region of Antarctica where an increasing trend of snow accumulation has been observed, along with rising temperatures. On the opposite side of the continent, the McMurdo Dry Valleys are among the driest areas in the world, generally classified as hyper-arid polar deserts. It is the only location on Earth known to contain dry permafrost consisting of loose soil, because its water content is very low and insufficient to cement the soil particles together. This is also the location where the oldest permafrost on Earth has been found.

The most extreme climatic conditions prevail inland at higher elevations, such as the University Valley. The



conditions resemble those on Mars, making the Dry Valleys a Mars analogue where scientists can study periglacial processes and geomorphic features and their use as a diagnostic for subsurface ice on Mars. In contrast to the Arctic, where active microbial populations can be found in permafrost soils, microbial activity is undetectable at the high elevation locations of the Dry Valleys. This lack of activity indicates the limits of life in terrestrial environments, carrying potential implications for life in extraterrestrial permafrost environments.





Over the Horizon

Climate change is having a profound effect on every part of the globe, but its effects are especially pronounced on the Earth's cryosphere. Readily observed and measured, the changes in snow and ice on land, glaciers, ice caps, and sea ice have long been evident. The effects on permafrost, however, are less conspicuous and have received less attention beyond the scientific community and Arctic residents. Nevertheless, permafrost is rapidly changing as a direct consequence of climate change, posing risks to Arctic peoples and the environment, and with significant implications for the entire planet.

The knowledge compiled in this atlas demonstrates how vital permafrost is for regulating the climate, maintaining unique ecosystems, supporting infrastructure, influencing hydrology, and sustaining the livelihoods of Indigenous communities.

- Climate regulation and climate science: Permafrost contains vast amounts of plant and animal matter, keeping them frozen for tens, even hundreds of thousands of years. When this organic matter thaws, it decomposes and releases carbon dioxide (CO₂) and methane (CH₄), both potent greenhouse gases. The release of these gases exacerbates global warming and climate change, contributing to a positive feedback loop that will last for centuries to come.
- Health: In addition to organic carbon, permafrost also stores contaminants. For example, permafrost soils store nearly twice as much mercury as all other soils, the ocean, and the atmosphere combined. As permafrost continues to thaw over the next century, this mercury becomes vulnerable to release into the environment. Other contaminants, including heavy metals, persistent organic pollutants,

and microbiological agents, are also locked in permafrost, as long as it stays frozen. However, when the permafrost thaws, these pose a risk for both human and animal health.

- Destabilisation and erosion: Permafrost thaw dramatically destabilises the ground. Coastal communities see their shorelines crumble away, lakes suddenly drain, and infrastructure threatened. This destabilisation carries costs for human and animal health, the economy, and society. The costs for infrastructure damage alone could exceed USD 276 billion over the next 40 years.
- *Ecosystem services and biodiversity:* Specialised flora and fauna adapted to cold and frozen conditions make permafrost their home. These ecosystems provide a habitat for species that are uniquely adapted to permafrost environments and contribute to global biodiversity. As permafrost changes, lakes may drain and disappear, ice-rich coastlines erode, and tundra vegetation is replaced by shrubs and trees. These and other habitat changes can have serious consequences for the species dependent upon them.
- *Water resources and quality:* Permafrost plays a role in regulating regional hydrology. It can act as a barrier to water movement and influence the flow of rivers and streams. Some permafrost regions store significant amounts of fresh water in the form of ice. As permafrost thaws, it can release this water, potentially impacting local water resources, and even sea levels.
- Indigenous communities: Indigenous Peoples
 possess unique knowledge and cultural practices
 associated with permafrost environments, and
 changes in permafrost conditions can profoundly
 affect their ways of life. Understanding the

impacts of ongoing change and determine the best adaptation strategies requires a respectful exchange between science practitioners and holders of local and traditional knowledge.

• *Earth's history:* Studying permafrost cores can provide insights into the past climate and provide information on the resilience of permafrost during warmer periods. This information can teach us about natural rates and consequences of change, unaffected by human agency.

Commonly, the social, physical, and health components of permafrost thaw have been studied in isolation, leading to inadequate policy options that ignore the holistic nature of the threat. For instance, contaminant release directly affects wildlife, which itself is harvested for food. Infrastructure failure directly threatens water and sanitation in communities, drastically impacting health and well-being. There is a need for an integrated and participatory approach to the complex issues at the nexus of climate change, permafrost thaw, infrastructure damage, contaminants, health and well-being, and for solutions founded on the cultural, natural, and social frameworks of local communities.

As climate change continues to affect permafrost regions, understanding and managing these ecosystems becomes increasingly important for the well-being of our planet and its inhabitants. This is also a time to rethink permafrost science, to foster co-design, and to share participation in observation, knowledge production, and their benefits. Some outstanding research topics are:

• Current projections of the impacts of permafrost thaw on the carbon cycle leave out important

processes and links. There is a need to incorporate the currently observed mobilisation of carbon on land and beneath the ocean into our models of change to better understand the multi-generational challenges of long-term feedbacks. Because permafrost thaw may trigger carbon loss that continues for centuries, these future emissions also need to be accounted for in modelling.

- Permafrost thaw, in its progressive and abrupt forms, will release pollutants, but the magnitude and the timing of the release remains unquantified.
- A new approach, including permafrost models projecting infrastructure lifetime into the twentysecond century, is needed to provide long-term vision and decision-making capacity to planners.
- There is a pressing need to evaluate the risks to wildlife and human health that result from changes in water quality due to permafrost thaw and to involve communities in developing nature-based solutions to improve their well-being.
- The interaction between human activities, climate change, and permafrost is creating numerous hazards which can have significant environmental, sociocultural, economic, and health-related impacts on Arctic communities. Greater effort is needed to develop adaptation, mitigation, and recovery measures that can be implemented in remote regions.
- A targeted effort to map out the interconnected impacts of permafrost thaw on One Health is needed to identify risks and to prevent and manage health risks in humans, animals, and our shared environment.
- Arctic Permafrost Ecosystem Services (APES) need to be identified and classified to underscore the global value of permafrost.

This atlas appears at a time when humanity's sense of permanence is uncertain. Human activity is pushing the limits of the planet to support us. It is in our best interest to preserve as much permafrost as possible. Achieving this requires a global societal transformation, breaking from our dependence on fossil fuels and reducing greenhouse gas emissions.

For those living on permafrost, urgent strategies are needed to cope with the ongoing changes. Once adapted to life on permafrost, people now face the need to adapt to a destabilised environment where homes, businesses, and transportation infrastructure are all at risk. Where permafrost has been actively relied on to restrict contaminants and pathogens, new solutions are required as permafrost thaw deepens and containment fails.

The most significant long-term impact of permafrost thaw for the global community is the substantial release of greenhouse gases for centuries to come, necessitating compensation by future generations. The duration and extent of this are still under investigation but from a permafrost perspective, current warming is already too much. Urgent action is needed to avert as much permafrost thaw as possible, and it is not too late if we act now. By significantly reducing our greenhouse gas emissions, we can protect the global climate from generations of uncontrollable permafrost emissions and work towards positive change for permafrost and the planet as a whole.

Glossary

- **Abrupt thaw** The rapid thaw of permafrost, usually by erosion, landslide or some other catastrophic process
- **Active layer** The top layer of permafrost soil that freezes and thaws

Active layer detachment slides – Slope failure(s) in permafrost areas, in which the active layer detaches from the frozen ground beneath and slides downslope

Aerobic – In the presence of oxygen, usually because air is available

Alaas – A depression or drained thaw lake typically covered by grasslands resulting from thawing of ice-rich permafrost

- **Anaerobic** In the absence of oxygen, usually because all oxygen has been consumed and air is not available
- **Arctic amplification** The phenomenon of faster warming in Arctic regions compared to the global average
- **Benthic** The bottom of a water body, such as an ocean, lake or river
- Bioaccumulate The gradual building up of a (harmful) chemical in an organism. In food chains, chemicals tend to become more highly concentrated in organisms higher up the chain
- **Carbon** The main element of life on Earth and the main element in greenhouse gases. It occurs in organic and inorganic compounds, mostly based on whether the carbon is bonded with hydrogen (as in most life-forms) or not (for example in minerals)
- Carbon cycle The movement of carbon between the different reservoirs (land, ocean, atmosphere) of the Earth
- **Carbon sequestration** The capturing and storing of carbon dioxide from the atmosphere

Carbon sink – Anything that absorbs and accumulates more carbon than it emits

- **Climate scenario** Results of computer modelling of the Earth's future climate that include estimates of future human greenhouse gas emissions. There are often called future "projections" or "pathways", and include low, moderate and high emission alternatives. Projections for future greenhouse gas emissions and their impacts on the climate, sometimes categorised into low/intermediate/high depending on the rate of temperature increase
- **Continuous permafrost** Permafrost that underlies 90 per cent or more of the ground in a region
- **Convection** A circular flow in gas or liquid that results from density differences, for example when warm water rises and cold water descends
- Discontinuous permafrost Permafrost that underlies 50-90 per cent of the ground in a region

Earth System Model – A computer model of the physical, chemical and biological processes of the Earth system

Erosion gullies – Trenches formed as soil material is washed away.
Positive feedback loop – A pair of processes, in which the result reinforces the cause

Frost heaving – The lifting of the ground surface during freezing *Glacial period* – see definition for Last Glacial Maximum *Gradual thaw* – The slow thawing of the permafrost by warming

Greenhouse gases – Chemicals in the atmosphere that absorb

- heat reflected from Earth's surface, trapping it and warming the atmosphere; the main greenhouse gases are water vapour, methane (CH₄, an organic compound) and carbon dioxide (CO₂, an inorganic carbon compound)
- **Ground subsidence** Sinking or a downward movement of Earth's surface; in permafrost regions, subsidence can result from the thaw of ground ice

High Arctic – The northernmost part of the Arctic, defined by its climate and vegetation

Hydraulic conductivity – The relative ease of which a fluid can pass through soil or rock

Ice wedge – A body of ice that forms when vertical cracks open in the ground during winter and subsequently fill with spring meltwater. As the water freezes, a vertical vein of ice is created which becomes wider every year as the process repeats itself, typically widest at the top. If several ice wedges grow large enough, and join, they create ice wedge polygons

Interglacial period – A geological period during which warm air temperatures shrink glaciers and ice caps. An interglacial period usually comes between colder glacial periods during which the ice masses grow.

Isolated permafrost – Permafrost that underlies less than 10 per cent of the ground in a region

Last Glacial Maximum – The geological period during which glaciers were at their largest around 25,000 years ago

Linear infrastructure – Features built along a continuous line, such as roads, railways, utilidors and pipelines

Methanotrophic microbes – Microorganisms such as bacteria or archaea that use methane as their source of carbon and chemical energy

Negative feedback loop – A pair of processes, in which the result dampens the cause

Outburst flood – A sudden release of a large volume of water for example from a naturally dammed lake. Such floods are not frequent but can cause a lot of damage

Permafrost – Ground with a temperature at or below o °C for at least two consecutive years.

- **Pingo** Large, roughly conical hill with an ice core, covered with soil and vegetation
- **Point infrastructure** Built features that are not connected, such as an airport, drilling pad or harbour
- **Primary production** Creation of chemical energy for living organisms from inorganic substrate, for example via photosynthesis
- **Retrogressive thaw slumps** A permafrost landform caused by the thawing of ice-rich permafrost, leading to a slope collapse and retreating headwall
- **Rock glacier** A permafrost landform consisting of rock debris frozen in or on top of ice
- **Sporadic permafrost** Permafrost that underlies 10-50 per cent of the ground in a region
- Talik An area of unfrozen ground within permafrost
- **The Paris Agreement** An international treaty adopted in 2015 by 194 countries. The treaty has a long-term goal to keep the mean global temperature warming well below 2 °C above preindustrial levels
- Thermal abrasion The destruction of frozen coastline by the mechanical and thermal action of waves
- **Thermal denudation** Erosion of shorelines caused by warming and thawing of permafrost above the water line
- **Thermokarst** Depressions in the landscape formed by permafrost thawing and ground ice melting, which lowers the ground surface (see ground subsidence). Such depressions may fill with water to form thermokarst ponds or lakes
- Thermosyphon A passive heat exchange device utilising convection to cool the ground
- *Turbidity* A measure of the relative clarity of a liquid
- **Yedoma** A carbon-rich and ice-rich class of permafrost deposits found in Siberia (Russia), and parts of Alaska (United States) and Yukon (Canada)
- **Zoonoses** Infectious diseases, also known as zoonotic diseases, that have jumped from animals to humans

Acronyms

- ACD Arctic Coastal Dynamics
- ACE Air Convection Embankment
- ACPR Arctic Circumpolar permafrost region
- AMAP Arctic Monitoring and Assessment Programme
- APPVI Arctic Permafrost Population Vulnerability Index
- AR6 Sixth Assessment Report
- CALM Circumpolar Active Layer Monitoring
- GRP Global Regional Product
- Gt Gigaton (109 t, 1015 g) = a billion tonnes
- GTN-P Global Terrestrial Network for Permafrost
- HKH Hindu Kush Himalaya
- IPA International Permafrost Association
- IPCC Intergovernmental Panel on Climate Change
- IPY International Polar Year
- NDVI Normalised Difference Vegetation Index
- NSIDC National Snow and Ice Data Center
- Pg Petagram (1015 g) = a billion tonnes
- POPs Persistent organic pollutants
- RCP Representative Concentration Pathway
- SDWG Arctic Council's Sustainable Development Working Group
- Tg Teragram (1012 g) = a million tonnes
- UNEP United Natio ns Environment Programme
- UNFCCC United Nations Framework Convention on Climate Change
- WMO World Meteorological Organization
- WWF World Wildlife Fund

References

Foreword

- European Commission and High Representative of the Union for Foreign Affairs and Security Policy (2021). Joint Communication to the European Parliament, the Council, the European Economic and Social Committee and the Committee of the Regions. A Stronger EU Engagement for a Peaceful, Sustainable and Prosperous Arctic (JOIN(2021) 27 Final), Brussels, 13.10.2021. https://www.eeas.europa.eu/sites/ default/files/2_en_act_part1_v7.pdf. Accessed 9 June 2023.
- Schuur, E.A.G., Abbott, B.W., Commane, R., Ernakovich, J., Euskirchen, E., Hugelius, G. et al. (2022). Permafrost and climate change: Carbon cycle feedbacks from the warming Arctic. Annual Review of Environment and Resources 47, 343–371. https://www.annualreviews.org/doi/10.1146/ annurev-environ-012220-011847.

Prologue

- Arctic Monitoring and Assessment Programme (2021). AMAP Arctic Climate Change Update 2021: Key Trends and Impacts. AMAP, Tromsø, Norway. viii+148. https://www.amap.no/ documents/doc/amap-arctic-climate-change-update-2021key-trends-and-impacts/3594.
- Intergovernmental Panel on Climate Change (2023). Sections. In Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Core Writing Team, Lee, H. and Romero, J. (eds.). Geneva, Switzerland. 35-115. https://www.ipcc.ch/report/ar6/syr/.
- Rantanen, M., Karpechko, A.Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K. et al. (2022). The Arctic has warmed nearly four times faster than the globe since 1979. Communications Earth & Environment 3(168). https://www. nature.com/articles/s43247-022-00498-3.

Earth's Freezer

Frozen grounds

Text

- Dobinski, W. (2011). Permafrost. Earth-Science Reviews 108(3– 4), 158–169. https://www.sciencedirect.com/science/article/ abs/pii/S0012825211000894.
- Obu, J. (2021). How much of the Earth's surface in underlain by permafrost? Journal of Geophysical Research: Earth Surface

126(5), e2021JF006123. https://agupubs.onlinelibrary.wiley. com/doi/10.1029/2021JF006123.

- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km2 scale. Earth-Science Reviews 193, 299–316. https://www.sciencedirect.com/science/article/ pii/S0012825218305907.
- Overduin, P.P., von Deimling, T.S., Miesner, F., Grigoriev, M.N., Ruppel, C., Vasiliev, A. et al. (2019). Submarine permafrost map in the Arctic modeled using 1-D transient heat flux (SuPerMAP). Journal of Geophysical Research: Oceans 124(6), 3490–3507. https://agupubs.onlinelibrary.wiley.com/doi/ full/10.1029/2018JC014675.
- Smith, S.L., Romanovsky, V.E., Lewkowicz, A.G., Burn, C.R., Allard, M., Clow, G.D. et al. (2010). Thermal state of permafrost in North America: A contribution to the International Polar Year. Permafrost Periglacial Processes 21(2), 117–135. https:// onlinelibrary.wiley.com/doi/10.1002/ppp.690.
- Van Everdingen, R.O. (ed.) (2005). Multi-language glossary of permafrost and related ground-ice terms. Calgary, Canada: International Permafrost Association. https:// globalcryospherewatch.org/reference/glossary_docs/ Glossary_of_Permafrost_and_Ground-Ice_IPA_2005.pdf.
- Way, R.G. (2018). Permafrost, 4 June. https:// indigenouspeoplesatlasofcanada.ca/article/tundra-andpermafrost/. Accessed 9 May 2023.

Maps

Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi.org/10.1016/j.earscirev.2019.04.023.
Overduin, P.P., Schneider von Deimling, T., Miesner, F., Grigoriev, M.N., Ruppel, C., Vasiliev, A. et al. (2019). Submarine permafrost map in the Arctic modeled using 1-D transient heat flux (SuPerMAP). Journal of Geophysical Research: Oceans 124(6), 3490–3507. https://doi.org/10.1029/2018JC014675.

Permafrost in profile

Text

Christiansen, H.H., Matsuoka, N. and Watanabe, T. (2016). Progress in understanding the dynamics, internal structure and palaeoenvironmental potential of ice wedges and sand wedges. Permafrost and Periglacial Processes 27(4), 365–376. https://onlinelibrary.wiley.com/doi/10.1002/ppp.1920.

- Dobinski, W. (2011). Permafrost. Earth-Science Reviews 108(3– 4), 158–169. https://www.sciencedirect.com/science/article/ abs/pii/S0012825211000894.
- Godin, E. and Fortier, D. (2012). Geomorphology of a thermoerosion gully, Bylot Island, Nunavut, Canada. Canadian Journal of Earth Sciences 49(8), 979–986. https:// cdnsciencepub.com/doi/10.1139/e2012-015.
- GRID-Arendal (2020). Coastal and offshore permafrost in a changing Arctic, 13 October. https://storymaps.arcgis.com/ stories/c163de04de7849cdb917fee88015dd73. Accessed 9 June 2023.
- Mackay, J.R. (1998). Pingo growth and collapse, Tuktoyaktuk Peninsula Area, Western Arctic Coast, Canada: A long-term field study. Géographie Physique et Quaternaire 52(3), 271–323. https://www.erudit.org/en/journals/gpq/1998-v52n3-gpq153/004847ar/.
- Olefeldt, D., Goswami, S., Grosse, G., Hayes, D., Hugelius, G., Kuhry, P. et al. (2016). Circumpolar distribution and carbon storage of thermokarst landscapes. Nature Communications 7, 13043. https://www.nature.com/ articles/ncomms13043.
- Paull, C.K., Ussler, W., Dallimore, S.R., Blasco, S.M., Lorenson, T.D., Melling, H. et al. (2007). Origin of pingo-like features on the Beaufort Sea shelf and their possible relationship to decomposing methane gas hydrates. Geophysical Research Letters 34(1), L01603. https://agupubs.onlinelibrary.wiley. com/doi/full/10.1029/2006GL027977.
- Prowse, T.D. and Brown, K. (2010). Appearing and disappearing lakes in the Arctic and their impacts on biodiversity. In Arctic Biodiversity Trends 2010: Selected Indicators of Change. Kurvits, T., Alfthan, B. and Mork, E. (eds.). Akureyri, Iceland: Ásprent Stell. Chapter 13. 68–70. https://www.caff.is/assessment-series/162-arctic-biodiversity-trends-2010-selected-indicators-of-change.
- United States of America, National Park Service (2017). Ice Wedges in Alaska's National Parks. [online video]. 9 December. https://www.youtube.com/watch?v=-_ IHbqLz530&ab_channel=AlaskaExtreme. Accessed 9 May 2023.
- Van Everdingen, R.O. (ed.) (2005). Multi-language glossary of permafrost and related ground-ice terms. Calgary, Canada: International Permafrost Association. https:// globalcryospherewatch.org/reference/glossary_docs/ Glossary_of_Permafrost_and_Ground-Ice_IPA_2005.pdf.

Frozen in time

Text

- Angelopoulos, M., Overduin, P.P., Miesner, F., Grigoriev, M.N. and Vasiliev, A.A. (2020). Recent advances in the study of Arctic submarine permafrost. Permafrost and Periglacial Processes 31(3), 442–453. https://onlinelibrary.wiley.com/ doi/full/10.1002/ppp.2061.
- Hughes, P.D., Gibbard, P.L. and Ehlers, J. (2013). Timing of glaciation during the last glacial cycle: Evaluating the concept of a global 'Last Glacial Maximum' (LGM). Earth-Science Reviews 125, 171–198. https://www.sciencedirect.com/ science/article/abs/pii/S0012825213001190.
- Osterkamp, T.E. and Burn, C.R. (2003). Permafrost. In Encyclopedia of Atmospheric Sciences Holton, J.R. (ed.). Academic Press. 1717–1729. https://www.sciencedirect.com/ science/article/abs/pii/B0122270908003110.
- National Snow and Ice Data Center (2020). Glaciers Quick Facts. https://nsidc.org/learn/parts-cryosphere/glaciers/glacierquick-facts. Accessed 9 May 2023.
- Vandenberghe, J., French, H.M., Gorbunov, A., Marchenko, S., Velichko, A.A., Jin, H. et al. (2014). The Last Permafrost Maximum (LPM) map of the Northern Hemisphere: Permafrost extent and mean annual air temperatures, 25–17 ka BP. Boreas 43(3), 652– 666. https://onlinelibrary.wiley.com/doi/10.1111/bor.12070.

Maps

- Ehlers, J., Gibbard, P.L. and Hughes, P.D. (eds.) (2011). Quaternary Glaciations – Extent and Chronology. Amsterdam:
- Elsevier. https://booksite.elsevier.com/9780444534477/. Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G. et al. (2007). Orbital and millennial Antarctic climate variability over the past 800,000 years. Science 317(5839), 793–796. https://doi.org/10.1126/science.1141038.
- Lindgren, A., Hugelius, G., Kuhry, P., Christensen, T.R. and Vandenberghe, J. (2015). GIS-based maps and area estimates of northern hemisphere permafrost extent during the Last Glacial Maximum. Permafrost and Periglacial Processes 27(1), 6–16. https://doi.org/10.1002/ppp.1851.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi. org/10.1016/j.earscirev.2019.04.023.
- Overduin, P.P., Schneider von Deimling, T., Miesner, F., Grigoriev, M.N., Ruppel, C., Vasiliev, A. et al. (2019). Submarine

permafrost map in the Arctic modeled using 1-D transient heat flux (SuPerMAP). Journal of Geophysical Research: Oceans 124(6), 3490–3507. https://doi.org/10.1029/2018JC014675.

An icy balance

Text

- Canada, Northwest Territories Geological Survey (no date). Permafrost thaw causes lake drainage and flow of debris, Peel Plateau, Northwest Territories, Canada. https://www. nwtgeoscience.ca/services/permafrost-thaw-slumps/videopermafrost-thaw-causes-lake-drainage-peel-plateau-nwt. Accessed 9 May 2023.
- Forbes, D.L., Craymer, M.R., James, T.S. and Whalen, D. (2022). Subsidence drives habitat loss in a large permafrost delta, Mackenzie River outlet to the Beaufort Sea, western Arctic Canada. Canadian Journal of Earth Sciences 59(11), 914–934. https://cdnsciencepub.com/doi/pdf/10.1139/cjes-2021-0127.
- Forbes, D.L., Manson, G.K., Whalen, D.J.R., Couture, N.J. and Hill, P.R. (2014). Coastal products of marine transgression in cold-temperate and high-latitude coastal-plain settings: Gulf of St Lawrence and Beaufort Sea. Geological Society, London, Special Publications 388, 131–163. https://www. lyellcollection.org/doi/10.1144/SP388.18.
- Jones, B.M., Grosse, G., Farquharson, L.M., Roy-Léveillée, P., Veremeeva, A., Kanevskiy, M.Z. et al. (2022). Lake and drained lake basin systems in lowland permafrost regions. Nature Reviews Earth & Environment 3, 85–98. https://www. nature.com/articles/s43017-021-00238-9.
- Kokelj, S.V., Kokoszka, J., van der Sluijs, J., Rudy, A.C.A., Tunnicliffe, J., Shakil, S. et al. (2021). Thaw-driven mass wasting couples slopes with downstream systems, and effects propagate through Arctic drainage networks. The Cryosphere 15(7), 3059–3081. https://tc.copernicus.org/articles/15/3059/2021/.
- Liljedahl, A., Boike, J., Daanen, R.P., Fedorov, A.N., Frost, G.V., Grosse, G. et al. (2016). Pan-Arctic ice-wedge degradation in warming permafrost and its influence on tundra hydrology. Nature Geoscience 9, 312–318. https://www.nature.com/ articles/nge02674.
- Nitze, I., Grosse, G., Jones, B.M., Romanovsky, V.E. and Boike, J. (2018). Remote sensing quantifies widespread abundance of permafrost region disturbances across the Arctic and Subarctic. Nature Communications 9, 5423. https://www. nature.com/articles/s41467-018-07663-3.
- Olefeldt, D., Goswami, S., Grosse, G., Hayes, D., Hugelius, G., Kuhry, P. et al. (2016). Circumpolar distribution and carbon

storage of thermokarst landscapes. Nature Communications 7, 13043. https://www.nature.com/articles/ncomms13043.

- Prowse, T.D. and Brown, K. (2010). Appearing and disappearing lakes in the Arctic and their impacts on biodiversity. In Arctic Biodiversity Trends 2010: Selected Indicators of Change. Kurvits, T., Alfthan, B. and Mork, E. (eds.). Akureyri, Iceland: Ásprent Stell. Chapter 13. 68–70. https://www.caff.is/assessment-series/162arctic-biodiversity-trends-2010-selected-indicators-of-change.
- Streletskiy, D. (2021). Permafrost degradation. In Snow and Ice-Related Hazards, Risks, and Disasters. Haeberli, W. and Whiteman, C. (eds.). Amsterdam, Netherlands, Oxford, UK and Cambridge, US: Elsevier. Chapter 10. 297–322. https://www.sciencedirect.com/science/article/abs/pii/ B9780128171295000214.
- Van Everdingen, R.O. (ed.) (2005). Multi-language glossary of permafrost and related ground-ice terms. Calgary, Canada: International Permafrost Association. https:// globalcryospherewatch.org/reference/glossary_docs/ Glossary_of_Permafrost_and_Ground-Ice_IPA_2005.pdf.

Maps

Brown, J., Ferrians, O., Heginbottom, J.A. and Melnikov, E. (2002). Circum-Arctic Map of Permafrost and Ground-Ice Conditions, Version 2. NASA National Snow and Ice Data Center Distributed Active Archive Center. https://nsidc.org/ sites/default/files/ggd318-v002-userguide.pdf. Accessed 20 June 2023. DOI: 10.7265/SKBG-KF16

What lies within

- Angelopoulos, M., Overduin, P.P., Miesner, F., Grigoriev, M.N. and Vasiliev, A.A. (2020). Recent advances in the study of Arctic submarine permafrost. Permafrost and Periglacial Processes 31(3), 442–453. https://onlinelibrary.wiley.com/ doi/full/10.1002/ppp.2061.
- Biskaborn, B.K., Lanckman, J.-P., Lantuit, H., Elger, K., Streletskiy, D.A., Cable, W.L. et al. (2015). The new database of the Global Terrestrial Network for Permafrost (GTN-P), Earth System Science Data 7(2), 245–259. https://essd.copernicus. org/articles/7/245/2015/.
- Biskaborn, B.K., Smith, S.L., Noetzli, J., Matthes, H., Vieira, G., Streletskiy, D.A. et al. (2019). Permafrost is warming at a global scale. Nature Communications 10, 264. https://www.nature.com/articles/s41467-018-08240-4.
- Dmitrenko, I.A., Kirillov, S.A., Tremblay, L.B., Kassens, H.,

Anisimov, O.A., Lavrov, S.A. et al. (2011). Recent changes in shelf hydrography in the Siberian Arctic: Potential for subsea permafrost instability. Journal of Geophysical Research: Oceans 116(C10), C10027. https://agupubs.onlinelibrary. wiley.com/doi/full/10.1029/2011JC007218.

- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J.W., Schuur, E.A.G., Ping, C.-L. et al. (2014). Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps. Biogeosciences 11(23), 6573–6593. https://bg.copernicus.org/articles/11/ 6573/2014/.
- Intergovernmental Panel on Climate Change (2013). Summary for Policymakers. In Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M.M.B., Allen, S.K., Boschung, J. et al. (eds.). Cambridge, UK and New York, US: Cambridge University Press. https:// www.ipcc.ch/site/assets/uploads/2018/03/WG1AR5_ SummaryVolume_FINAL.pdf.
- Schuur, T. (2019). Permafrost and the global carbon cycle, 30 October. https://arctic.noaa.gov/Report-Card/Report-Card-2019/ArtMID/7916/ArticleID/844/Permafrost-and-the-Global-Carbon-Cycle. Accessed 9 May 2023.
- Schuur, E.A.G., McGuire, A.D., Schädel, C., Grosse, G., Harden, J.W., Hayes, D.J. et al. (2015). Climate change and the permafrost carbon feedback. Nature 520, 171–179. https:// www.nature.com/articles/nature14338.

Maps

- Hugelius, G., Bockheim, J.G., Camill, P., Elberling, B., Grosse, G., Harden, J.W. et al. (2013). A new data set for estimating organic carbon storage to 3 m depth in soils of the northern circumpolar permafrost region. Earth System Science Data 5(2), 393–402. https://doi.org/10.5194/essd-5-393-2013.
- Miesner, F., Overduin, P.P., Grosse, G., Strauss, J., Langer, M., Westermann, S. et al. (2023). Subsea permafrost organic carbon stocks are large and of dominantly low reactivity. Scientific Reports 13, 9425. https://doi.org/10.1038/s41598-023-36471-z.
- Strauss, J., Laboor, S., Fedorov, A.N., Fortier, D., Froese, D.G., Fuchs, M. et al. (2016). Database of Ice-Rich Yedoma Permafrost (IRYP). https://doi.org/10.1594/PANGAEA.861733. Accessed 20 June 2023.

When ice grows up

Text

Bennett, M.R. (2009). Pingos. In Encyclopedia of Paleoclimatology and Ancient Environments. Gornitz, V. (ed.). Dordrecht, Netherlands: Springer. 783–784. https://link.springer.com/ referenceworkentry/10.1007/978-1-4020-4411-3 187.

- Mackay, J.R. (1979). Pingos of the Tuktoyaktuk Peninsula Area, Northwest Territories. Géographie Physique et Quaternaire 33(1), 3–61. https://www.erudit.org/en/journals/gpq/1979v33-n1-gpq1495571/1000322ar/.
- Mackay, J.R. (1986). Growth of Ibyuk Pingo, Western Arctic Coast, Canada, and some implications for environmental reconstructions. Quaternary Research 26(1), 68–80. https://www.cambridge.org/core/ journals/quaternary-research/article/abs/growth-ofibyuk-pingo-western-arctic-coast-canada-and-someimplications-for-environmental-reconstructions/ BD8B61AFAB4048A7C8DC0392C354132B.
- Mackay, J.R. (1998). Pingo growth and collapse, Tuktoyaktuk Peninsula Area, Western Arctic Coast, Canada: A long-term field study. Géographie Physique et Quaternaire 52(3), 271–323. https://www.erudit.org/en/journals/gpq/1998-v52n3-gpq153/004847ar/.

Maps

Vieira, G., Pina, P., Whalen, D., Freitas, P., Pinheiro, D., Fraser, P. et al. (2019). Pingo Canadian Landmark. Ultra High Resolution Aerial Orthomosaic [image]. https://doi.org/10.5281/zenodo.4298906.

Drilling down

Text

- Biskaborn, B.K., Lanckman, J.-P., Lantuit, H., Elger, K., Streletskiy, D.A., Cable, W.L. et al. (2015). The new database of the Global Terrestrial Network for Permafrost (GTN-P). Earth System Science Data 7(2), 245–259. https://essd.copernicus. org/articles/7/245/2015/.
- International Permafrost Association (2010). IPA-IPY Thermal State of Permafrost (TSP) Snapshot Borehole Inventory, Version 1 (G02190). National Snow and Ice Data Center. https://nsidc. org/data/g02190/versions/1. Accessed 9 May 2023.
- Konstantinov, P., Zhelezniak, M., Basharin, N., Misailov, I. and Andreeva, A. (2020). Establishment of permafrost thermal monitoring sites in East Siberia. Land 9(12), 476. https:// www.mdpi.com/2073-445X/9/12/476.
- Pika, P., Tanski, G., Ulrich, M., Roy, L.-P., Calmels, F., Lantuit, H. et al. (2022). Landscape-related ground ice variability on the Yukon coastal plain inferred from computed tomography and remote sensing. EGU General Assembly 2022. Vienna, Austria, 23–27 May 2022. Vienna: European Geosciences Union. https://meetingorganizer.copernicus.org/EGU22/ EGU22-10196.html.

Maps

Pika, P., Tanski, G., Ulrich, M., Roy, L.-P., Calmels, F., Lantuit, H. et al. (in prep.). A multi scale approach assessing excess ground ice volume in permafrost in NW Canada.

Un/settled

Text

- Bartsch, A., Pointner, G., Nitze, I., Efimova, A., Jakober, D., Ley, S. et al. (2021). Expanding infrastructure and growing anthropogenic impacts along Arctic coasts. Environmental Research Letters 16(11), 115013. https://iopscience.iop.org/ article/10.1088/1748-9326/ac3176.
- Istomin, K., and Habeck, J.O. (2019). Permafrost zone soils and the traditional land use of the aboriginal populations of north-eastern European Russia and western Siberia: Research problem statement. Vestnik arkheologii, antropologii i etnografii 1(44), 108–119. https://iasc.info/about/publications-documents/publications-list/642-permafrost-zone-soils-and-the-traditional-land-use-of-the-aboriginal-populations-of-north-eastern-european-russia-and-western-siberia-research-problem-statement.
- Peterson, R.A. and Wendler, K. (2011). Preservation of traditional ice cellars in permafrost: A modeling study of zero-energy solutions for enhancing thermal resilience. Fairbanks, US: University of Alaska. https://sites.google.com/a/alaska.edu/ thermally-resilient-permafrost-cellars/.
- Ramage, J., Jungsberg, L., Wang, S., Westermann, S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https://link. springer.com/article/10.1007/s11111-020-00370-6.

Maps

Ramage, J., Jungsberg, L., Wang, S., Westermann. S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https:// doi.org/10.1007/S1111-020-00370-6.

Frozen states I

- Huntington, H.P., Anisimova, O., Christensen, T., Fenge, T., Hoel,
 A.H., Jung, T. et al. (2013). Provisioning and cultural services.
 In Arctic Biodiversity Assessment: Status and Trends in Arctic Biodiversity. Meltofte, H. (ed.). Odder, Denmark: Narayana
 Press. Chapter 18. 592–627. https://www.caff.is/assessment-series/233-arctic-biodiversity-assessment-2013.
- Istomin, K.V. and Habeck, J.O. (2016). Permafrost and indigenous land use in the northern Urals: Komi and Nenets reindeer husbandry. Polar Science, 10(3), 278–287. https://www. sciencedirect.com/science/article/pii/S1873965216300615.
- Ramage, J., Jungsberg, L., Wang, S., Westermann, S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https://link. springer.com/article/10.1007/s1111-020-00370-6.
- Streletskiy, D.A., Suter, L.J., Shiklomanov, N.I., Porfiriev, B.N. and Eliseev, D.O. (2019). Assessment of climate change impacts on

buildings, structures and infrastructure in the Russian regions on permafrost. Environmental Research Letters 14(2), 025003. https://iopscience.iop.org/article/10.1088/1748-9326/aaf5e6.

Maps

- Oak Ridge National Laboratory and East View Cartographic, Inc. (2019). LandScan 2019 Global Population Database. https://searchworks.stanford.edu/view/13736097. Accessed 20 June 2023.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi. org/10.1016/j.earscirev.2019.04.023.
- Ramage, J., Jungsberg, L., Wang, S., Westermann. S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https:// doi.org/10.1007/s1111-020-00370-6.

Frozen states II

Text

- Fritz, M., Vonk, J.E. and Lantuit, H. (2017). Collapsing Arctic coastlines. Nature Climate Change 7, 6–7. https://www.nature.com/articles/nclimate3188.
- GRID-Arendal (2020). Coastal and offshore permafrost in a changing Arctic, 13 October. https://storymaps.arcgis.com/stories/ c163de04de7849cdb917fee88015dd73. Accessed 9 May 2023.
- Irrgang, A.M., Bendixen, M., Farquharson, L.M., Baranskaya, A.V., Erikson, L.H., Gibbs, A.E. et al. (2022). Drivers, dynamics and impacts of changing Arctic coasts. Nature Reviews Earth & Environment 3, 39–54. https://www.nature.com/articles/ 543017-021-00232-1.
- Kanigan, J.C.N. and Kokelj, S.V. (2010). Review of current research on drilling-mud sumps in permafrost terrain, Mackenzie Delta region, NWT, Canada. GeoCalgary 2010: The 63rd Canadian Geotechnical Conference and 6th Canadian Permafrost Conference. Calgary, Canada, 12–15 September 2010. University of Calgary. 1473–1479. https://pubs.aina.ucalgary.ca/cpc/CPC6-1473.pdf.
- Ramage, J., Jungsberg, L., Wang, S., Westermann, S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https://link. springer.com/article/10.1007/s11111-020-00370-6.

Maps

- Oak Ridge National Laboratory and East View Cartographic, Inc. (2019). LandScan 2019 Global Population Database. https://searchworks.stanford.edu/view/13736097. Accessed 20 June 2023.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern

Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi.org/10.1016/j.earscirev.2019.04.023. Ramage, J., Jungsberg, L., Wang, S., Westermann. S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https:// doi.org/10.1007/s11111-020-00370-6.

Frozen states III

Text

- Christiansen, H.H., Gilbert, G.L., Neumann, U., Demidov,
 N., Guglielmin, M., Isaksen, K. et al. (2021). Ground ice content, drilling methods and equipment and permafrost dynamics in Svalbard 2016-2019 (PermaSval). In SESS report 2020. Moreno-Ibáñez et al. (eds). Longyearbyen: Svalbard Integrated Arctic Earth Observing System. Chapter 12. 259-275. https://doi.org/10.5281/zenod0.4294095
- Humlum, O., Instanes, A. and Sollid, J.L. (2003). Permafrost in Svalbard: A review of research history, climatic background and engineering challenges. Polar Research 22(2), 191–215. https://polarresearch.net/index.php/polar/article/view/2108.
- Ingeman-Nielsen, T. and Foged, N. (2018). Kangerlussuaq Airport: Establishing a borehole database in support of future operational and maintenance decisions. In Proceedings of the conference AIC2018: Transportation Infrastructure Engineering in Cold Regions, 74-75. BYG DTU Rapport No. R-388.
- Magnin, F., Etzelmüller, B., Westermann, S., Isaksen, K., Hilger, P. and Hermanns, R.L. (2019). Permafrost distribution in steep rock slopes in Norway: Measurements, statistical modelling and geomorphological processes. Earth Surface Dynamics 7, 1019–1040. https://esurf.copernicus.org/articles/7/1019/2019/.
 Meltofte, H. and Thing, H. (eds.) (1996). Zackenberg Ecological
- Research Operations, 1st Annual Report, 1995. Copenhagen: Danish Polar Center. https://g-e-m.dk/fileadmin/Resources/DMU/ GEM/Zackenberg/Nye_Zac_files/ZERO_Annual_Report1995.pdf. Quaile, I. (2020). Zackenberg revisited – Greenland climate
- research station under threat from permafrost thaw, 14 July. https://www.rcinet.ca/eye-on-the-arctic/2020/07/14/blogzackenberg-revisited-greenland-climate-research-stationunder-threat-from-permafrost-thaw/. Accessed 9 May 2023.
- Ramage, J., Jungsberg, L., Wang, S., Westermann, S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https://link. springer.com/article/10.1007/s11111-020-00370-6.

Maps

Oak Ridge National Laboratory and East View Cartographic, Inc. (2019). LandScan 2019 Global Population Database. https://searchworks. stanford.edu/view/13736097. Accessed 20 June 2023. Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi. org/10.1016/j.earscirev.2019.04.023.

Ramage, J., Jungsberg, L., Wang, S., Westermann. S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https:// doi.org/10.1007/S11111-020-00370-6.

Awakening Giant

Warming up, warming down

Text

- Biskaborn, B.K., Smith, S.L., Noetzli, J., Matthes, H., Vieira, G., Streletskiy, D.A. et al. (2019). Permafrost is warming at a global scale. Nature Communications 10, 264. https://www.nature.com/articles/s41467-018-08240-4.
- Intergovernmental Panel on Climate Change (2021). Climate Change 2021: The Physical Science Basis: Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Chen, Y. et al. (eds.). Cambridge, UK and New York, US: Cambridge University Press. https://oceanrep.geomar.de/id/ eprint/58039/1/IPCC_AR6_WGI_FullReport.pdf.
- Pihl, E., Alfredsson, E., Bengtsson, M., Bowen, K., Cástan Broto,
 V., Chou, K. et al. (2021). Ten new insights in climate science
 2020 a horizon scan. Global Sustainability 4, e5, 1-18.
 https://doi.org/10.1017/sus.2021.2
- Romanovsky, V., Isaksen, K., Drozdov, D., Anisimov, O., Instanes, A., Leibman, M. et al. (2017). Changing permafrost and its impacts. In Snow, Water, Ice and Permafrost in the Arctic (SWIPA) 2017. Symon, C. (ed.). Oslo: Arctic Monitoring and Assessment Programme. Chapter 4. 64–102. https://www. amap.no/documents/download/2987/inline.
- Smith, S.L., O'Neill, H.B., Isaksen, K., Noetzli, J. and Romanovsky, V.E. (2022). The changing thermal state of permafrost. Nature Reviews Earth & Environment 3, 10–23. https://www.nature.com/articles/s43017-021-00240-1.

Maps

- Biskaborn, B.K., Smith, S.L., Noetzli, J., Matthes, H., Vieira, G., Streletskiy, D.A. et al. (2019). Permafrost is warming at a global scale. Nature Communications 10, 264. https://doi. org/10.1038/s41467-018-08240-4.
- Romanovsky, V. (2021). Past, recent, and future changes in permafrost in Alaska: Measurements and modeling results. Cryosphere Transformation and Geotechnical Safety in the Arctic. Salekhard, Russia and Online, 8–12 November 2021. Fairbanks, Alaska: Arctic Research Consortium of the United States.

The chill is gone

Text

- Brown, J., Hinkel, K.M. and Nelson, F.E. (2000). The circumpolar active layer monitoring (calm) program:
 Research designs and initial results. Polar Geography 24(3), 166–258. https://www.tandfonline.com/doi/abs/10.1080/10889370009377698.
- Romanovsky, V., Isaksen, K., Drozdov, D., Anisimov, O., Instanes, A., Leibman, M. et al. (2017). Changing permafrost and its impacts. In Snow, Water, Ice and Permafrost in the Arctic (SWIPA) 2017. Symon, C. (ed.). Oslo: Arctic Monitoring and Assessment Programme. Chapter 4. 64–102. https:// www.amap.no/documents/download/2987/inline.
- Romanovsky, V.E., Smith, S.L., Isaksen, K., Nyland, K.E., Kholodov, A.L., Shiklomanov, N.I., et al. (2020). Terrestrial Permafrost. In State of the Climate in 2019. Bulletin of the American Meteorological Society 101(8), S265–S269. https://journals. ametsoc.org/view/journals/bams/101/8/2020BAMSStateof theClimate.xml?tab_body=pdf.
- Smith, S.L., O'Neill, H.B., Isaksen, K., Noetzli, J. and Romanovsky, V.E. (2022). The changing thermal state of permafrost. Nature Reviews Earth & Environment 3, 10–23. https://www.nature.com/articles/s43017-021-00240-1.
- Strand, S.M., Christiansen, H.H., Johansson, M., Åkerman, J. and Humlum, O. (2020). Active layer thickening and controls on interannual variability in the Nordic Arctic compared to the circum-Arctic. Permafrost and Periglacial Processes 32(1), 47–58. https://onlinelibrary.wiley.com/doi/10.1002/ppp.2088.

Maps

Obu, J., Westermann, S., Barboux, C., Bartsch, A., Delaloye, R., Grosse, G. et al. (2021). ESA Permafrost Climate Change Initiative (Permafrost_cci): Permafrost Active Layer Thickness for the Northern Hemisphere, v3.0. NERC EDS Centre for Environment Data Analysis. https://catalogue.ceda.ac.uk/ uuid/67a3f8c8dc914ef99f7f08ebod997e23. Accessed 20 June 2023. DOI: 10.5285/67a3f8c8dc914ef99f7f08ebod997e23

Disappearing act

Text

- Chadburn, S.E., Burke, E.J., Cox, P.M., Friedlingstein, P., Hugelius, G. and Westermann, S. (2017). An observationbased constraint on permafrost loss as a function of global warming. Nature Climate Change 7, 340–344. https://www. nature.com/articles/nclimate3262.
- Intergovernmental Panel on Climate Change (2021). Climate Change 2021: The Physical Science Basis: Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-

Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Chen, Y. et al. (eds.). Cambridge, UK and New York, US: Cambridge University Press. https://oceanrep.geomar.de/ id/eprint/58039/1/IPCC_AR6_WGI_FullReport.pdf.

- Obu, J. (2021). How much of the Earth's surface in underlain by permafrost? Journal of Geophysical Research: Earth Surface 126(5), e2021JF006123. https://agupubs.onlinelibrary.wiley. com/doi/10.1029/2021JF006123.
- Vasiliev, A.A., Drozdov, D.S., Gravis, A.G., Malkova, G.V., Nyland, K.E. and Streletskiy, D.A. (2020). Permafrost degradation in the Western Russian Arctic. Environmental Research Letters 15(4), 045001. https://iopscience.iop.org/ article/10.1088/1748-9326/ab6f12.

Maps

Chadburn, S., Burke, E.J., Cox, P., Friedlingstein, P., Hugelius, G. and Westermann, S. (2017). Estimated Future Permafrost Maps Constrained by Observed Relationships, with Link to Model Result Files in NetCDF Format. PANGAEA. https://doi.pangaea. de/10.1594/PANGAEA.873192. Accessed 20 June 2023.

Microorganisms, macro effects

Text

- Jansson, J.K. and Hofmockel, K.S. (2019). Soil microbiomes and climate change. Nature Reviews Microbiology 18, 35–46. https://www.nature.com/articles/s41579-019-0265-7.
- Schuur, E.A.G., Abbott, B.W., Commane, R., Ernakovich, J., Euskirchen, E., Hugelius, G. et al. (2022). Permafrost and climate change: Carbon cycle feedbacks from the warming Arctic. Annual Review of Environment and Resources 47, 343–371. https://www.annualreviews.org/doi/10.1146/ annurev-environ-012220-011847.
- Schuur, E.A.G., McGuire, A.D., Schädel, C., Grosse, G., Harden, J.W., Hayes, D.J. et al. (2015). Climate change and the permafrost carbon feedback. Nature 520, 171–179. https:// www.nature.com/articles/nature14338.

Maps

- Scheel, M. (2021). It's getting hot in here: Ancient microbes in thawing permafrost, 22 October.
- https://blogs.egu.eu/divisions/cr/2021/10/22/its-gettinghot-in-here-ancient-microbes-in-thawing-permafrost/. Accessed 20 June 2023.
- Wagner, D. (2021). Methanosarcina soligelidi A multiresistant archaeon from Siberian permafrost. In Microbial Life in the Cryosphere and Its Feedback on Global Change: Life in Extreme Environments vol. 7. Liebner, S. and Ganzert, L. (eds.). Berlin and Boston, United States: De Gruyter. Chapter 5. 113–140. https://doi.org/10.1515/ 9783110497083-005.

Faster, deeper, stronger I

Text

- Miner, K.R., Turetsky, M.R., Malina, E., Bartsch, A., Tamminen, J., McGuire, A.D. et al. (2022). Permafrost carbon emissions in a changing Arctic. Nature Reviews Earth & Environment 3, 55–67. https://www.nature.com/articles/s43017-021-00230-3.
- Packalen, M.S., Finkelstein, S.A. and McLaughlin, J.W. (2014). Carbon storage and potential methane production in the Hudson Bay Lowlands since mid-Holocene peat initiation. Nature Communications 5, 4078. https://www.nature.com/ articles/ncomms5078.
- Packalen, M.S., Finkelstein, S.A. and McLaughlin, J.W. (2016). Climate and peat type in relation to spatial variation of the peatland carbon mass in the Hudson Bay Lowlands, Canada. Journal of Geophysical Research: Biogeosciences 121(4), 1104–1117. https://agupubs.onlinelibrary.wiley.com/doi/ full/10.1002/2015JG002938.
- Turetsky, M.R., Abbott, B.W., Jones, M.C., Anthony, K.T., Olefeldt, D., Schuur, E.A.G. et al. (2020). Carbon release through abrupt permafrost thaw. Nature Geoscience 13, 138–143. https://www.nature.com/articles/s41561-019-0526-0.

Maps

- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J.W., Schuur, E.A.G., Ping, C.-L. et al. (2014). Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps. Biogeosciences 11(23), 6573–6593. https://doi.org/10.5194/bg-11-6573-2014.
- Olefeldt, D., Goswami, S., Grosse, G., Hayes, D., Hugelius, G., Kuhry, P. et al. (2016). Circumpolar distribution and carbon storage of thermokarst landscapes. Nature Communications 7, 13043. https://doi.org/10.1038/ncomms13043.

Faster, deeper, stronger II

- Kremenetski, K.V., Velichko, A.A., Borisova, O.K., MacDonald, G.M., Smith, L.C., Frey, K.E. et al. (2003). Peatlands of the Western Siberian lowlands: Current knowledge on zonation, carbon content and Late Quaternary history. Quaternary Science Reviews 22(5–7), 703–723. https://www.sciencedirect. com/science/article/abs/pii/S0277379102001968.
- Schirmeister, L., Grosse, G., Wetterich, S., Overduin, P.P., Strauss, J., Schuur, E.A.G. et al. (2011). Fossil organic matter characteristics in permafrost deposits of the northeast Siberian Arctic. Journal of Geophysical Research: Biogeosciences 116(G2), GooMo2. https:// agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2011JG001647.
- Streletskiy, D.A., Suter, L.J., Shiklomanov, N.I., Porfiriev,B.N. and Eliseev, D.O. (2019). Assessment of climate change impacts on buildings, structures and infrastructure

in the Russian regions on permafrost. Environmental Research Letters 14, 025003. https://iopscience.iop.org/ article/10.1088/1748-9326/aaf5e6#erlaaf5e6s2.

Turetsky, M.R., Abbott, B.W., Jones, M.C., Anthony, K.W., Olefeldt, D., Schuur, E.A.G. et al. (2019). Permafrost collapse is accelerating carbon release. Nature 569, 32–34. https:// www.nature.com/articles/d41586-019-01313-4.

Vasiliev, A.A., Drozdov, D.S., Gravis, A.G., Malkova, G.V., Nyland K.E. and Streletskiy, D.A. (2020). Permafrost degradation in the Western Russian Arctic. Environmental Research Letters 15(4), 045001. https://iopscience.iop.org/ article/10.1088/1748-9326/ab6f12.

Maps

Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J.W., Schuur, E.A.G., Ping, C.-L. et al. (2014). Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps. Biogeosciences 11(23), 6573–6593. https://doi.org/10.5194/bg-11-6573-2014.

Olefeldt, D., Goswami, S., Grosse, G., Hayes, D., Hugelius, G., Kuhry, P. et al. (2016). Circumpolar distribution and carbon storage of thermokarst landscapes. Nature Communications 7, 13043. https://doi.org/10.1038/ncomms13043.

Crossing the threshold

Text

Canadell, J.G., Monteiro, P.M.S., Costa, M.H., Cotrim da Cunha, L., Cox, P.M., Eliseev, A.V. et al. (2021). Global carbon and other biogeochemical cycles and feedbacks. In Climate Change 2021: The Physical Science Basis: Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Berger, S. et al. (eds.). Cambridge, UK and New York, US: Cambridge University Press. Chapter 5. 673–816. https://oceanrep.geomar.de/id/eprint/58039/1/ IPCC_AR6_WGI_FullReport.pdf.

Intergovernmental Panel on Climate Change (2019). Technical summary. In The Ocean and Cryosphere in a Changing Climate: Special Report of the Intergovernmental Panel on Climate Change. Pörtner, H.-O., Roberts, D.C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E. et al. (eds.). Cambridge, UK and New York, US: Cambridge University Press. 39–70. https://www.cambridge.org/core/services/aop-cambridge-core/content/view/Ao5E6C9F8638FA7CE1748DE2EB7B491B/9781009157971AR. pdf/The_Ocean_and_Cryosphere_in_a_Changing_Climate.pdf.

Kleinen, T. and Brovkin, V. (2018). Pathway-dependent fate of permafrost region carbon. Environmental Research Letters 13(9), 094001. https://iopscience.iop.org/article/10.1088/ 1748-9326/aad824. Turetsky, M.R., Abbott, B.W., Jones, M.C., Anthony, K.W., Olefeldt, D., Schuur, E.A.G. et al. (2020). Carbon release through abrupt permafrost thaw. Nature Geoscience 13, 138– 143. https://www.nature.com/articles/s41561-019-0526-0. United Nations. Framework Convention on Climate Change

(2015). Adoption of the Paris Agreement. 12 December. FCCC/ CP/2015/L.9/Rev.1. https://unfccc.int/resource/docs/2015/ cop21/eng/lo9r01.pdf.

Wilkenskjeld, S., Miesner, F., Overduin, P.P., Puglini, M. and Brovkin, V. (2022). Strong increase in thawing of subsea permafrost in the 22nd century caused by anthropogenic climate change. The Cryosphere 16(3), 1057–1069. https:// tc.copernicus.org/articles/16/1057/2022/.

Maps

Kleinen, T. and Brovkin, V. (2018). Pathway-dependent fate of permafrost region carbon. Environmental Research Letters 13(9), 094001.https://doi.org/10.1088/1748-9326/aad824.
Ridolfi, E., Geels, A., Lefebvre, C. and Arndt, S. (unpublished).

Moving Grounds

Frost and flora

Text

- Circumpolar Arctic Vegetation Mapping Team (2023). Raster Circumpolar Arctic Vegetation Map, Version 2. Scale 1:7,000,000. Mendeley Data. https://www.geobotany.uaf. edu/cavm/.
- Druckenmiller, M. L., Thoman, R. L. and Moon, T. A. (2022). Arctic Report Card 2022: Executive Summary. National Oceanic and Atmospheric Administration. https://doi. org/10.25923/yix6-r184
- Heijmans, M.M.P.D., Magnússon, R.Í., Lara, M.J., Frost, G.V., Myers-Smith, I.H., van Huissteden, J. et al. (2022). Tundra vegetation change and impacts on permafrost. Nature Reviews Earth & Environment 3, 68–84. https://www.nature. com/articles/s43017-021-00233-0.
- Raynolds, M.K., Walker, D.A., Balser, A., Bay, C., Campbell, M., Cherosov, M.M. et al. (2019). A raster version of the Circumpolar Arctic Vegetation Map (CAVM). Remote Sensing of Environment 232, 111297. https://www.sciencedirect.com/ science/article/pii/S0034425719303165.
- Scholten, R.C., Jandt, R., Miller, E.A., Rogers, B.M. and Veraverbeke, S. (2021). Overwintering fires in boreal forests. Nature 593, 399– 404. https://www.nature.com/articles/S41586-021-03437-y.
- Smith, S.L., O'Neill, H.B., Isaksen, K., Noetzli, J. and Romanovsky, V.E. (2022). The changing thermal state of permafrost. Nature Reviews Earth & Environment 3, 10–23. https://www.nature.com/articles/s43017-021-00240-1.
 United Nations Environment Programme (2022). Spreading like

wildfire: The rising threat of extraordinary landscape fires: A rapid response assessment. Nairobi: United Nations Environment Programme. https://www.unep.org/resources/report/spreadingwildfire-rising-threat-extraordinary-landscape-fires.

Maps

- Frost, G.V., Macander, M.J., Bhatt, U.S., Berner, L.T., Bjerke, J.W., Epstein, H.E. et al. (2021). Tundra Greenness. In Arctic Report Card 2021, NOAA Technical Report OAR ARC; 21-08. United States: NOAA Institutional Repository. 1–8. https:// doi.org/10.25923/8n78-wp73.
- Li, J., Holmgren, M. and Xu, C. (2021). Greening vs browning? Surface water cover mediates how tundra and boreal ecosystems respond to climate warming. Environmental Research Letters 16(10), 104004. https://doi.org/10.1088/1748-9326/ac2376.
- Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G V N., Underwood, E.C. et al. (2001). Terrestrial ecoregions of the world: A new map of life on earth: A new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity. BioScience 51(11), 933–938. https://doi. org/10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2.

Fire on ice

- Chapin III, F.S., Mcguire, A.D., Randerson, J., Pielke, R., Baldocchi, D., Hobbie, S.E., et al. (2000). Arctic and boreal ecosystems of western North America as components of the climate system. Global Change Biology 6(S1), 211–223. https://onlinelibrary. wiley.com/doi/10.1046/j.1365-2486.2000.06022.x.
- Conard, S.G. and Ponomarev, E. (2020). Fire in the North: The 2020 Siberian fire season, October–December. https://www.iawfonline. org/article/fire-in-the-north-the-2020-siberian-fire-season/. Accessed 10 May 2023.
- Food and Agriculture Organization of the United Nations (2020). Peatland mapping and monitoring: Recommendations and technical overview. Rome: Food and Agriculture Organization of the United Nations. https://www.fao.org/3/CA8200EN/ CA8200EN.pdf.
- Helbig, M., Waddington, J.M., Alekseychik, P., Amiro, B.D.,
 Aurela, M., Barr, A.G. et al. (2020). Increasing contribution of peatlands to boreal evapotranspiration in a warming climate.
 Nature Climate Change 10, 555–560. https://www.nature.
 com/articles/s41558-020-0763-7.pdf.
- Hugelius, G., Loisel, J., Chadburn, S., Jackson, R.B., Jones, M., MacDonald, G. et al. (2020). Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw. PNAS 117(34), 20438–20446. https://www.pnas.org/doi/ full/10.1073/pnas.1916387117.
- Intergovernmental Panel on Climate Change (2021). Climate Change 2021: The Physical Science Basis: Contribution of Working Group

I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Chen, Y. et al. (eds.). Cambridge, UK and New York, US: Cambridge University Press. https://oceanrep. geomar.de/id/eprint/58039/1/IPCC_AR6_WGI_FullReport.pdf.

- United Nations Environment Programme (2019). Permafrost peatlands: Losing ground in a warming world. In Frontiers 2018/19: Emerging Issues of Environmental Concern. Nairobi: United Nations Environment Programme. Chapter 3. 38–48. https://wedocs.unep.org/bitstream/handle/20.500.11822/ 27542/Frontiers1819_ch3.pdf.
- United Nations Environment Programme (2022). Global peatlands assessment: The state of the world's peatlands: Evidence for action toward the conservation, restoration, and sustainable management of peatlands: Main report. Nairobi: United Nations Environment Programme. https://globalpeatlands.org/ sites/default/files/2022-12/peatland_assessment.pdf.
- Witze, A. (2020). The Arctic is burning like never before and that's bad news for climate change. Nature 585, 336-337. https://www.nature.com/articles/d41586-020-02568-y. Accessed 10 May 2023.
- Zhang, Z., Wang, C., Lv, D., Hay, W.W., Wang, T. and Cao, S. (2020). Precession-scale climate forcing of peatland wildfires during the early middle Jurassic greenhouse period. Global and Planetary Change 184, 103051. https://doi.org/10.1016/j. gloplacha.2019.103051

Maps

- Greifswald Mire Centre (2023). The Global Peatland Database. https://www.greifswaldmoor.de/global-peatland-databaseen.html. Accessed 20 June 2023.
- NASA Earth Observations (2020). Active Fires (1 Month Terra/ MODIS). NASA Earth Observations. https://neo.gsfc.nasa. gov/view.php?datasetId=MOD14A1_M_FIRE&year=2020. Accessed 20 June 2023.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi. org/10.1016/j.earscirev.2019.04.023.

Overduin, P.P., Schneider von Deimling, T., Miesner, F., Grigoriev, M.N., Ruppel, C., Vasiliev, A. et al. (2019). Submarine permafrost map in the Arctic modeled using 1-D transient heat flux (SuPerMAP). Journal of Geophysical Research: Oceans 124(6), 3490–3507. https://doi.org/10.1029/2018JC014675.

State of matter

Text

Douglas, T.A., Turetsky, M.R. and Koven, C.D. (2020). Increased rainfall stimulates permafrost thaw across a variety of Interior

Alaskan boreal ecosystems. Npj Climate and Atmospheric Science 3, 28. https://www.nature.com/articles/s41612-020-0130-4.

- Fritz, M., Wolter, J., Rudaya, N., Palagushkina, O., Nazarova, L., Obu, J. et al. (2016). Holocene ice-wedge polygon development in northern Yukon permafrost peatlands (Canada). Quaternary Science Reviews 147, 279–297. https://www.sciencedirect. com/science/article/abs/pii/S0277379116300403.
- Liljedahl, A., Boike, J., Daanen, R.P., Fedorov, A.N., Frost, G.V., Grosse, G. et al. (2016). Pan-Arctic ice-wedge degradation in warming permafrost and its influence on tundra hydrology. Nature Geoscience 9, 312–318. https://www.nature.com/ articles/nge02674.
- O'Neill, H.B. and Burn, C.R. (2017). Impacts of variations in snow cover on permafrost stability, including simulated snow management, Dempster Highway, Peel Plateau, Northwest Territories. Arctic Science 3(2), 150–178. https:// cdnsciencepub.com/doi/10.1139/as-2016-0036.
- Smith, S.L., O'Neill, H.B., Isaksen, K., Noetzli, J. and Romanovsky, V.E. (2022). The changing thermal state of permafrost. Nature Reviews Earth & Environment 3, 10–23. https://www.nature.com/articles/s43017-021-00240-1.
- Speetjens, N.J., Hugelius, G., Gumbricht, T., Lantuit, H., Berghuijs, W.R., Pika, P.A. et al. (2023) The pan-Arctic catchment database (ARCADE). Earth System Science Data 15(2), 541–554. https:// essd.copernicus.org/articles/15/541/2023/.
- Vonk, J.E., Tank, S.E. and Walvoord, M.A. (2019). Integrating hydrology and biogeochemistry across frozen landscapes. Nature Communications 10, 5377. https://www.nature.com/ articles/s41467-019-13361-5.
- Vonk, J.E., Speetjens, N.J. and Poste, A.E. (2023). Small watersheds may play a disproportionate role in arctic landocean fluxes. Nature Communications 14, 3442. https://www. nature.com/articles/s41467-023-39209-7.
- Zhang, S.-M., Mu, C.-C., Li, Z.-L., Dong, W.-W., Wang, X.-Y., Streletskaya, I. et al. (2021). Export of nutrients and suspended solids from major Arctic rivers and their response to permafrost degradation. Advances in Climate Change Research 12(4), 466–474. https://www.sciencedirect.com/ science/article/pii/S1674927821000824.
- Zheng, L., Overeem, I., Wang, K. and Clow, G. D. (2019). Changing Arctic river dynamics cause localized permafrost thaw. Journal of Geophysical Research: Earth Surface 124(9), 2324–2344. https://agupubs.onlinelibrary.wiley.com/doi/ full/10.1029/2019JF005060.

Maps

McCrystall, M.R., Stroeve, J., Serreze, M., Forbes, B.C. and Screen, J.A. (2021). New climate models reveal faster and larger increases in Arctic precipitation than previously projected. Nature Communications 12, 6765. https://doi. org/10.1038/541467-021-27031-y.

- Sathyendranath, S., Brewin, R.J.W., Brockmann, C., Brotas, V., Calton, B., Chuprin, A. et al. (2019). An ocean-colour time series for use in climate studies: The experience of the Ocean-Colour Climate Change Initiative (OC-CCI). Sensors 19(19), 4285. https://doi.org/10.3390/s19194285.
- Speetjens, N.J., Hugelius, G., Gumbricht, T., Lantuit, H., Berhuijs, W.R., Pika, P.A. et al. (2023). The pan-Arctic catchment database (ARCADE). Articles 15(2), 541–554. https://doi.org/10.5194/essd-15-541-2023.

The rivers run through it

- Beel, C.R., Heslop, J.K., Orwin, J.F., Pope, M.A., Schevers, A.J., Hung, J.K.Y. et al. (2021). Emerging dominance of summer rainfall driving High Arctic terrestrial-aquatic connectivity. Nature Communications 12, 1448. https://doi.org/10.1038/ 541467-021-21759-3.
- Forbes, D.L. (2019). Arctic deltas and estuaries: A Canadian perspective. In Coasts and Estuaries: The Future. Wolanski, E., Day, J., Elliott, M. and Ramachandran, R. (eds.). Amsterdam, Netherlands, Oxford, UK and Cambridge, US: Elsevier. Chapter 8. 123–147. https://doi.org/10.1016/B978-0-12-814003-1.00008-3.
- Fuchs, M., Nitze, I., Strauss, J., Günther, F., Wetterich, S., Kizyakov, A. et al. (2020). Rapid fluvio-thermal erosion of a Yedoma permafrost cliff in the Lena River Delta. Frontiers in Earth Science 8, 336. https://doi.org/10.3389/feart.2020.00336.
- Holmes, R.M., McClelland, J.W., Peterson, B.J., Tank, S.E., Bulygina, E., Eglinton, T.I. et al. (2012). Seasonal and Annual fluxes of nutrients and organic matter from large rivers to the Arctic Ocean and surrounding seas. Estuaries and Coasts 35, 369–382. https://doi.org/10.1007/s12237-011-9386-6.
- Juhls, B., Antonova, S., Angelopoulos, M., Bobrov, N., Grigoriev, M., Langer, M. et al. (2021). Serpentine (floating) ice channels and their interaction with riverbed permafrost in the Lena River Delta, Russia. Frontiers in Earth Science 9, 689941. https://doi.org/10.3389/feart.2021.689941.
- Kokelj, S.V., Kokoszka, J., van der Sluijs, J., Rudy, A.C.A., Tunnicliffe, J., Shakil, S. et al. (2021). Thaw-driven mass wasting couples slopes with downstream systems, and effects propagate through Arctic drainage networks. The Cryosphere 15(7), 3059–3081. https://doi.org/10.5194/tc-15-3059-2021.
- Overeem, I., Nienhuis, J.H. and Piliouras, A. (2022). Icedominated Arctic deltas. Nature Reviews Earth & Environment 3, 225–240. https://doi.org/10.1038/543017-022-00268-x.
- Semiletov, I., Pipko, I., Gustafsson, Ö., Anderson, L.G., Sergienko, V., Pugach, S. et al. (2016). Acidification of East Siberian Arctic Shelf waters through addition of freshwater and terrestrial carbon. Nature Geoscience 9, 361–365. https://doi.org/10.1038/nge02695
- Speetjens, N.J., Hugelius, G., Gumbricht, T., Lantuit, H., Berghuijs,

W.R., Pika, P.A. et al. (2023) The pan-Arctic catchment database (ARCADE). Earth System Science Data 15(2), 541–554. https://doi.org/10.5194/essd-15-541-2023.

Wagner, A., Lohmann, G. and Prange, M. (2011). Arctic river discharge trends since 7 ka BP. Global and Planetary Change 79(1–2), 48–60. https://doi.org/10.1016/j.gloplacha.2011.07.006.

Maps

European Union Agency for the Space Programme (2020). Copernicus Data Space Ecosystem: Satellite Imagery for 9 September 2020. https://dataspace.copernicus.eu/. Accessed 20 June 2023.

Along the edge of the world

Text

- Boisson, A. and Allard, M. (2020). Boulder barricades in an emerging hypertidal environment: The case of Ungava Bay, northern Quebec, Canada. Geografiska Annaler: Series A, Physical Geography 102(1), 33–56. https://www.tandfonline. com/doi/full/10.1080/04353676.2020.1711665.
- Boisson, A., Allard, M. and Sarrazin, D. (2020). Permafrost aggradation along the emerging eastern coast of Hudson Bay, Nunavik (northern Quebec, Canada). Permafrost and Perglacial Processes 31(1), 128–140. https://onlinelibrary. wiley.com/doi/10.1002/ppp.2033.
- Forbes, D.L. (2019). Arctic deltas and estuaries: A Canadian perspective. In: Coasts and Estuaries: The Future. Wolanski, E., Day, J., Elliott, M. and Ramachandran, R. (eds.). Amsterdam, Netherlands, Oxford, UK and Cambridge, US: Elsevier. Chapter 8. 123–147. https://www.sciencedirect.com/science/article/abs/pii/B9780128140031000083.
- Forbes, D.L. and Taylor, R.B. (1994). Ice in the shore zone and the geomorphology of cold coasts. Progress in Physical Geography: Earth and Environment 18(1), 59–89. https:// journals.sagepub.com/doi/10.1177/030913339401800104.
- Fritz, M., Vonk, J.E. and Lantuit, H. (2017). Collapsing Arctic coastlines. Nature Climate Change 7, 6–7. https://www.nature.com/articles/nclimate3188.
- Hansell, R.I.C., Scott, P.A., Staniforth, R. and Svoboda, J. (1983). Permafrost development in the intertidal zone at Churchill, Manitoba: A possible mechanism for accelerated beach uplift. Arctic 36(2), 198–203. https://journalhosting.ucalgary. ca/index.php/arctic/article/view/65315.
- Hansom, J.D., Forbes, D.L. and Etienne, S. (2014). The rock coasts of polar and sub-polar regions. In Rock Coast Geomorphology: A Global Synthesis. Kennedy, D. M., Stephenson, W.J. and Naylor, L.A. (eds.). Bath, UK: Geological Society of London. 263–281. https://pubs.geoscienceworld. org/gsl/books/book/1781/chapter-abstract/107679778/Therock-coasts-of-polar-and-sub-polar-regions.

- Hatcher, S.V., Forbes, D.L. and Manson, G.K. (2021). Boulderstrewn flats in a high-latitude macrotidal embayment, Baffin Island: geomorphology, formation, and future stability. Canadian Journal of Earth Sciences 59(11), 866–883. https:// doi.org/10.1139/cjes-2021-0019
- Irrgang, A.M., Bendixen, M., Farquharson, L.M., Baranskaya, A.V., Erikson, L.H., Gibbs, A.E. et al. (2022). Drivers, dynamics and impacts of changing Arctic coasts. Nature Reviews Earth & Environment 3, 39–54. https://www.nature.com/articles/ \$43017-021-00232-1.
- Jones, B.M, Arp, C.D., Jorgenson, M.T., HInkel, K.M., Schmutz, J.A. and Flint, P.L. (2009). Increase in the rate and uniformity of coastline erosion in Arctic Alaska. Geophysical Research Letters 36, L03503. https://doi. org/10.1029/2008GL036205
- Lantuit, H., Overduin, P.P., Couture, N., Wetterich, S. Aré, F., Atkinson, D. et al. (2012). The Arctic Coastal Dynamics Database: A new classification scheme and statistics on Arctic permafrost coastlines. Estuaries and Coasts 35, 383–400. https://link.springer.com/article/10.1007/s12237-010-9362-6.
- National Snow and Ice Data Center (2021). Arctic sea ice at highest minimum since 2014, 22 September. https://nsidc. org/arcticseaicenews/2021/09/arctic-sea-ice-at-highestminimum-since-2014/. Accessed 10 May 2023.

Maps

- Lantuit, H., Overduin, P.P., Couture, N., Wetterich, S., Aré, F., Atkinson, D. et al. (2012). The Arctic Coastal Dynamics database: A new classification scheme and statistics on Arctic permafrost coastlines. Estuaries and Coasts 35, 383–400. https://doi.org/10.1007/S12237-010-9362-6. Obu, J., Westermann, S., Barboux, C., Bartsch, A., Delaloye, R., Grosse, G. et al. (2021). ESA Permafrost Climate Change Initiative (Permafrost_cci): Permafrost Active Layer Thickness for the Northern Hemisphere, v3.0. NERC EDS Centre for Environment Data Analysis. https://catalogue.ceda.ac.uk/ uuid/67a3f8c8dc914ef99f7f08ebod997e23. Accessed 20 June
- 2023. DOI: 10.5285/67a3f8c8dc914ef99f7f08eb0d997e23 Ramage, J., Jungsberg, L., Wang, S., Westermann. S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https:// doi.org/10.1007/S11111-020-00370-6.

Wear and tear

Text

Aré, F.E. (1988). Thermal abrasion of sea coasts (part I).
Polar Geography and Geology 12(1), 1–86. https://www. tandfonline.com/doi/abs/10.1080/10889378809377343.
Atkinson, D.A. (2005). Observed storminess patterns and trends in the circum-Arctic coastal regime. Geo-Marine Letters, 25, 98-109. https://doi.org/10.1007/s00367-004-0191-0

- Grosse, G., Nitze, I. Sachs, T., Gessner, M. (2019). Master tracks in different resolutions during POLAR 6 campaign P6_219_ThawTrend_Air_2019. Alfred Wegener Institute Research Unit Potsdam, PANGAEA, https://doi.org/10.1594/PANGAEA.905792. Accessed 04 April 2023.
- Fritz, M., Vonk, J.E. and Lantuit, H. (2017). Collapsing Arctic coastlines. Nature Climate Change 7, 6–7. https://www.nature.com/articles/nclimate3188.
- Günther, F., Overduin, P.P., Sandakov, A.V., Grosse, G. and Grigoriev, M.N. (2013). Short- and long-term thermo-erosion of ice-rich permafrost coasts in the Laptev Sea region. Biogeosciences 10(6), 4297–4318. https://bg.copernicus.org/ articles/10/4297/2013/.
- Irrgang, A.M., Bendixen, M., Farquharson, L.M., Baranskaya, A.V., Erikson, L.H., Gibbs, A.E. et al. (2022). Drivers, dynamics and impacts of changing Arctic coasts. Nature Reviews Earth & Environment 3, 39–54. https://www.nature.com/articles/ \$43017-021-00232-1.
- Jones, B.M., Farquharson, L.M., Baughman, C.A., Buzard, R.M., Arp, C.D., Grosse, G. et al. (2018). A decade of remotely sensed observations highlight complex processes linked to coastal permafrost bluff erosion in the Arctic. Environmental Research Letters 13(11), 115001. https://iopscience.iop.org/ article/10.1088/1748-9326/aae471/meta.
- Tanski, G., Bröder, L., Wagner, D., Knoblauch, C., Lantuit, H., Beer, C. et al. (2021). Permafrost carbon and CO2 pathways differ at contrasting coastal erosion sites in the Canadian Arctic. Frontiers in Earth Science 9, 630493. https://www. frontiersin.org/articles/10.3389/feart.2021.630493/full.
- Tanski, G., Wagner, D., Knoblauch, C., Fritz, M., Sachs, T. and Lantuit, H. (2019). Rapid CO2 release from eroding permafrost in seawater. Geophysical Research Letters 46(20), 11244–11252. https:// agupubs.onlinelibrary.wiley.com/doi/10.1029/2019GL084303.
- Terhaar, J., Lauerwald, R., Regnier, P., Gruber, N. and Bopp, L. (2021). Around one third of current Arctic Ocean primary production sustained by rivers and coastal erosion. Nature Communications 12, 169. https://www.nature.com/articles/s41467-020-20470-z.
- Wegner, C., Bennett, K. E., de Vernal, A., Forwick, M., Fritz, M., Heikkilä, M. et al. (2015). Variability in transport of terrigenous material on the shelves and the deep Arctic Ocean during the Holocene. Polar Research 34, 1. https:// polarresearch.net/index.php/polar/article/view/3254.

Maps

- Fritz, M., Vonk, J.E. and Lantuit, H. (2017). Collapsing Arctic coastlines. Nature Climate Change 7, 6–7. https://doi.org/ 10.1038/nclimate3188.
- Grosse, G., Nitze, I., Sachs, T. and Gessner, M. (2019). Master Tracks in Different Resolutions During POLAR 6 Campaign P6_219_ThawTrend_Air_2019. Alfred Wegener Institute –

Research Unit Potsdam, PANGAEA. https://doi.org/10.1594/ PANGAEA.905792. Accessed 20 June 2023.

Irrgang, A.M., Bendixen, M., Farquharson, L.M., Baranskaya, A.V., Erikson, L.H., Gibbs, A.E. et al. (2022). Drivers, dynamics and impacts of changing Arctic coasts. Nature Reviews Earth & Environment 3, 39–54. https://doi.org/10.1038/s43017-021-00232-1.

Eating away at the landscape

Text

- Bernhard, P., Zwieback, S. and Hajnsek, I. (2022). Accelerated mobilization of organic carbon from retrogressive thaw slumps on the northern Taymyr Peninsula. The Cryosphere 16(7), 2819–2835. https://tc.copernicus.org/ articles/16/2819/2022/.
- Costard, F., Dupeyrat, L., Séjourné, A., Bouchard, F., Fedorov, A. and Saint-Bézar, B. (2020). Retrogressive thaw slumps on icerich permafrost under degradation: Results from a large-scale laboratory simulation. Geophysical Research Letters 48(1), e2020GL091070. https://agupubs.onlinelibrary.wiley.com/ doi/10.1029/2020GL091070.
- Turetsky, M.R., Abbott, B.W., Jones, M.C., Anthony, K.W., Olefeldt, D., Schuur E.A.G. et al. (2020). Carbon release through abrupt permafrost thaw. Nature Geoscience 13, 138– 143. https://www.nature.com/articles/s41561-019-0526-0.

Maps

Vieira, G. and Pina, P. (Unpublished). Qikiqtaryuk Slump D ultra-high resolution aerial orthomosaic, July 2018. Centre for Geographical Studies of the Institute of Geography and Spatial Planning, University of Lisbon and National Resources Canada.

Losing ground

Text

Fritz, M., Vonk, J.E. and Lantuit, H. (2017). Collapsing Arctic coastlines. Nature Climate Change 7, 6–7. https://www.nature.com/articles/nclimate3188.

Lantuit, H., Overduin, P.P., Couture, N., Wetterich, S., Aré, F., Atkinson, D. et al. (2011). The Arctic Coastal Dynamics Database: A new classification scheme and statistics on Arctic permafrost coastlines. Estuaries and Coasts 35, 383–400. https://link.springer.com/article/10.1007/s12237-010-9362-6.

Nielsen, D.M., Pieper, P., Barkhordarian, A., Overduin, P., Ilyina, T., Brovkin, V. et al. (2022). Increase in Arctic coastal erosion and its sensitivity to warming in the twenty-first century. Nature Climate Change 12, 263–270. https://www.nature. com/articles/s41558-022-01281-0. Tanski, G., Bröder, L., Wagner, D., Knoblauch, C., Lantuit, H., Beer, C. et al. (2021). Permafrost carbon and CO2 pathways differ at contrasting coastal erosion sites in the Canadian Arctic. Frontiers in Earth Science 9, 207. https://www. frontiersin.org/articles/10.3389/feart.2021.630493/full. Terhaar, J., Lauerwald, R., Regnier, P., Gruber, N. and Bopp,

L. (2021). Around one third of current Arctic Ocean primary production sustained by rivers and coastal erosion. Nature Communications 12, 169. https://www.nature.com/articles/ \$41467-020-20470-Z.

Maps

- Lantuit, H., Overduin, P.P., Couture, N., Wetterich, S., Aré, F., Atkinson, D. et al. (2012). The Arctic Coastal Dynamics database: A new classification scheme and statistics on Arctic permafrost coastlines. Estuaries and Coasts 35, 383-400. https://doi.org/10.1007/S12237-010-9362-6.
- Nielsen, D.M., Pieper, P., Barkhordarian, A., Overduin, P., Ilyina, T., Brovkin, V. et al. (2022). Increase in Arctic coastal erosion and its sensitivity to warming in the twenty-first century. Nature Climate Change 12, 263–270. https://doi.org/10.1038/ \$41558-022-01281-0.
- Obu, J., Westermann, S., Barboux, C., Bartsch, A., Delaloye, R., Grosse, G. et al. (2021). ESA Permafrost Climate Change Initiative (Permafrost_cci): Permafrost Active Layer Thickness for the Northern Hemisphere, v3.0. NERC EDS Centre for Environment Data Analysis. https://catalogue.ceda.ac.uk/ uuid/67a3f8c8dc914ef99f7f08eb0d997e23. Accessed 20 June 2023. DOI: 10.5285/67a3f8c8dc914ef99f7f08eb0d997e23
- Ramage, J., Jungsberg, L., Wang, S., Westermann. S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https:// doi.org/10.1007/S11111-020-00370-6.

Beneath the waves

Text

- Angelopoulos, M., Overduin, P.P., Miesner, F., Grigoriev, M.N. and Vasiliev, A.A. (2020). Recent advances in the study of Arctic submarine permafrost. Permafrost and Periglacial Processes 31(3), 442–453. https://onlinelibrary.wiley.com/ doi/full/10.1002/ppp.2061.
- GRID-Arendal (2020). Coastal and offshore permafrost in a changing Arctic, 13 October. https://storymaps.arcgis.com/ stories/c163de04de7849cdb917fee88015dd73. Accessed 9 May 2023.
- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J.W., Schuur, E.A.G., Ping, C.-L. et al. (2014). Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps. Biogeosciences 11(23), 6573–6593. https://bg.copernicus.org/articles/11/6573/2014/.

- Miesner, F., Overduin, P.P., Grosse, G., Strauss, J., Langer, M., Westermann, S. et al. (2023). Subsea permafrost organic carbon stocks are large and of dominantly low reactivity. Scientific Reports, 13, 9425. https://doi.org/10.1038/s41598-023-36471-z
- Overduin, P.P., von Deimling, T.S., Miesner, F., Grigoriev, M.N., Ruppel, C., Vasiliev, A. et al. (2019). Submarine permafrost map in the Arctic modeled using 1-D transient heat flux (SuPerMAP). Journal of Geophysical Research: Oceans 124(6), 3490–3507. https://agupubs.onlinelibrary.wiley.com/doi/ full/10.1029/2018JC014675.
- Paull, C.K., Dallimore, S.R., Caress, D.W., Gwiazda, R., Melling,
 H., Riedel, M. et al. (2015). Active mud volcanoes on the continental slope of the Canadian Beaufort Sea. Geochemistry, Geophysics, Geosystems 16(9), 3160–3181. https://agupubs. onlinelibrary.wiley.com/doi/10.1002/2015GC005928.
- Ruppel, C. (2015). Permafrost-associated gas hydrate: is it really approximately 1% of the global system? Journal of Chemical & Engineering Data 60(2), 429–436. https://pubs.acs.org/ doi/10.1021/je500770m.
- Schuur, E.A.G., Abbott, B.W., Commane, R., Ernakovich, J., Euskirchen, E., Hugelius, G. et al. (2022). Permafrost and climate change: Carbon cycle feedbacks from the warming Arctic. Annual Review of Environment and Resources 47, 343–371. https://www.annualreviews.org/doi/10.1146/ annurev-environ-012220-011847.
- Wild, B., Shakhova, N., Dudarev, O., Ruban, A., Kosmach, D., Tumskoy, V. et al. (2022). Organic matter composition and greenhouse gas production of thawing subsea permafrost in the Laptev Sea. Nature Communications 13, 5057. https:// www.nature.com/articles/s41467-022-32696-0.

Maps

- Majorowicz, J.A. and Hannigan, P.K. (2000). Stability zone of natural gas hydrates in a permafrost-bearing region of the Beaufort—Mackenzie Basin: Study of a feasible energy source1 (Geological Survey of Canada Contribution No.1999275). Natural Resources Research 9, 3–26. https:// doi.org/10.1023/A:1010105628952.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi.org/10.1016/j.earscirev.2019.04.023.
- Overduin, P.P., Schneider von Deimling, T., Miesner, F., Grigoriev, M.N., Ruppel, C., Vasiliev, A. et al. (2019). Submarine permafrost map in the Arctic modeled using 1-D transient heat flux (SuPerMAP). Journal of Geophysical Research: Oceans 124(6), 3490–3507. https://doi.org/10.1029/2018JC014675.
- Paull, C.K., Dallimore, S.R., Jin, Y.K., Caress, D.W., Gwiazda, R., Anderson, K. et al. (2022). Rapid seafloor changes associated

with the degradation of Arctic submarine permafrost. PNAS 119(12), e2119105119. https://doi.org/10.1073/pnas.2119105119.

Arctic Ripples

Feeling the heat

Text

- de Grandpré, I., Fortier, D. and Stephani, E. (2012). Degradation of permafrost beneath a road embankment enhanced by heat advected in groundwater. Canadian Journal of Earth Sciences 49(8), 953–962. https://cdnsciencepub.com/doi/10.1139/ e2012-018.
- Hjort, J., Streletskiy, D., Doré, G., Wu, Q., Bjella, K. and Luoto, M. (2022). Impacts of permafrost degradation on infrastructure. Nature Reviews Earth & Environment 3, 24–38. https://www. nature.com/articles/s43017-021-00247-8.
- Hjort, J., Karjalainen, O., Aalto, J., Westermann, S., Romanovsky, V.E., Nelson, F.E. et al. (2018). Degrading permafrost puts Arctic infrastructure at risk by mid-century. Nature Communications 9, 5147. https://www.nature.com/articles/ \$41467-018-07557-4.
- Streletskiy, D.A., Shiklomanov, N.I. and Nelson, F.E. (2012). Permafrost, infrastructure, and climate change: A GIS-based landscape approach to geotechnical modeling. Arctic, Antarctic, and Alpine Research 44(3), 368–380. https://www. tandfonline.com/doi/full/10.1657/1938-4246-44.3.368.
- Streletskiy, D.A., Suter, L.J., Shiklomanov, N.I., Porfiriev, B.N. and Eliseev, D.O. (2019). Assessment of climate change impacts on buildings, structures and infrastructure in the Russian regions on permafrost. Environmental Research Letters 14(2), 025003. https://iopscience.iop.org/ article/10.1088/1748-9326/aaf5e6
- Ramage, J., Jungsberg, L., Wang, S., Westermann, S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https://link. springer.com/article/10.1007/s11111-020-00370-6.
- Shur, Y. and Goering, D.J. (2009). Climate change and foundations of buildings in permafrost regions. In Permafrost Soils. Margesin, R. (ed.). Soil Biology 16, 251–260. Berlin and Heidelberg: Springer. https://link.springer.com/ chapter/10.1007/978-3-540-69371-0_17.

Maps

Obu, J., Westermann, S., Barboux, C., Bartsch, A., Delaloye, R., Grosse, G. et al. (2021). ESA Permafrost Climate Change Initiative (Permafrost_cci): Permafrost Active Layer Thickness for the Northern Hemisphere, v3.0. NERC EDS Centre for Environment Data Analysis. https://catalogue.ceda.ac.uk/ uuid/67a3f8c8dc914ef99f7f08ebod997e23. Accessed 20 June 2023. DOI: 10.5285/67a3f8c8dc914ef99f7f08ebod997e23 Ramage, J., Jungsberg, L., Wang, S., Westermann. S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https:// doi.org/10.1007/S1111-020-00370-6.

Risky business I

Text

- Allard, M., Lemay, M., Barrette, C., L'Hérault, E. and Sarrazin, D. (2012). Permafrost and climate change in Nunavik and Nunatsiavut: Importance for municipal and transportation infrastructures. In Nunavik and Nunatsiavut: From Science to Policy – An Integrated Regional Impact Study (IRIS) of Climate Change and Modernization. Allard, M. and Lemay, M. (eds.). Quebéc, Canada: ArcticNet Incorporated. Chapter 6. 171–197. https://www.researchgate.net/publication/264454915_ Chapter_6_Permafrost_and_climate_change_in_Nunavik_ and_Nunatsiavut_Importance_for_municipal_and_ transportation_infrastructures. Accessed 30 June 2023.
- De Guzman, E.M.B., Alfaro, M.C., Doré, G., Arenson, L.U. and Piamsalee, A. (2021). Performance of highway embankments in the Arctic constructed under winter conditions. Canadian Geotechnical Journal 58(5), 722–736. https://cdnsciencepub. com/doi/pdf/10.1139/cgj-2019-0121.
- Calmels, F., Roy, L.-P., Grandmont, K. and Pugh R. (2018). Summary of Climate- and Geohazard-related Vulnerabilities for the Dempster Highway Corridor. Whitehorse, Yukon: Yukon University Research Centre. https://yukonu.arcabc.ca/ islandora/object/yukonu%3A28/datastream/PDF/view.
- Calmels, F., Roy, L.-P., Laurent, C., Pelletier, M., Kinnear, L., Benkert, B. et al. (2015). Vulnerability of the North Alaska Highway to Permafrost Thaw: A Field Guide and Data Synthesis. Halladay, P. (ed). Whitehorse, Yukon: Yukon University Research Centre. https://www.yukonu.ca/sites/ default/files/inline-files/permafrost_report.pdf.
- Hivon, E.G. and Sego, D.C. (1993). Distribution of saline permafrost in the Northwest Territories, Canada. Canadian Geotechnical Journal 30(3), 506–514. https://cdnsciencepub. com/doi/10.1139/t93-043.
- Hjort, J., Streletskiy, D., Doré, G., Wu, Q., Bjella, K. and Luoto, M. (2022). Impacts of permafrost degradation on infrastructure. Nature Reviews Earth & Environment 3, 24–38. https://www. nature.com/articles/s43017-021-00247-8.
- Hjort, J., Karjalainen, O., Aalto, J., Westermann, S., Romanovsky, V.E., Nelson, F.E. et al. (2018). Degrading permafrost puts Arctic infrastructure at risk by mid-century. Nature Communications 9, 5147. https://www.nature.com/articles/s41467-018-07557-4.
- Ingeman-Nielsen, T., Kandiliotis, P., Lola, M., Foged, N. and Stenseng, L. (2018). Kangerlussuaq Airport Deformations and construction failures in Arctic Permafrost areas. ARTEC International Conference AIC 2018: Transportation Infrastructure

Engineering in Cold Regions. Sisimut, Greenland, 1-3 May 2018. 96–97. https://orbit.dtu.dk/en/publications/kangerlussuaqairport-deformations-and-construction-failures-in-a.

- Jungsberg, L., Herslund, L.B., Nilsson, K., Wang, S., Tomaškovičová, S., Madsen, K. et al. (2021). Adaptive capacity to manage permafrost degradation in Northwest Greenland. Polar Geography 45(1), 58–76. https://www. tandfonline.com/doi/full/10.1080/1088937X.2021.1995067
- Lamoureux, S., Forbes, D.L., Bell, T., Manson, G.K. et al. (2015). The impact of climate change on infrastructure in the western and central Canadian Arctic. In: From Science to Policy in the Western and Central Canadian Arctic: an Integrated Regional Impact Study (IRIS) of Climate Change and Modernization. Stern, G. and Gaden, A., (eds). Québec, Canada: ArcticNet Incorporated. Chapter 7. 300-341. https://arcticnet.ulaval.ca/ wp-content/uploads/2022/06/IRIS_FromScience_ArcticNet_ Ir_.pdf. Accessed 24 February 2023.
- Lemay, M., Allard, M., Bell, T. et al. (2018). Living on permafrost: background and case studies for adapting community infrastructure in a changing climate. In: From Science to Policy in the Eastern Canadian Arctic: an Integrated Regional Impact Study (IRIS) of Climate Change and Modernization. Bell, T. and Brown, T., (eds). Québec, Canada: ArcticNet, Incorporated. Chapter 15. 210-249, https://arcticnet.ulaval. ca/wp-content/uploads/2022/06/29170_IRIS_East_fullreport web compressed.pdf. Accessed 24 February 2023.
- Ma, D., Motagh, M., Liu, G., Zhang, R., Wang, X., Zhang, B. et al. (2022). Thaw settlement monitoring and active layer thickness retrieval using time series COSMO-SkyMed imagery in Iqaluit Airport. Remote Sensing 14(9), 2156. https://www. mdpi.com/2072-4292/14/9/2156.
- Melvin, A.M., Larsen, P., Boehlert, B., Neumann, J.E., Chinowsky, P., Espinet, X. et al. (2016). Climate change damages to Alaska public infrastructure and the economics of proactive adaptation. PNAS 114(2), E122–E131. https://www.pnas.org/ doi/epdf/10.1073/pnas.1611056113.
- Ramage, J., Jungsberg, L., Wang, S., Westermann, S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https:// doi.org/10.1007/S11111-020-00370-6.
- Romanovsky, V. (2011). High-resolution permafrost modeling and planning implications for Greenland and Alaska, 31 May. https://www.arcus.org/witness-the-arctic/2011/2/ article/1651. Accessed 7 June 2023.
- Streletskiy, D.A., Clemens, S., Lanckman, J.P. and Shiklomanov, N.I. (2023). The costs of Arctic infrastructure damages due to permafrost degradation. Environmental Research Letters 18, 014006. https://iopscience.iop.org/ article/10.1088/1748-9326/acab18/meta
- Suter, L., Streletskiy, D. and Shiklomanov, N. (2019). Assessment of the cost of climate change impacts on critical

infrastructure in the circumpolar Arctic. Polar Geography 42(4), 267–286. https://www.tandfonline.com/doi/abs/10.10 80/1088937X.2019.1686082.

Maps

- Hjort, J., Karjalainen, O., Aalto, J., Westermann, S., Romanovsky, V.E., Nelson, F.E. et al. (2018). Degrading permafrost puts Arctic infrastructure at risk by mid-century. Nature Communications 9, 5147. https://doi.org/10.1038/s41467-018-07557-4.
- Karjalainen, O., Aalto, J., Luoto, M., Westermann, S., Romanovsky, V.E., Nelson, F.E. et al. (2019). Circumpolar permafrost maps and geohazard indices for near-future infrastructure risk assessments. Scientific Data 6, 190037. https://doi.org/10.1038/sdata.2019.37.
- Ramage, J., Jungsberg, L., Wang, S., Westermann. S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https:// doi.org/10.1007/S11111-020-00370-6.

Risky business II

Text

Hjort, J., Streletskiy, D., Doré, G., Wu, Q., Bjella, K. and Luoto, M. (2022). Impacts of permafrost degradation on infrastructure. Nature Reviews Earth & Environment 3, 24–38. https://www. nature.com/articles/s43017-021-00247-8.

- Kotov, P.I. and Khilimonyuk, V.Z. (2021). Building stability on permafrost in Vorkuta, Russia. Geography, Environment, Sustainability 14(4), 67–74. https://ges.rgo.ru/jour/article/ view/2069.
- Shiklomanov, N.I., Streletskiy, D.A., Swales, T.B. and Kokorev, V.A. (2017). Climate change and stability of urban infrastructure in Russian permafrost regions: Prognostic assessment based on GCM climate projections. Geographical Review 107(1), 125–142. https://www.tandfonline.com/doi/full/10.1111/gere.12214.
- Shur, Y. and Goering, D.J. (2009). Climate change and foundations of buildings in permafrost regions. In Permafrost Soils. Margesin, R. (ed.). Soil Biology 16, 251–260. Berlin and Heidelberg: Springer. https://link.springer.com/ chapter/10.1007/978-3-540-69371-0_17.

Maps

Hjort, J., Karjalainen, O., Aalto, J., Westermann, S., Romanovsky, V.E., Nelson, F.E. et al. (2018). Degrading permafrost puts Arctic infrastructure at risk by mid-century. Nature Communications 9, 5147. https://doi.org/10.1038/s41467-018-07557-4.

Karjalainen, O., Aalto, J., Luoto, M., Westermann, S., Romanovsky, V.E., Nelson, F.E. et al. (2019). Circumpolar permafrost maps and geohazard indices for near-future infrastructure risk assessments. Scientific Data 6, 190037. https://doi.org/10.1038/sdata.2019.37.

Ramage, J., Jungsberg, L., Wang, S., Westermann. S., Lantuit, H. and Heleniak, T. (2021). Population living on permafrost in the Arctic. Population and Environment 43, 22–38. https://doi. org/10.1007/S11111-020-00370-6.

Terra infirma I

Text

Barnhart, K.R., Overeem, I. and Anderson, R.S. (2014). The effect of changing sea ice on the physical vulnerability of Arctic coasts. Cryosphere 8(5), 1777–1799. https://tc.copernicus. org/articles/8/1777/2014/.

- Bartsch, A., Pointner, G., Nitze, I., Efimova, A., Jakober, D., Ley, S. et al. (2021). Expanding infrastructure and growing anthropogenic impacts along Arctic coasts. Environmental Research Letters 16, 115013. https://iopscience.iop.org/ article/10.1088/1748-9326/ac3176/pdf.
- Chuvilin, E.M., Yakushev, V.S. and. Perlova, E.V. (2000). Gas and possible gas hydrates in the permafrost of Bovanenkovo gas field, Yamal Peninsula, West Siberia. Polarforschung 68, 215–219. https://core.ac.uk/download/pdf/11771406.pdf.
- Chuvilin, E., Tipenko, G., Bukhanov, B., Istomin, V. and Pissarenko, D. (2022). Simulating thermal interaction of gas production wells with relict gas hydrate-bearing permafrost. Geosciences 12(3), 115. https://www.mdpi.com/2076-3263/12/3/115.
- Fritz, M., Vonk, J.E. and Lantuit, H. (2017). Collapsing Arctic coastlines. Nature Climate Change 7, 6–7. https://www.nature.com/articles/nclimate3188.
- Leibman, M., Khomutov, A. and Kizyakov, A. (2014). Cryogenic landslides in the west-Siberian plain of Russia: Classification, mechanisms, and landforms. In Landslides in Cold Regions in the Context of Climate Change. Shan, W., Guo, Y., Wang, F., Marui, H., Strom, A. (eds). Cham, Switzerland: Springer. 143–162. https://doi.org/10.1007/978-3-319-00867-7 11.
- Tsentr, A. (2020). TEK Rossii—2019 [Russia's Fuel and Energy Complex 2019]. Analytical Center for the Government of the Russian Federation.
- Vasiliev, A. (2021). An Arctic success story: Yamal Nenets Autonomous Region, 9 March. https://www.arcticcircle.org/ journal/an-arctic-success-story-yamal-nenets-autonomousregion. Accessed 7 June 2023.
- Yakushev, V.S. (2020). Natural gas liberations around production wells at Russian Arctic gas fields. Geosciences 10(5), 184. https://www.mdpi.com/2076-3263/10/5/184.

Maps

Bartsch, A., Pointner, G., Nitze, I., Efimova, A., Jakober, D., Ley, S. et al. (2021). Expanding infrastructure and growing anthropogenic impacts along Arctic coasts. Environmental Research Letters 16(11), 115013. https://dx.doi.org/10.1088/1748-9326/ac3176.
Obu, J., Westermann, S., Barboux, C., Bartsch, A., Delaloye,

R., Grosse, G. et al. (2021). ESA Permafrost Climate Change

Initiative (Permafrost_cci): Permafrost Active Layer Thickness for the Northern Hemisphere, v3.o. NERC EDS Centre for Environment Data Analysis. https://catalogue.ceda.ac.uk/ uuid/67a3f8c8dc914ef99f7f08ebod997e23. Accessed 20 June 2023. DOI: 10.5285/67a3f8c8dc914ef99f7f08ebod997e23. Protection of the Arctic Marine Environment (2021). Arctic Ship

Traffic Data for 2 March 2021. Protection of the Arctic Marine Environment. https://pame.is/index.php/projects/arcticmarine-shipping/astd. Accessed 20 June 2023.

Terra infirma II

Text

- Bouchard, C. (2020). Arctic Airports and Aerodromes As Critical Infrastructure. Peterborough: North American Arctic Defence and Security Network. https://www.naadsn.ca/wp-content/ uploads/2020/11/Airports_CI_2020_11_05.pdf.
- Tanguy, R., Whalen, D., Prates, G., Pina, P., Freitas, P., Bergstedt, H., and Vieira, G. (2023). Permafrost degradation in the ice-wedge tundra terrace of Paulatuk Peninsula (Darnley Bay, Canada). Geomorphology 435, 108754. https://doi. org/10.1016/j.geomorph.2023.108754
- Tanguy, R., Whalen, D., Prates, G., and Vieira, G. (2023). Shoreline change rates and land to sea sediment and soil organic carbon transfer in eastern Parry Peninsula from 1965 to 2020 (Amundsen Gulf, Canada). Arctic Science 9(3), 506-525. https://doi.org/10.1139/AS-2022-0028

Maps

Vieira, G., Whalen, D., Pina, P., Freitas, P. and Pinheiro,
D. (Unpublished). Paulatuk ultra-high resolution aerial orthomosaic, July 2019. Centre for Geographical Studies of the Institute of Geography and Spatial Planning, University of Lisbon and National Resources Canada.

Terra infirma III

- Brown, J. (1969). Ionic Concentration Gradients in Permafrost, Barrow, Alaska. Hanover, New Hampshire, US: United States Army Corps of Engineers, Cold Regions Research and Engineering Laboratory. https://erdc-library.erdc.dren.mil/jspui/ bitstream/11681/5827/1/CRREL-Research-Report-272.pdf.
- D'Oro, R. (2019). Failing ice cellars signal changes in Alaska whaling towns, 25 November. https://apnews.com/article/usnews-ap-top-news-whales-climate-change-weekend-readsea84d035f9ff46d099ef63fc21b95a2f. Accessed 7 June 2023.
- Frost, K. (2021). Climate change is melting Arctic ice cellars, 1 June. https://civileats.com/2021/06/01/climate-change-ismelting-arctic-ice-cellars/. Accessed 7 June 2023.
 Maslakov, A.A., Nyland, K.E., Komova, N.N., Yurov, F.D.,

Yoshikawa, K. and Kraev, G.N. (2020). Community ice cellars In Eastern Chukotka: Climatic and anthropogenic influences on structural stability. Geography, Environment, Sustainability 13(3), 49–56. https://ges.rgo.ru/jour/ article/view/1192.

- Nyland, K.E., Klene, A.E., Brown, J., Shiklomanov, N.I., Nelson, F.E., Streletskiy, D.A. et al. (2017). Traditional Iñupiat ice cellars (SIĠlUAQ) in Barrow, Alaska: Characteristics, temperature monitoring, and distribution. Geographical Review 107(1), 143–158. https://www.tandfonline.com/doi/ full/10.1111/j.1931-0846.2016.12204.x.
- Ray, P. H., Murdoch, J., Riley, C. V., Gray, A., Dall, W. H. and Schott, C. A. (1885). Report of the International Polar Expedition to Point Barrow, Alaska. Washington, D.C.: United States Government Printing Office. https://www.biodiversitylibrary. org/item/76077#page/9/mode/1up.

Maps

Yoshikawa, K., Maslakov, A.A., Kraev, G., Ikuta, H., Romanovsky, V.E., George, J.C. et al. (2022). Food storage in permafrost and seasonally frozen ground in Chukotka and Alaska communities. Archives 75(2), 149–290. https://doi. org/10.14430/arctic75259.

Terra infirma IV

Text

- Avannaata Kommunia (2021). Urban Development Ilulissat Nord: Nordre Næs and Kangerluarsuk / Brede Bay Sites. Ilulissat, Greenland. https://www.avannaata.gl/-/media/ avannaata/filer/nyheder/2022/strukturplan pdf?la=da&hash= DFD68E806DF3399F83B5B4BDFDC6099.
- Finnsson, P.T. (2018). The threat of permafrost thaw is looming in the Arctic, 15 October. https://nordregio.org/nordregio-magazine/ issues/arctic-changes-and-challenges/the-threat-of-permafrostthaw-is-looming-in-the-arctic/. Accessed 8 June 2023.

Ingemann-Nielsen, personal communication

- Ingeman-Nielsen, T. (2018). Oedometer tests on frozen cores from Ilulissat, Greenland. Lyngby, Denmark: Technical University of Denmark, Department of Civil Engineering.
- United Nations Educational, Scientific and Cultural Organization (2004). Ilulissat Icefjord. https://whc.unesco.org/en/list/ 1149/. Accessed 8 June 2023.

Maps

- ASIAQ Greenland Survey (2021). The technical base map. Scale: 1: 2,750 and 1: 5,000. ASIAQ Greenland Survey. https:// www.asiaq-greenlandsurvey.gl/kortlaegning-geografiskinformation/the-technical-base-map/. Accessed 20 June 2023.
- Porter, C., Morin, P., Howat, I., Noh, M.-J., Bates, B., Peterman, K. et al. (2018). ArcticDEM, Version 3. Harvard Dataverse. https://

doi.org/10.7910/DVN/OHHUKH. Accessed 20 June 2023. Scheer, J., Caduff, R., How, P., Marcer, M., Strozzi, T., Bartsch, A. et al. (2023). Thaw-season InSAR surface displacements and frost susceptibility mapping to support community-scale planning in Ilulissat, West-Greenland. Preprints.org 2023051683. https://doi.org/10.20944/ preprints202305.1683.V1.

Nothing in isolation

Text

Brubaker, M., Berner, J., Chavan, R. and Warren, J. (2011). Climate change and health effects in Northwest Alaska. Global Health Action 4(1), 8445. https://www.tandfonline. com/doi/full/10.3402/gha.v4i0.8445.

- Cunsolo Willox, A., Harper, S.L., Ford, J.D., Edge, V.L., Landman, K., Houle, K. et al. (2013). Climate change and mental health: an exploratory case study from Rigolet, Nunatsiavut, Canada. Climatic Change 121, 255–270. https://link.springer.com/ article/10.1007/s10584-013-0875-4.
- Healey, G.K. (2017). What if our health care systems embodied the values of our communities? A reflection from Nunavut, 20 June. https://www.thearcticinstitute.org/health-care-systemsvalues-communities-nunavut/. Accessed 8 June 2023.
- Ruscio, B.A., Brubaker, M., Glasser, J., Hueston, W. and Hennessy, T.W. (2015). One Health – A strategy for resilience in a changing Arctic. International Journal of Circumpolar Health 74(1), 27913. https://www.tandfonline.com/doi/ full/10.3402/ijch.v74.27913.
- Timlin, U., Ingimundarson, J.H., Jungsberg, L., Kauppila, S., Larsen, J.N., Nordström, T. et al. (2021). Living conditions and mental wellness in a changing climate and environment: Focus on community voices and perceived environmental and adaptation factors in Greenland. Heliyon 7(4), E06862. https://www.cell.com/heliyon/fulltext/S2405-8440(21)00965-8.
- Timlin, U., Kauppila, S., Jungsberg, L., Nordström, T., Schmidt-Pedersen, K., Kyngäs, H. et al. (2021). Perception of health challenges, self-rated health and feeling of empowerment in a changing climate and environment with permafrost thawing. Sci 3(2), 28. https://www.mdpi.com/2413-4155/3/2/28.
- Timlin, U., Meyer, A., Nordström, T. and Rautio, A. (2022). Permafrost thaw challenges and life in Svalbard. Current Research in Environmental Sustainability 4, 100122. https://www.sciencedirect.com/science/article/pii/ S2666049021001006.
- Timlin, U., Ramage, J., Gartler, S., Nordström, T. and Rautio, A. (2022). Self-rated health, life balance and feeling of empowerment when facing impacts of permafrost thaw – A case study from Northern Canada. Atmosphere 13(5), 789. https://www.mdpi.com/2073-4433/13/5/789.

Toxic grounds

- Arctic Monitoring and Assessment Programme (2013). Transport of contaminants to the Arctic - atmospheric and aquatic transport. https://www.amap.no/documents/doc/transportof-contaminants-to-the-arctic-atmospheric-and-acquatictransport/944. Accessed 8 June 2023.
- Arctic Monitoring and Assessment Programme (2018). AMAP Assessment 2018: Biological Effects of Contaminants on Arctic Fish and Wildlife. Tromsø, Norway. https://www. amap.no/documents/doc/amap-assessment-2018biological-effects-of-contaminants-on-arctic-wildlife-andfish/166.
- Arctic Monitoring and Assessment Programme (2021). POPs and Chemicals of Emerging Arctic Concern: Influence of Climate Change. Summary for Policymakers. Tromsø, Norway. https://www.amap.no/documents/doc/pops-and-chemicalsof-emerging-arctic-concern-influence-of-climate-change.summary-for-policy-makers/3511.
- Dastoor, A., Angot, H., Bieser, J., Christensen, J.H., Douglas, T.A., Heimbürger-Boavida, L.-E. et al. (2022). Arctic mercury cycling. Nature Reviews Earth & Environment 3, 270–286. https://www.nature.com/articles/s43017-022-00269-w.
- Langer, M., von Deimling, T.S., Westermann, S., Rolph, R., Rutte, R., Antonova, S. et al. (2023). Thawing permafrost poses environmental threat to thousands of sites with legacy industrial contamination. Nature Communications 14, 1721. https://www.nature.com/articles/s41467-023-37276-4.
- Lim, A.G., Jiskra, M., Sonke, J.E., Loiko, S.V., Kosykh, N. and Pokrovsky, O.S. (2020). A revised pan-Arctic permafrost soil Hg pool based on Western Siberian peat Hg and carbon observations. Biogeosciences 17(12), 3083–3097. https:// bg.copernicus.org/articles/17/3083/2020/.
- Rudnicka-Kępa, P. and Zaborska, A. (2021). Sources, fate and distribution of inorganic contaminants in the Svalbard area, representative of a typical Arctic critical environment – A review. Environmental Monitoring and Assessment 193, 724. https://link.springer.com/article/10.1007/S10661-021-09305-6.
- Schuster, P.F., Schaefer, K.M., Aiken, G.R., Antweiler, R.C., Dewild, J.F., Gryziec, J.D. et al. (2018). Permafrost stores a globally significant amount of mercury. Geophysical Research Letters 45(3), 1463–1471. https://agupubs.onlinelibrary.wiley. com/doi/10.1002/2017GL075571.
- Wang, X., Yuan, W., Lin, C.-J., Zhang, L., Zhang, H. and Feng, X. (2019). Climate and vegetation as primary drivers for global mercury storage in surface soil. Environmental Science & Technology 53(18), 10665–10675. https://pubs.acs.org/ doi/10.1021/acs.est.9b02386.

Maps

Schuster, P.F., Schaefer, K.M., Aiken, G.R., Antweiler, R.C., Dewild, J.F., Gryziec, J.D. et al. (2018). Permafrost stores a globally significant amount of mercury. Geophysical Research Letters 45(3), 1463–1471. https://doi. org/10.1002/2017GL075571.

Coming back to life

Text

- Biagini, P., Thèves, C., Géraut, A., Keyser, C., Nikolaeva, D., Gérard, P. et al. (2012). Variola virus in a 300-year-old Siberian mummy. The New England Journal of Medicine 367, 2057– 2059. https://www.nejm.org/doi/full/10.1056/NEJMc1208124.
- Carrasco-Hernandez, R., Jácome, R., López Vidal, Y. and Ponce de León, S. (2017). Are RNA viruses candidate agents for the next global pandemic? A Review. ILAR Journal 58(3), 343–358. https://academic.oup.com/ilarjournal/ article/58/3/343/4107390.
- Charlier, P., Héry-Arnaud, G., Coppens, Y., Malaurie, J., Hoang-Oppermann, V., Deps, P. et al. (2020). Global warming and planetary health: An open letter to the WHO from scientific and indigenous people urging for paleo-microbiology studies. Infection, Genetics and Evolution 82, 104284. https://www. sciencedirect.com/science/article/abs/pii/S1567134820301155.
- Christie, A. (2021). Blast from the past: Thawing permafrost could lead to future pandemics. Cambridge Journal of Science & Policy 2(2), 8 https://doi.org/10.17863/CAM.74501.
- Duffy, S. (2018). Why are RNA virus mutation rates so damn high? PLOS Biology 16(8), e300003. https://journals.plos. org/plosbiology/article?id=10.1371/journal.pbio.300003.
- Enderle, R., Stenlid, J. and Vasaitis, R. (2019). An overview of ash (Fraxinus spp.) and the ash dieback disease in Europe. CABI Reviews 14, 1–12. https://www.cabidigitallibrary.org/ doi/10.1079/PAVSNNR201914025.
- Ezhova, E., Orlov, D., Suhonen, E., Kaverin, D., Mahura, A., Gennadinik, V. et al. (2021). climatic factors influencing the anthrax outbreak of 2016 in Siberia, Russia. EcoHealth 18, 217–228. https://link.springer.com/article/10.1007/s10393-021-01549-5.
- Legendre, M., Lartigue, A., Bertaux, L., Jeudy, S., Bartoli, J., Lescot, M. et al. (2015). In-depth study of Mollivirus sibericum, a new 30,000-y-old giant virus infecting Acanthamoeba. PNAS 112(38), E5327–E5335. https://doi.org/10.1073/pnas.1510795112.
- Revich, B., Tokarevich, N. and Parkinson, A. J. (2012). Climate change and zoonotic infections in the Russian Arctic. International Journal of Circumpolar Health 71(1), 18792. https://doi.org/10.3402/ijch.v71i0.18792.
- Schütte, U.M.E., Henning, J.A., Ye, Y., Bowling, A., Ford, J., Genet, H. et al. (2019). Effect of permafrost thaw on plant and soil fungal community in a boreal forest: Does fungal community change

mediate plant productivity response? Journal of Ecology 107(4), 1737–1752. https://doi.org/10.1111/1365-2745.13139.

Stocks, J.J., Metheringham, C.L., Plumb, W.J., Lee, S.J., Kelly, L.J., Nichols, R.A. et al. (2019). Genomic basis of European ash tree resistance to ash dieback fungus. Nature Ecology & Evolution 3, 1686–1696. https://doi.org/10.1038/s41559-019-1036-6.
Taubenberger, J.K., Hultin, J.V. and Morens, D.M. (2007). Discovery and characterization of the 1918 pandemic influenza virus in historical context. Antiviral Therapy 12(4 Pt B), 581–91. https://pubmed.ncbi.nlm.nih.gov/17944266/.

Maps

- Arctic Council, Conservation of Arctic Flora and Fauna Working Group (2001). CAFF Map No. 28 – Distribution of wild reindeer, caribou, and domestic reindeer in the Arctic. Arctic Portal Library. http://library.arcticportal.org/1358/.
- Stella, E., Mari, L., Gabrieli, J., Barbante, C. and Bertuzzo, E. (2021). Mapping environmental suitability for anthrax reemergence in the Arctic. Environmental Research Letters 16(10), 105013. https:// iopscience.iop.org/article/10.1088/1748-9326/ac2527/meta.

Frozen assets I

Text

- Hope, C. and Schaefer, K. (2016). Economic impacts of carbon dioxide and methane released from thawing permafrost. Nature Climate Change 6, 56–59. https://doi.org/10.1038/nclimate2807.
- Jungsberg, L., Ramage, J., Cordier, M., Vanderlinden, J-P., Vasivlevskaya, A. (2021). 7.5 Report on Permafrost Thawing, Natural Resource-based Economy, and Arctic Coastal Communities' Ability to Face the Associated Risks: An Integrated Strategic Framework. Nunataryuk: WP7–Natural Resources, Economy and Coastal Community Planning.
- Larsen, J.N. and Fondahl, G. (eds.) (2015). Arctic Human Development Report: Regional Processes and Global Linkages. Copenhagen: Nordic Council of Ministers. http://norden.divaportal.org/smash/get/diva2:788965/FULLTEXT03.pdf.
- Melvin, A.M., Larsen, P., Boehlert, B., Neumann, J.E., Chinowsky, P., Espinet, X. et al. (2016). Climate change damages to Alaska public infrastructure and the economics of proactive adaptation. PNAS 114(2), E122–E131. https://doi.org/10.1073/ pnas.1611056113.
- Yumashev, D., Hope, C., Shaefer, K., Riemann-Campe, K., Iglesias-Suarez, F., Jafarov, E. et al. (2019). Climate policy implications of nonlinear decline of Arctic land permafrost and other cryosphere elements. Nature communications 10(1900). https://doi.org/10.1038/s41467-019-09863-x

Maps

Jungsberg, L., Ramage, J., Cordier, M., Vanderlinden, J-P., Vasivlevskaya, A. (eds.) (2021). 7.5 Report on Permafrost Thawing, Natural Resource-based Economy, and Arctic Coastal Communities' Ability to Face the Associated Risks: An Integrated Strategic Framework. Deliverable Report, Grant Agreement number: 773421, H2020-BG-2016-2017/H2020-BG-2017-1. Nordregio, Stockholm.

Frozen assets II

Text

- Glomsrød, S., Duhaime, G. and Aslaksen, I. (eds.) (2017). The Economy of the North 2015. Oslo: Statistics Norway. https:// www.ssb.no/en/natur-og-miljo/artikler-og-publikasjoner/_ attachment/299576?_ts=15aea97d740.
- Jungsberg, L., Ramage, J., Cordier, M., Vanderlinden, J-P., Vasivlevskaya, A. (2021). 7.5 Report on Permafrost Thawing, Natural Resource-based Economy, and Arctic Coastal Communities' Ability to Face the Associated Risks: An Integrated Strategic Framework. Nunataryuk: WP7–Natural Resources, Economy and Coastal Community Planning.
- Ramage, J., Jungsberg, L., Wang, S., Gartler, S., Meyer, A., Cordier, M. et al. (2020). Synthesis Report: Identifying and Recognizing the Value of Provisioning Permafrost Ecosystem Services in the Arctic. Nunataryuk: WP7–Natural Resources, Economy and Coastal Community Planning.

Maps

- Northwest Territories Bureau of Statistics (2018). Households Eating Meat or Fish Obtained from Hunting or Fishing by Community (1998 to 2018). https://www.statsnwt.ca/ Traditional%20Activities/. Accessed 20 June 2023.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi.org/10.1016/j.earscirev.2019.04.023.
- Ramage, J., Jungsberg, L., Meyer, A. and Gartler, S. (2022). 'No longer solid': Perceived impacts of permafrost thaw in three Arctic communities. Polar Geography 45(3), 226–239. https:// doi.org/10.1080/1088937X.2022.2105973.

Cultural homeland

- Crate, S.A. (2006). Cows, Kin and Globalization. An Ethnography of Sustainability. Walnut Creek. Walnut Creek, United States: AltaMira Press. https://rowman.com/ISBN/9780759107403/ Cows-Kin-and-Globalization-An-Ethnography-of-Sustainability.
- Crate, S.A. (2016). Gone with cows and kin? Climate, globalization, and youth alienation in Siberia. In Anthropology and Climate Change: From Actions to Transformations. Crate, S.A. and Nuttall, M. (eds.). Oxford, United Kingdom:

Routledge. 139–161. https://www.taylorfrancis.com/chapters/ edit/10.4324/9781315530338-11/gone-cows-kin-climateglobalization-youth-alienation-siberia.

- Crate, S.A. (2021). Once Upon the Permafrost: Knowing Culture And Climate Change in Siberia. Tucson, United States: University of Arizona Press. https://uapress.arizona.edu/ book/once-upon-the-permafrost.
- Crate, S., Ulrich, M., Habeck, J.O., Desyatkin, A.R., Desyatkin, R.V., Fedorov, A.N. et al. (2017). Permafrost livelihoods: A transdisciplinary review and analysis of thermokarst-based systems of indigenous land use. Anthropocene 18, 89–104. https://doi.org/10.1016/j.ancene.2017.06.001.
- Czerniawska, J., and Chlachula, J. (2020). Climate-change induced permafrost degradation in Yakutia, East Siberia. Arctic 73(4), 509–528. https://www.jstor.org/stable/26991438.
- Fedorov, A. N. (2022). Permafrost landscape research in the northeast of Eurasia. Earth 3 (1), 460–478. https://doi. org/10.3390/earth3010028eds.
- Gavril'ev, P.P. (1991). Melioratsiia merzlotnykh zemel' v lakutii [Melioration of permafrost grounds in Yakutia} Nauka, Novosibirsk. (In Russian)
- Gabyshev, M. et al. (1929). O zemel'noi reforme v lakutii: k VI vseiakutskomu s"ezdu sovetov [Russian edition].
- Hubberten, H.W., Andreev, A., Astakhov, V.I., Demidov, I., Dowdeswell, J.A., Henriksen, M. et al. (2004). The periglacial climate and environment in northern Eurasia during the Last Glaciation. Quaternary Science Reviews 23(11-13), 1333–1357. https://doi.org/10.1016/j.quascirev.2003.12.012.
- Maj, E. (2009). The horse of Sakha: Ethnic symbol in postcommunist Sakha Republic (lakutiia). Sibirica 8 (1), 68–74. https://doi.org/10.3167/sib.2009.080105.
- Matveev, I.A. (1989). Atlas selskogo khozyaistva Yakutskoi ASSR. [Agricultural atlas of the Yakut ASSR]. Moskow: Glavnoe upravlenie geodezii i kartographii pri sovete ministerstvo SSSR. https://www.loc.gov/item/2018694181/.
- Pekarski, E.K. 1958 [1899]. Slovar lakutskogo lazyka. [Dictionary of the Sakha language]. St Petersburg: Academy of Science.
- Solovyev, P.A. (1973). Thermokarst phenomena and landforms due to frost heaving in central Yakutia. Periglacial Bulletin 23, 135–155.
- Takakura, H. 2015. Arctic Pastoralist Sakha: Ethnography of Evolution and Microadaptation in Siberia. Melbourne: Trans Pacific Press. https://transpacificpress.com/products/arcticpastoralist-sakha?variant=32554245324885.
- Takakura, H. (2016). Limits of pastoral adaptation to permafrost regions caused by climate change among the Sakha people in the middle basin of Lena River. Polar Science 10(3), 395–403. https://doi.org/10.1016/j.polar.2016.04.003.

Maps

Crate, S., Ulrich, M., Habeck, J.O., Desyatkin, A.R., Desyatkin, R.V., Fedorov, A.N. et al. (2017). Permafrost livelihoods: A transdisciplinary review and analysis of thermokarst-based systems of indigenous land use. Anthropocene 18, 89–104. https://doi.org/10.1016/j.ancene.2017.06.001.

European Union Agency for the Space Programme (2020). Copernicus Data Space Ecosystem: Satellite Imagery for 3 June 2022. European Union Agency for the Space Programme. https://dataspace.copernicus.eu/. Accessed 20 June 2023.

Holding Tight

Bumpy road ahead

Text

- Allard, M., Lemay, M., Barrette, C., L'Hérault, E. and Sarrazin, D. (2012). Permafrost and climate change in Nunavik and Nunatsiavut: Importance for municipal and transportation infrastructures. In Nunavik and Nunatsiavut: From Science to Policy – An Integrated Regional Impact Study (IRIS) of Climate Change and Modernization. Allard, M. and Lemay, M. (eds.). Quebéc, Canada: ArcticNet Incorporated. Chapter 6. 171–197. https://www.researchgate.net/publication/264454915_ Chapter_6_Permafrost_and_climate_change_in_Nunavik_ and_Nunatsiavut_Importance_for_municipal_and_ transportation_infrastructures. Accessed 30 June 2023.
- Bartsch, A., Pointner, G., Nitze, I., Efimova, A., Jakober, D., Ley, S. et al. (2021). Expanding infrastructure and growing anthropogenic impacts along Arctic coasts. Environmental Research Letters 16(11), 115013. https://iopscience.iop.org/ article/10.1088/1748-9326/ac3176/pdf.
- De Guzman, E.M. (2020). Structural Stability of Highway Embankments in the Arctic Corridor. Winnipeg, Canada: Department of Civil Engineering, University of Manitoba. https://mspace.lib.umanitoba.ca/handle/1993/35094.
- Doré, G., Niu, F. and Brooks, H. (2016). Adaptation methods for transportation infrastructure built on degrading permafrost. Permafrost and Periglacial Processes 27(4), 352–364. https:// onlinelibrary.wiley.com/doi/10.1002/ppp.1919.
- Hjort, J., Streletskiy, D., Doré, G., Wu, Q., Bjella, K. and Luoto, M. (2022). Impacts of permafrost degradation on infrastructure. Nature Reviews Earth & Environment 3(1), 24–38. https:// www.nature.com/articles/s43017-021-00247-8.
- Lepage, J.M. and Doré, G. (2016). Experimentation Of Mitigation Techniques to Reduce the Effects of Permafrost Degradation on Transportation Infrastructures at Beaver Creek Experimental Road Site (Alaska Highway, Yukon). Québec, Canada: Department of Civil Engineering, Laval University. https://pubs.aina.ucalgary.ca/cpc/CPC6-526.pdf.
- Shneider von Deimling, T., Lee, H., Ingeman-Nielsen, T.,
 Westermann, S., Romanovsky, V., Lamoureux, S. et al.
 (2021). Consequences of permafrost degradation for Arctic infrastructure Bridging the model gap between regional and

engineering scales. The Cryosphere 15(5), 2451–2471. https://tc.copernicus.org/articles/15/2451/2021/.

- Stern, G. and Gaden, A. (eds.) (2015). The impact of climate change on infrastructure in the western and central Canadian Arctic. In From Science to Policy in the Western and Central Canadian Arctic: An Integrated Regional Impact Study (IRIS) of Climate Change and Modernization. Stern, G. and Gaden, A. (eds.). Quebéc, Canada: ArcticNet Incorporated. Chapter 7. 300–341. https://arcticnet.ulaval.ca/wp-content/ uploads/2022/06/IRIS_FromScience_ArcticNet_Ir_.pdf.
- Streletskiy, D.A., Clemens, S., Lanckman, J.P. and Shiklomanov, N.I. (2023). The costs of Arctic infrastructure damages due to permafrost degradation. Environmental Research Letters 18, 014006. https://iopscience.iop.org/ article/10.1088/1748-9326/acab18/meta

Maps

- Hjort, J., Streletskiy, D., Doré, G., Wu, Q., Bjella, K. and Luoto, M. (2022). Impacts of permafrost degradation on infrastructure. Nature Reviews Earth & Environment 3, 24–38. https://doi. org/10.1038/543017-021-00247-8.
- Streletskiy, D.A., Clemens, S., Lanckman, J.P. and Shiklomanov, N.I. (2023). The costs of Arctic infrastructure damages due to permafrost degradation. Environmental Research Letters 18, 014006. https://iopscience.iop.org/article/10.1088/1748-9326/ acab18/meta

Undermined

- Booshehrian, A., Wan, R. and Su, X. (2020). Hydraulic variations in permafrost due to open-pit mining and climate change: A case study in the Canadian Arctic. Acta Geotechnica 15, 883–905. https://link.springer.com/article/10.1007/511440-019-00786-x.
- Davies, M. (2011). Filtered dry stacked tailings The fundamentals. Tailings and Mine Waste. Vancouver, BC, 6–9 November 2011. Vancouver: University of British Columbia. 1–9. https://open. library.ubc.ca/media/stream/pdf/59368/1.0107683/1.
- Glomsrød, S., Duhaime, G. and Aslaksen, I. (eds.) (2021). The Economy of the North. ECONOR 2020. Oslo: Statistisk sentralbyrå, Statistics Norway. https://www.chaireconditionautochtone. fss.ulaval.ca/doc/Publication/ECONOR-2020-FinalReport-20May2021_LOW.pdf
- International Council on Mining and Metals (2019). Adapting to Climate Change: Building Resilience in the Mining and Metals Industry. London. https://www.icmm.com/website/ publications/pdfs/environmental-stewardship/2019/ guidance_changing-climate.pdf.
- Kolb, J., Rosa, D., Thrane, K., & Keiding, J. K. (2016). Greenland.
 In R. Boyd, T. Bjerkgård, B. Nordahl, & H. Schiellerup (Eds.),
 Mineral resources in the Arctic (pp. 162-189). (Geological

Survey of Norway Special Publication.). Geological Survey of Norway. https://www.ngu.no/upload/Aktuelt/CircumArtic/ Mineral_Resources_Arctic_Shortver_Eng.pdf

- Medeiros, A.S., Wood, P., Wesche, S.D., Bakaic, M. and Peters, J.F. (2017). Water security for northern peoples: Review of threats to Arctic freshwater systems in Nunavut, Canada. Regional Environmental Change 17(3), 635–647. https://link. springer.com/article/10.1007/s10113-016-1084-2.
- Stephani, E., Angelopouloss, M., Musial, M., Anderson, S., Gince, P. and Kissiova, M. (2016). Investigation and management of permafrost and groundwater conditions at mining sites: Case studies in continuous permafrost of Alaska and Canada. Eleventh International Conference on Permafrost: Exploring Permafrost in a Future Earth. Potsdam, Germany, 20–24 June 2016. 1146–1147. https://simmakers.com/wp-content/ uploads/icop2016/icop2016_book_of_abstracts.pdf

Maps

- Boyd, R., Bjerkgård, T., Nordahl, B. and Schiellerup, H. (eds.) (2016). Mineral Resources in the Arctic: An Introduction. Trondheim, Norway: NGU – Geological Survey of Norway. https://www.ngu.no/upload/Aktuelt/CircumArtic/Mineral_ Resources_Arctic_Shortver_Eng.pdf.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi.org/10.1016/j.earscirev.2019.04.023.

Keeping the light on

Text

- Biev, A.A. and Serova, N.A. (2020). Features of the organization of fuel supplies to the Arctic regions of Russia: Assessment of transport conditions. IOP Conference Series: Earth and Environmental Science 539, 012017. https://iopscience.iop. org/article/10.1088/1755-1315/539/1/012017/pdf.
- Cherry, J.E., Knapp, C., Trainor, S., Ray, A.J., Tedesche, M. and Walker, S. (2017). Planning for climate change impacts on hydropower in the Far North. Hydrology and Earth System Sciences 21(1), 133–151. https://hess.copernicus.org/ articles/21/133/2017/hess-21-133-2017.pdf.
- Dilley, L M. and Hulse, L. (2007). Foundation Design of Wind Turbines in Southwestern Alaska, a Case Study. Anchorage, Alaska: Institute of the North. https://helicalpileworld.com/Ren-13%20Foundation%20Design%200f%20Wind%20Turbines.pdf.
- Intergovernmental Panel on Climate Change (2021). Climate Change 2021: The Physical Science Basis: Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Chen,

Y. et al. (eds.). Cambridge, UK and New York, US: Cambridge University Press. https://www.ipcc.ch/report/ar6/wg1/. Lenio. M. (2020). Nunavut residents need support for renewables.

- 22 October. https://wwf.ca/stories/nunavut-residents-needsupport-for-renewables/. Accessed 31 May 2023.
- Natural Resources Canada (2021). Glencore RAGLAN Mine Renewable Electricity Smart-Grid Pilot Demonstration, 9 March. https://www.nrcan.gc.ca/science-and-data/fundingpartnerships/funding-opportunities/current-investments/ glencore-raglan-mine-renewable-electricity-smart-grid-pilotdemonstration/16662. Accessed 31 May 2023.
- Nogovitsyn, D.D., Sheina, Z.M. and Sergeeva, L.P. (2020). Influence of the Vilyui hydropower plant on the regime of the Vilyui River. IOP Conference Series: Earth and Environmental Science 579, 012031. https://iopscience.iop. org/article/10.1088/1755-1315/579/1/012031/pdf.
- Panfilov, A.A. (2018). Features of calculation schemes and methods for design of wind turbine foundations for Arctic conditions. 2018 International Ural Conference on Green Energy (UralCon). Chelyabinsk, Russia, 4–8 October 2018. 122–126. https://ieeexplore.ieee.org/document/8544376.
 Salonen, H. (2021). All habits die hard: Exploring the path dependence and lock-ins of outdated energy systems in the Russian Arctic.
- Energy Research & Social Science 78, 102149. https://www. sciencedirect.com/science/article/pii/S2214629621002425.
- Zheng, M., Yang, Z.J., Yang, S. and Still, B. (2017). Modeling and mitigation of excessive dynamic responses of wind turbines founded in warm permafrost. Engineering Structures 148, 36–46. https://www.sciencedirect.com/science/article/abs/ pii/S0141029617301876.

Maps

- Norwegian Directorate for Cultural Heritage (2022). Kulturminner – Lokaliteter, Enkeltimmen og Sikringssoner. Geonorge. https://kartkatalog.geonorge.no/metadata/kulturminnerlokaliteter-enkeltminner-og-sikringssoner/c72906a0-2bc2-41d7-bea2-c92d368e3c49. Accessed 20 June 2023.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi.org/10.1016/j.earscirev.2019.04.023.

No time to waste

Text

Arctic Council (2021). Tackling waste pollution in the Arctic with community empowerment, 10 May. https://arctic-council.org/news/tackling-waste-pollution-in-the-arctic-with-community-empowerment/. Accessed 31 May 2023.
Burns, C., Orttung, R.W., Shaiman, M., Silinsky, L. and Zhang,

E. (2021). Solid waste management in the Arctic. Waste Management 126, 340–350. https://www.sciencedirect.com/ science/article/abs/pii/S0956053X2100163X.

- Grebenets, V.I., Tolmanov, V.A., lurov, F.D. and Groisman, P.Y. (2021). The problem of storage of solid waste in permafrost. Environmental Research Letters 16(10), 105007. https:// iopscience.iop.org/article/10.1088/1748-9326/ac2375/pdf.
- Hjort, J., Streletskiy, D., Doré, G., Wu, Q., Bjella, K. and Luoto, M. (2022). Impacts of permafrost degradation on infrastructure. Nature Reviews Earth & Environment 3, 24–38. https://www. nature.com/articles/s43017-021-00247-8.
- Magee, G.L. and Rice, W.J. (2002). Rethinking landfill development and operation in permafrost regions. 11th International Conference on Cold Regions Engineering: Cold Regions Impacts on Transportation and Infrastructure. Anchorage, Alaska, 20–22 May 2002. 910–921. Reston, Virginia, United States: American Society of Civil Engineers. https://ascelibrary.org/doi/ abs/10.1061/40621%28254%2979.
- Sustainable Development Working Group (2019). Best Waste Management Practices for Small and Remote Arctic Communities: A Study of Best Waste Management Practices Used in Remote Arctic Communities in Alaska, Canada, and Finland. Tromsø, Norway: Arctic Council. https://oaarchive.arctic-council.org/ bitstream/handle/11374/2297/SAOFI204_2019_RUKA_07-01-06_SDWG_Report-Waste-Management-Best-Practices.pdf. United States of America, Alaska Administrative Code, Title
- 18 Environmental Conservation, Chapter 60 Solid Waste Management, Article 2 – Waste Disposal Permit, Section 18 AAC 60.228 – Exemptions for Freezeback Landfills. https:// www.law.cornell.edu/regulations/alaska/18-AAC-60.228.

Modern history

- Markkula, I., Turunen, M. and Rasmus, S. (2019). A review of climate change impacts on the ecosystem services in the Saami Homeland in Finland. Science of the Total Environment 692, 1070–1085. https://www.sciencedirect.com/science/ article/abs/pii/S0048969719333868.
- Nicu, I.C., Tanyas, H., Rubensdotter, L. and Lombardo, L. (2022). A glimpse into the northernmost thermo-erosion gullies in Svalbard archipelago and their implications for Arctic cultural heritage. Catena 212, 106105. https://www.sciencedirect. com/science/article/pii/S0341816222000911.
- Norway, Svalbard Environmental Protection Act 2001, Act of 15 June 2001, No. 79, Relating to the Protection of the Environment in Svalbard. https://www.regjeringen.no/en/dokumenter/ svalbard-environmental-protection-act/id173945/.
- Sinitsyn, A., Westermann, S., Arlov, T.B., Meyer, A., Lutz, J., Bekele, Y.W. et al. (2022). PCCH-Arctic – Polar climate and cultural heritage – Preservation and restoration management:

Project presentation and current results. Frostdagen 2022 at Norwegian Geotechnical Society. Oslo, 24 November 2022. Oslo: SINTEF Community. https://sintef.brage.unit.no/sintefxmlui/handle/11250/3035461.

Statistics Norway (2016). This is Svalbard 2016: What the Figures Say. Oslo: Statistisk sentralbyrå, Statistics Norway. https:// www.ssb.no/en/befolkning/artikler-og-publikasjoner/_ attachment/294354?_ts=15a12de02c0.

Maps

- Langer, M., Schneider von Deimling, T., Westermann, S., Rolph, R., Rutte, R., Antonova, S. et al. (2023). Thawing permafrost poses environmental threat to thousands of sites with legacy industrial contamination. Nature Communications 14, 1721. https://doi.org/10.1038/s41467-023-37276-4.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi. org/10.1016/j.earscirev.2019.04.023.

Going South

A planetary perspective

Text

- Haeberli, W., Noetzli, J., Arenson, L., Delaloye, R., Gärtner-Roer, I., Gruber, S. et al. (2010). Mountain permafrost: Development and challenges of a young research field. Journal of Glaciology 56(200), 1043–1058. https://doi. org/10.3189/002214311796406121.
- Yoshikawa, K., Schorghofer, N. and Klasner, F. (2023). Permafrost and seasonal frost thermal dynamics over fifty years on tropical Maunakea volcano, Hawai'i. Arctic, Antarctic, and Alpine Research. 55, 1. https://doi.org/10.1080 /15230430.2023.2186485

Maps

- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi. org/10.1016/j.earscirev.2019.04.023.
- Obu, J., Westermann, S., Kääb, A. and Bartsch, A. (2019). Ground Temperature Map, 2000–2016, Andes, New Zealand and East African Plateau Permafrost. University of Oslo, PANGAEA. https://doi.pangaea.de/10.1594/PANGAEA.905512. Accessed 20 June 2023.
- Obu, J., Westermann, S., Vieira, G., Abramov, A., Balks, M.R., Bartsch, A. et al. (2020). Pan-Antarctic map of near-surface permafrost temperatures at 1 km² scale. The Cryosphere

14(2), 497–519. https://doi.org/10.5194/tc-14-497-2020. Overduin, P.P., Schneider von Deimling, T., Miesner, F., Grigoriev, M.N., Ruppel, C., Vasiliev, A. et al. (2019). Submarine permafrost map in the Arctic modeled using 1-D transient heat flux (SuPerMAP). Journal of Geophysical Research: Oceans 124(6), 3490–3507. https://doi.org/10.1029/2018JC014675.

Frozen giants

Text

- Biskaborn, B.K., Smith, S.L., Noetzli, J., Matthes, H., Vieira, G., Streletskiy, D.A. et al. (2019). Permafrost is warming at a global scale. Nature Communications 10(264). https://www. nature.com/articles/s41467-018-08240-4.
- Hock, R., Rasul, G., Adler, C., Cáceres, B., Gruber, S., Hirabayashi, Y. et al. (2019). High mountain areas. In The Ocean and Cryosphere in a Changing Climate: Special Report of the Intergovernmental Panel on Climate Change. Pörtner, H.-O., Masson-Delmotte, V., Roberts, D.C., Tignor, M., Zhai, P., Poloczanska, E. et al. (eds.). Cambridge, UK and New York, US: Cambridge University Press. 39–70. https:// www.ipcc.ch/srocc/.
- Jin, X., Jin, H., Luo, D., Sheng Y., Wu, Q., Wu, J. et al. (2022). Impacts of permafrost degradation on hydrology and vegetation in the source area of the Yellow River on northeastern Qinghai-Tibet Plateau, southwest China. Frontiers in Earth Science 10. https:// doi.org/10.3389/feart.2022.845824.
- Jones, D.B., Harrison, S., Anderson, K. and Betts, R.A. (2018). Mountain rock glaciers contain globally significant water stores. Scientific Reports 8(1), 1–10. https://doi.org/10.1038/ 541598-018-21244-w.
- Schneider von Deimling, T., Lee, H., Ingeman-Nielsen, T., Westermann, S., Romanovsky, V., Lamoureux, S. et al. (2021). Consequences of permafrost degradation for Arctic infrastructure – Bridging the model gap between regional and engineering scales. The Cryosphere 15(5), 2451-2471. https:// doi.org/10.5194/tc-15-2451-2021.

Maps

- NASA Shuttle Radar Topography Mission (2013).Shuttle Radar Topography Mission (SRTM) Global. OpenTopography. https:// portal.opentopography.org/datasetMetadata?otCollectionID= OT.042013.4326.1 Accessed 20 June 2023. https://doi.org/ 10.5069/G9445JDF.
- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi.org/10.1016/j.earscirev.2019.04.023.
- Obu, J., Westermann, S., Kääb, A. and Bartsch, A. (2019). Ground Temperature Map, 2000–2016, Andes, New

Zealand and East African Plateau Permafrost. University of Oslo, PANGAEA. https://doi.pangaea.de/10.1594/ PANGAEA.905512. Accessed 20 June 2023.

Obu, J., Westermann, S., Vieira, G., Abramov, A., Balks, M.R., Bartsch, A. et al. (2020). Pan-Antarctic map of near-surface permafrost temperatures at 1 km² scale. The Cryosphere 14(2), 497–519. https://doi.org/10.5194/tc-14-497-2020.

The view from the top

- Arenson, L.U., Harrington, J.S., Koenig, C.E. and Wainstein, P.A. (2022). Mountain permafrost hydrology—A practical review following studies from the Andes. Geosciences 12(2), 48. https://doi.org/10.3390/geosciences12020048.
- Baldis, C.T. and Liaudat, T.D. (2019). Rockslides and rock avalanches in the Central Andes of Argentina and their possible association with permafrost degradation. Permafrost and Periglacial Processes 30(4), 330–347. https://doi. org/10.1002/ppp.2024.
- Barsch, D. (1996). Rockglaciers and the high mountain environment. In Rockglaciers – Indicators for the Present and Former Geoecology in High Mountain Environments. Springer Series in Physical Environment vol. 16. Berlin: Springer Berlin Heidelberg. Chapter 10. 231–268. https://doi. org/10.1007/978-3-642-80093-1_11.
- Bellisario, A., Ferrando, F. and Janke, J. (2013). Water resources in Chile: The critical relation between glaciers and mining for sustainable water management. Investigaciones Geográficas 46(2013), 3–24. https://doi.org/10.5354/0719-5370.2013.30288.
- Boeckli, L., Brenning, A., Gruber, S. and Noetzli, J. (2012). Permafrost distribution in the European Alps: Calculation and evaluation of an index map and summary statistics. The Cryosphere 6(4), 807-820. https://doi.org/10.5194/ tc-6-807-2012.
- Brain, K.A. (2017). The impacts of mining on the livelihoods in the Andes: A critical overview. The Extractive Industries and Society 4(2), 410–418. https://doi.org/10.1016/j. exis.2017.03.001.
- Gruber, S. and Haeberli, W. (2009). Mountain permafrost. In Permafrost Soils. Margesin, R. (ed.). Soil Biology 16, 33–44. https://doi.org/10.1007/978-3-540-69371-0_3.
- Gruber, S., Fleiner, R., Guegan E. et al. (2017). Review article: Inferring permafrost and permafrost thaw in the mountains of the Hindu Kush Himalaya region. The Cryosphere, 11, pp. 81-99. https://doi.org/10.5194/tc-11-81-2017
- Hjort, J., Streletskiy, D., Doré, G., Wu, Q., Bjella, K. and Luoto M. (2022). Impacts of permafrost degradation on infrastructure. Nature Reviews Earth & Environment 3(1), 24–38. https://doi. org/10.1038/s43017-021-00247-8.

- Jin, X., Jin, H., Luo, D., Sheng Y., Wu, Q., Wu, J. et al. (2022). Impacts of permafrost degradation on hydrology and vegetation in the source area of the Yellow River on northeastern Qinghai-Tibet Plateau, southwest China. Frontiers in Earth Science 10. https://doi.org/10.3389/feart.2022.845824.
- Jones, D.B., Harrison, S., Anderson, K. and Betts, R.A. (2018). Mountain rock glaciers contain globally significant water stores. Scientific Reports 8(1), 1–10. https://doi.org/10.1038/ 541598-018-21244-w.
- Kääb, A., Kaufmann, V., Ladstädter, R. and Eiken, T. (2003).
 Rock glacier dynamics: Implications from high-resolution measurements of surface velocity fields. Eighth International Conference on Permafrost. Phillips, M., Springman, S.M. and Arenson, L.U. (eds.). Zürich, 21–25 July 2003. Lisse, Netherlands: Swets & Zeitlinger. 501–506. https://www.staff. tugraz.at/viktor.kaufmann/Chapter_089.pdf.
- Lu, Q., Zhao, D. and Wu, S. (2017). Simulated responses of permafrost distribution to climate change on the Qinghai– Tibet Plateau. Scientific Reports 7(3845). https://doi. org/10.1038/s41598-017-04140-7.
- Peng, X., Zhang, T., Frauenfeld, O.W., Wang, K., Luo, D., Cao, B. et al. (2018). Spatiotemporal changes in active layer thickness under contemporary and projected climate in the northern hemisphere. Journal of Climate 31(1), 251–266. https://doi.org/10.1175/JCLI-D-16-0721.1.
- Rangecroft, S., Suggitt, A.J., Anderson, K. and Harrison, S. (2016). Future climate warming and changes to mountain permafrost in the Bolivian Andes. Climatic Change 137(1–2), 231–243. https://doi.org/10.1007/S10584-016-1655-8.
- Ruiz, L. and Trombotto, D. (2012). Mountain permafrost distribution in the Andes of Chubut (Argentina) based on a statistical model. Proceedings of the Tenth International Conference on Permafrost. Hinkel, K.M. (ed.). Salekhard, Russian Federation, 25–29 June 2012. Salekhard: The Northern Publisher. 365–370. https://www.permafrost.org/conference-proceedings/.
- Streletskiy, D. (2021). Permafrost degradation. In Snow and ice-related hazards, risks, and disasters. Haeberli, W. and Whiteman, C. (eds.). Hazards and Disasters Series, 297–322. https://doi.org/10.1016/B978-0-12-817129-5.00021-4.
- Wei, D., Qi, Y., Ma, Y. and Wang, X. (2021). Plant uptake of CO2 outpaces losses from permafrost and plant respiration on the Tibetan Plateau. PNAS 118(33), e2015283118. https://doi. org/10.1073/pnas.2015283118.
- Zhang G., Nan, Z., Hu, N., Yin, Z., Zhao, L., Cheng, G. et al. (2022). Qinghai-Tibet Plateau permafrost at risk in the late 21st century. Earth's Future 10(6), e2022EF002652. https:// doi.org/10.1029/2022EF002652.
- Zou, D., Zhai, L., Sheng, Y., Chen, J., Hu, G., Wu, T. et al. (2017). A new map of permafrost distribution on the Tibetan Plateau. The Cryosphere 11(6), 2527–2542. https://doi.org/10.5194/ tc-11-2527-2017.

Maps

- Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H.H., Dashtseren, A. et al. (2019). Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale. Earth-Science Reviews 193, 299–316. https://doi.org/10.1016/j.earscirev.2019.04.023. Obu, J., Westermann, S., Kääb, A. and Bartsch, A. (2019). Ground
- Temperature Map, 2000–2016, Andes, New Zealand and East African Plateau Permafrost. University of Oslo, PANGAEA. https://doi.pangaea.de/10.1594/PANGAEA.905512. Accessed
- 20 June 2023.

Europe's frozen heart

Text

- Allen, S. and Huggel, C. (2013). Extremely warm temperatures as a potential cause of recent high mountain rockfall. Global and Planetary Change 107, 59–69. https://doi.org/10.1016/j. gloplacha.2013.04.007.
- Boeckli, L., Brenning, A., Gruber, S. and Noetzli, J. (2012). Permafrost distribution in the European Alps: Calculation and evaluation of an index map and summary statistics. The Cryosphere 6(4), 807-820. https://doi.org/10.5194/ tc-6-807-2012.
- Guerin, A., Ravanel, L., Matasci, B., Jaboyedoff, M. And Deline, P. (2020). The three-stage rock failure dynamics of the Drus (Mont Blanc massif, France) since the June 2005 large event. Scientific Reports 10, 17330. https://doi.org/10.1038/541598-020-74162-1.
- Haeberli, W. and Gruber, S. (2009). Global warming and mountain permafrost. In Permafrost Soils. Margesin, R. (ed.). Soil Biology 16, 33–44. https://doi.org/10.1007/978-3-540-69371-0_14.
- Haeberli, W., Barsch, D., Brown, J., Guodong, C., Corte, A.E., Dramis, F. et al. (1998). Mapping, modelling and monitoring of mountain permafrost: a review of ongoing programs. Mountain permafrost working group report. Proceedings of the Seventh International Conference on Permafrost. Lewkowicz, A.G. and Allard, M. (eds.). Yellowknife, Canada, 23–27 June 1998. Ottawa and Québec: University of Ottawa and Université Laval. 233–246. https://www.uspermafrost. org/conference-proceedings.
- Hock, R., Rasul, G., Adler, C., Cáceres, B., Gruber, S.,
 Hirabayashi, Y. et al. (2019). High mountain areas. In The Ocean and Cryosphere in a Changing Climate: Special Report of the Intergovernmental Panel on Climate Change.
 Pörtner, H.-O., Masson-Delmotte, V., Roberts, D.C., Tignor, M., Zhai, P., Poloczanska, E. et al. (eds.). Cambridge, UK and New York, US: Cambridge University Press. 39–70. https://www.ipcc.ch/srocc/.
- Mourey, J., Perrin-Malterre, C. and Ravanel, L. (2020). Strategies used by French Alpine guides to adapt to the effects of

climate change. Journal of Outdoor Recreation and Tourism 29, 100278. https://doi.org/10.1016/j.jort.2020.100278. Weber, S., Beutel, J., Forni, R.D., Geiger, A., Gruber, S., Hasler, A. et al. (2019). A decade of detailed observations (2008–2018) in steep bedrock permafrost at the Matterhorn Hörnligrat (Zermatt, CH). Earth System Science Data 11(3), 1203–1237. https://doi.org/10.5194/essd-11-1203-2019.

Maps

- Guerin, A., Ravanel, L., Matasci, B., Jaboyedoff, M. and Deline, P. (2020). The three-stage rock failure dynamics of the Drus (Mont Blanc massif, France) since the June 2005 large event. Scientific Reports 10, 17330. https://doi.org/10.1038/s41598-020-74162-1.
- Obu, J., Westermann, S., Kääb, A. and Bartsch, A. (2019). Ground Temperature Map, 2000–2016, Andes, New Zealand and East African Plateau Permafrost. University of Oslo, PANGAEA.
- https://doi.pangaea.de/10.1594/PANGAEA.905512. Accessed 20 June 2023.

The ends of the Earth I

- Burton-Johnson, A., Black, M., Fretwell, P.T. and Kaluza-Gilbert, J. (2016). An automated methodology for differentiating rock from snow, clouds and sea in Antarctica from Landsat 8 imagery: A new rock outcrop map and area estimation for the entire Antarctic continent. The Cryosphere 10, 1665–1677. https://doi.org/10.5194/tc-10-1665-2016.
- Convey, P., Chown, S.L., Clarke, A., Barnes, D.K.A., Cummings, V., Ducklow, H. et al. (2014). The spatial structure of Antarctic biodiversity. Ecological Monographs 84(2), 203–244. https:// doi.org/10.1890/12-2216.1.
- Gray, A., Krolikowski, M., Fretwell, P., Convey, P., Peck, L.S., Mendelova, M. et al. (2020). Remote sensing reveals Antarctic green snow algae as important terrestrial carbon sink. Nature Communications 11(2527). https://doi.org/10.1038/541467-020-16018-w.
- Guglielmin, M., Fratte, M.D. and Cannone, N. (2014). Permafrost warming and vegetation changes in continental Antarctica. Environmental Research Letter 9(4), 045001. https://doi. org/10.1088/1748-9326/9/4/045001.
- Medley, B., McConnell, J.R., Neumann, T.A., Reijmer, C.H., Chellman, N., Sigl, M. et al. (2018). Temperature and snowfall in western Queen Maud Land increasing faster than climate model projections. Geophysical Research Letters 45(3), 1472–1480. https://doi. org/10.1002/2017GL075992.
- Obu, J., Westermann, S., Vieira, G., Abramov, A., Balks, M.R., Bartsch, A. et al. (2020). Pan-Antarctic map of near-surface permafrost temperatures at 1 km2 scale. The Cryosphere

14(2), 497-519. https://doi.org/10.5194/tc-14-497-2020.

Philippe, M., Tison, J.-L., Fjøsne, K., Hubbard, B., Kjær, H.A., Lenaerts, J.T.M. et al. (2016). Ice core evidence for a 20th century increase in surface mass balance in coastal Dronning Maud Land, East Antarctica. The Cryosphere 10(5), 2501– 2516. https://doi.org/10.5194/tc-10-2501-2016.

Potapowicz, J., Szumińska, D., Szopińska, M. and Polkowska, Ż. (2019). The influence of global climate change on the environmental fate of anthropogenic pollution released from the permafrost: Part I. Case study of Antarctica. Science of the Total Environment 651, 1534–1548. https://doi.org/10.1016/j. scitotenv.2018.09.168.

Turner, J., Lu, H., King, J., Marshall, G.J., Phillips, T., Bannister, D. et al. (2021). Extreme temperatures in the Antarctic. Journal of Climate 34(7), 2653–2668. https://doi.org/10.1175/ JCLI-D-20-0538.1.

Vieira, G., Bockheim, J., Guglielmin, M., Balks, M., Abramov, A.A., Boelhouwers, J. et al. (2010). Thermal state of permafrost and active-layer monitoring in the Antarctic: Advances during the international polar year 2007–2009. Permafrost and Periglacial Process 21(1), 182–197. https:// doi.org/10.1002/ppp.685.

Maps

- Matsuoka, K., Skoglund, A., Roth, G., de Pomereu, J., Griffiths, H., Headland, R. et al. (2021). Quantarctica, an integrated mapping environment for Antarctica, the Southern Ocean, and sub-Antarctic islands. Environmental Modelling & Software 140, 105015. https://doi.org/10.1016/j. envsoft.2021.105015.
- Obu, J., Westermann, S., Vieira, G., Abramov, A., Balks, M.R., Bartsch, A. et al. (2020). Pan-Antarctic map of near-surface permafrost temperatures at 1 km² scale. The Cryosphere 14(2), 497–519. https://doi.org/10.5194/tc-14-497-2020.

The ends of the Earth II

Text

Melting History (2020). The waiting game II : A climate change issue? Melting History. https://meltinghistory.org/blog/ the-waiting-game-ii-a-climate-change-issue. [Accessed 15 March 2022].

Obu, J., Westermann, S., Vieira, G., Abramov, A., Balks, M.R., Bartsch, A. et al. (2020). Pan-Antarctic map of near-surface permafrost temperatures at 1km2 scale, The Cryosphere 14(2), 497–519. https://doi.org/10.5194/tc-14-497-2020.

Ramos, M., Vieira, G., de Pablo, M.A., Molina, A. and Jimenez, J.J. (2020). Transition from a subaerial to a subnival permafrost temperature regime following increased snow cover (Livingston Island, maritime Antarctic). Atmosphere 11(12), 1332. https://doi.org/10.3390/atmos11121332. Turner, J., Lu, H., King, J., Marshall, G.J., Phillips, T., Bannister, D. et al. (2021). Extreme temperatures in the Antarctic. Journal of Climate 34(7), 2653–2668. https://doi.org/10.1175/ JCLI-D-20-0538.1.

Maps

- Mapping Application for Penguin Populations and Projected Dynamics (2021). MAPPPD v4.1. Oceanites Inc. https://www. penguinmap.com/mapppd/. Accessed 20 June 2023.
- Matsuoka, K., Skoglund, A., Roth, G., de Pomereu, J., Griffiths, H., Headland, R. et al. (2021). Quantarctica, an integrated mapping environment for Antarctica, the Southern Ocean, and sub-Antarctic islands. Environmental Modelling & Software 140, 105015. https://doi.org/10.1016/j. envsoft.2021.105015.
- Obu, J., Westermann, S., Vieira, G., Abramov, A., Balks, M.R., Bartsch, A. et al. (2020). Pan-Antarctic map of near-surface permafrost temperatures at 1 km² scale. The Cryosphere 14(2), 497–519. https://doi.org/10.5194/tc-14-497-2020.

The ends of the Earth III

Text

- Goordial, J., Davila, A., Lacelle, D., Pollard, W., Marinova, M.M., Greer, C.W. et al. (2016). Nearing the cold-arid limits of microbial life in permafrost of an upper dry valley, Antarctica. The ISME Journal 10(7), 1613–1624. https://doi.org/10.1038/ ismej.2015.239.
- Heldmann J.L., Pollard, W., McKay, C.P., Marinova, M. M., Davila, A., Williams, K. E. et al. (2013). The high elevation Dry Valleys in Antarctica as analog sites for subsurface ice on Mars. Planetary and Space Science 85, 53-58. https://doi. org/10.1016/j.pss.2013.05.019.
- Marchant D.R. and Head III, J.W. (2007). Antarctic dry valleys: Microclimate zonation, variable geomorphic processes, and implications for assessing climate change on Mars. Icarus 192(1), 187–222. https://doi.org/10.1016/j. icarus.2007.06.018.
- Medley, B., McConnell, J.R., Neumann, T.A., Reijmer, C.H., Chellman, N., Sigl, M. et al. (2018). Temperature and snowfall in western Queen Maud Land increasing faster than climate model projections. Geophysical Research Letters 45(3), 1472–1480. https://doi.org/10.1002/2017GL075992.
- Philippe, M., Tison, J.-L., Fjøsne, K., Hubbard, B., Kjær, H.A., Lenaerts, J.T.M. et al. (2016). Ice core evidence for a 20th century increase in surface mass balance in coastal Dronning Maud Land, East Antarctica, The Cryosphere 10(5), 2501– 2516. https://doi.org/10.5194/tc-10-2501-2016.
- Verret, M., Dickinson, W., Lacelle, D., Fisher, D., Norton, K., Chorley, H. et al. (2021). Cryostratigraphy of mid-Miocene

permafrost at Friis Hills, McMurdo Dry Valleys of Antarctica. Antarctic Science 33(2), 174–188. https://doi.org/10.1017/ S0954102020000619.

Maps

- Matsuoka, K., Skoglund, A., Roth, G., de Pomereu, J., Griffiths, H., Headland, R. et al. (2021). Quantarctica, an integrated mapping environment for Antarctica, the Southern Ocean, and sub-Antarctic islands. Environmental Modelling & Software 140, 105015. https://doi.org/10.1016/j. envsoft.2021.105015.
- Obu, J., Westermann, S., Vieira, G., Abramov, A., Balks, M.R., Bartsch, A. et al. (2020). Pan-Antarctic map of near-surface permafrost temperatures at 1 km² scale. The Cryosphere 14(2), 497–519. https://doi.org/10.5194/tc-14-497-2020.

Glossary and Acronyms

van Everdingen, R.O. (2005). Multi-language Glossary of Permafrost and Related Ground-Ice Terms. Calgary, Canada: Arctic Institute of North America and University of Calgary. https://globalcryospherewatch.org/reference/ glossary_docs/Glossary_of_Permafrost_and_Ground-Ice_ IPA_2005.pdf.

Authors and acknowledgements

NUNATARYUK Advisory Board

Donald Forbes, Emeritus Research Scientist at the Geological Survey of Canada and the Bedford Institute of Oceanography Johnnie (JD) Storr, Director, Hunters and Trappers Committee, Aklavik, Canada Margareta Johansson, Coordinator of the EU Project INTERACT

Nicole Biebow, Coordinator of the EU Polar-Net 2 project

Cartography and Design Advisory Board

Alexander M. Tait, National Geographic Society Margaret Pearce, Independent cartographer Oliver Uberti Creative Adolfo Arranz, South China Morning Post Lauren C. Tierney, The Washington Post

Editorial team

Levi Westerveld, GRID-Arendal Tiina Kurvits, GRID-Arendal Tina Schoolmeester, GRID-Arendal Oda Bortheim Mulelid, GRID-Arendal Torjus Solheim Eckhoff, GRID-Arendal Pier Paul Overduin, AWI Michael Fritz, AWI Hugues Lantuit, AWI

Cartography

Levi Westerveld, GRID-Arendal Alexander M. Tait, National Geographic Society (page 23 and 75)

Illustrations Olga Borjon-Privé (Oluko)

Photography Katie Orlinsky

Layout GRID-Arendal

Copy-editing

Strategic Agenda Guendalina De Luigi, GRID-Arendal

Cover page

Permafrost thaw ponds at BGR, Sheldrake River catchment, Umiujaq (Nunavik, Quebec, Canada). Drone survey and orthomosaic modified from G. Vieira, C. Mora and P. Freitas, Project THAWPOND, PROPOLAR, CEG/IGOT, University of Lisbon and CEN, Laval University.

Experts and contributors (in alphabetical order)

Portraits

Interviews carried out by Torjus Solheim Eckhoff, Susanna Gartler, Thomas Ingeman-Nielsen and Levi Westerveld.

Interviews edited by Guendalina De Luigi, Torjus Solheim Eckhoff, Tiina Kurvits, Tina Schoolmeester and Levi Westerveld.

Interviewees

- Gwen Healey Akearok, Executive Director, Qaujigiartiit Health Research Centre, Iqaluit, Nunavut Canada Angus Alunik, Inuvialuit Settlement Region,
- Community-Based Monitoring Program, Inuvik, NWT
- Annett Bartsch, CEO and senior scientist at b.goes GmbH, Austria Susanna Gartler, PhD student, University
- of Vienna, Austria
- Palle Jerimiassen, Mayor of Avannaata Municipality, Kalaallit Nunaat (Greenland)
- Jessi Pascal, Environmental Coordinator at Gwich'in Renewable Resources Board,
- Aklavik, NWT Canada Ingrid Rekkavik, Advisor at Kings Bay AS,
- NyÅlesund, Norway Vyacheslav Shadrin, Chair of Council of
- Yukaghir Elders, Nelemnoye, Russia Dmitry Streletstkiy, Associate Professor of Geography and International Affairs at the George Washington University, United States

Earth's Freezer: Introduction to Permafrost

Lead author: Tiina Kurvits, GRID-Arendal Michael Fritz Leneisja Jungsberg Amelie Lindgren Frederieke Miesner Pier Paul Overduin Philip Pika Justine Ramage Gonçalo Vieira

Awakening Giant: Permafrost and Climate Change

Lead author: Tiina Kurvits, GRID-Arendal Sandra Arndt Victor Brovkin Sarah Chadburn Michael Fritz Gustaf Hugelius Thomas Kleinen Constance Lefebvre Susanne Liebner Pier Paul Overduin Emilia Ridolfi Maria Scheel Jorien Vonk Dirk Wagner

Moving Grounds: Permafrost Changes

Lead author: Tiina Kurvits, GRID-Arendal Michael Fritz Örjan Gustafsson Bennet Juhls Frederieke Meisner David Nielsen Pier Paul Overduin Tina Schoolmeester Niek Speetjens Gonçalo Vieira

Jorien Vonk

Arctic Ripples: Impacts of Permafrost Thaw

Lead author: Björn Alfthan, GRID-Arendal Khaled Abass Susan A. Crate

Otto Habeck Ian Hiort Thomas Ingeman-Nielsen Lenesja Jungsberg Olli Karjalainen Franz Krause Joan Nymand Larsen Justine Ramage Arja Rautio Rebecca Rolph Peter Schweitzer Elisa Stella **Dmitry Streletskiv Rodrigue Tanguy** Ulla Timlin Mathias Ulrich Anna Vasilevskava Goncalo Vieira Sebastian Westermann

Holding Tight: Adaptation to Permafrost Thaw

Lead author: Torjus Solheim Eckhoff, GRID-Arendal Lukas Arenson Scott Dallimore Guy Dore Jan Hjort Arne Instanes Alexandra Meyer Peter Schweitzer

Going South: Permafrost in Other Areas

Lead author: Anna Sinisalo, GRID-Arendal Elaine Baker Michael Fritz Stephan Gruber Peter Harris Florence Magnin Gonçalo Vieira Ludovic Ravanel

Over the Horizon

Lead author: Pier Paul Overduin, AWI Tiina Kurvits Tina Schoolmeester



The Arctic Permafrost Atlas was written and edited by GRID-Arendal together with all NUNATARYUK project partners, scientists, and earlycareer scientists. It presents state-of-the-art knowledge about permafrost and the impacts of permafrost thaw on human settlements in the Arctic.

The Arctic Permafrost Atlas is a key deliverable of the NUNATARYUK project. NUNATARYUK is an international permafrost research project consisting of 26 partners from 14 countries working together to understand how thawing permafrost on land, along the coast and below the sea changes the global climate and life for people in the Arctic.

The editors would like to acknowledge the valuable contributions of the following people: Arne Aalberg, Marcel Babin, Daniel Baptista, Annett Bartsch, Clément Bertin, Kerstin Brembach, Lisa Bröder, João Canário, Nitin Chaudhary, Bernardo da Costa, Natalia Doloisio, Jade Falardeau, Frank Fell, Vincent Le Fouest, Pedro Freitas, Mathias Göckede, Rafael Gonçalves-Araujo, Gustav Grimstad, Timothy E. Heleniak, Knut Vilhelm Høyland, Jón Haukur Ingimundarson, Konstantin Klein, Rachele Lodi, Victoria Martin, Carla Mora, Georg Pointner, Olga Povoroznyuk, Sandra Raab, Andreas Richter, Torsten Sachs, Niklaas Schmidt, George Tanski, Jean-Paul Vanderlinden, Julia Wagner, Henrike Walther, Birgit Wild, Stiig Wilkenskeld, Dustin Whalen and Alberto Zocchi.



This project is funded by the European Union's Horizon 2020 research and innovation programme under grant agreement No. 773421.



In May 2022, three members of the Advisory Board for Cartography and Design gathered at GRID-Arendal's offices to work intensively together on all the maps and graphics and to discuss matters of layout and design for a week. Their insights and assistance have been of invaluable importance for the quality of this publication.

We would like to thank both IASC and the IPA for providing financial support to make this possible.



External review

Christopher Burn, President, International Permafrost Association Hanne Hvidtfeldt Christiansen, UNIS, UArctic Thematic Network Donald L. Forbes, Emeritus Research Scientist at the Geological Survey of Canada and the Bedford Institute of Oceanography Peter Harris, Managing Director, GRID-Arendal Andrea Hinwood, Chief Scientist, UNEP Emily Jenkins, Professor, University of Saskatchewan (pages 104–105) Lorna Little, GRID-Arendal Miles Macmillan-Lawler, Chief Scientist, GRID-Arendal Merritt Turetsky, Professor, University of Colorado Pages 10-11, 18-19, 32-33, 46-51, 54-55, 60-61, 64–65, 98–99 and 102–105 anonymously reviewed through the AMAP Secretariat.

NUNATARYUK Partners

ARTIST SPOTLIGHT Olga Borjon-Privé (Oluko)


I was born and grew up on the Taimyr Peninsula, located in the Arctic part of Siberia. My mother was a teacher by profession and an artist at heart, and my father loved to draw. Thanks to him, I picked up a pencil and began to draw characters that I had invented or read about in books.

I studied at the Norilsk College of Arts and the State University of Saint Petersburg, where I received an artist diploma and a teacher's diploma of decorative and applied arts. I worked for three years at the Taimyr House of Popular Creativity and for 10 years as a teacher at the Norilsk College of Arts.


I use different materials and techniques to create my works. When I paint, I use acrylic. When creating graphic works, I use ink, watercolour, and soft materials like pastels or charcoal. For the illustrations for the Arctic Permafrost Atlas I used a graphics tablet, which allowed me to achieve a variety of effects thanks to a large selection of brushes. I am inspired by traditional folk art, folklore, history, and the people I meet. I'm also inspired by nature: landscapes, textures, and the animal world.

From the very beginning of this project, I worked with great enthusiasm to create illustrations for it. Although the people portrayed in the Atlas come from different parts of the Arctic, I understood their problems and feelings perfectly. The theme of permafrost is common to us – the people who live or are born there. Permafrost is an integral part of my homeland. To my people, the ground is like deerskin, the trees and plants like hairs, and the people like insects. As I learned the stories of others who are experiencing environmental change, I became more deeply inspired by it and understood how important it is not to be silent about it. I wanted my illustrations to talk not only about the problems, but also tell readers about the place or culture of the people living in these regions. By studying the places, people, and the traditions mentioned in the texts, each illustration became an independent plot.


Facebook: Olga Borjon-Privé (Porotova) Instagram: olukodolgan



Scientists use maps to assess the consequences of both permafrost thaw and human activities. The threat to the environment is also hidden in landfills that arose from previous human industries: for instance, containers that contained fuels and lubricants are beginning to collapse. I have depicted the tundra in stains against the background of the rusty surface of an oil barrel. Numerous lakes are represented as human organs, and nature as a living organism.



This illustration is enclosed in a circle, in the form of a traditional Yakut calendar, which I took as the basis for creating the plot. The elements of the composition are located around the centre, the red colour symbolising the sun. Climate change entails changes in the traditional way of life of people: the number of fishers and hunters is decreasing, and more and more people are collecting mammoth tusks. In northern and eastern Siberian cultures, the mammoth has long been associated with evil or strong spirits who warn of danger.



This illustration was inspired by a traditional Inuit parka. Its smooth lines and shapes suggested an outline in which I could depict the main elements of the composition. I have imagined a map on which I have placed mountains, rivers, fishing boats, roads, as well as silhouettes of a man with a dog and birds. This is part of the life of the inhabitants of Aklavik – their activities, their place. With the onslaught of warming, permafrost is retreating, exposing its pressure points, primarily through the state of infrastructure and then affecting the lifestyle of its inhabitants.




The thawing of permafrost due to global warming is a serious change that worries people living in the Arctic, as it affects their lifestyle and puts it at risk. This thawing occurs gradually, and since permafrost is underground, it is not very noticeable. However, the thawing of glaciers directly shows the process of climate change on Earth. In the illustration, I have tried to show the beauty of nature against the background of glaciers, an integral part of the Arctic landscape.



The accident next to Norilsk has served as the main plot for the creation of this illustration. I have drawn a tundra landscape in spring, in which I have imagined a river in the form of a bending deer. This river is depicted with an oil product that got into it because of an accident, and therefore the deer-river meanders as it suffers from torment. I have worked for many years in Norilsk. With sadness in my heart, I understand that this tragedy will have serious consequences for the entire ecosystem of my native land and for the dietary health of my people.



This illustration depicts the situation when, due to global warming, water occupies more and more space and pushes people out of their homes. At the centre of the composition is a man contemplating the house he has lost. The reflection of the sky on the water in the silhouette of the house corresponds to the man reflecting on his memories of the place he once lived.



"It surprised me how diverse permafrost is. It is sort of a living, breathing, moving creature." Suzanne compares permafrost with a living, breathing, moving creature. This became the main plot of my illustration. The philosophy of the local peoples, the main message of which is to live in harmony with the surrounding world, shows us that permafrost is the basis of life in the Arctic. I would like to show this idea of unity of humanity with nature and how much we depend on it.



The elements of this illustration are arranged so that together they depict an inuksuk. An inuksuk is a traditional Inuit stone monument, used as a landmark for travellers and installed in sacred places and burial sites. Permafrost is represented here in the form of a block of ice: a fragile basis for the life of local residents. The moon and stars against the background of the polar sky silently observe the changes that have already begun.



Ingrid's statement about the buildings in Ny-Ålesund inspired me to create this illustration. They open a window to the past and tell us about people's hopes and perseverance. Therefore, my main element is a pattern on a frozen window in the form of a flock of birds as a symbol of undying hope. I depicted coal houses in the background to convey that Ny-Ålesund was a mining settlement and, despite the fact that coal mining stopped, the houses there are kept in memory of this time.

ARTIST SPOTLIGHT Katie Orlinsky

Photographer Katie Orlinsky has spent the last 15 years covering news stories and feature assignments around the world for publications like National Geographic, The New York Times, and The New Yorker. For the last eight years, the majority of her work has explored how the climate crisis is challenging and transforming the relationship between people, animals, and the land. Katie's photography has earned awards from World Press Photo, Pictures of the Year International, Alexia Foundation, Visa Pour L'image, PDN, and the Art Director's Club. She received a BA in Political Science from Colorado College and an MS in Journalism from Columbia University. Katie teaches as a visiting professor at New York University and the University of Alaska Fairbanks.

www.katieorlinskyphoto.com @Katie Orlinsky



After a successful hunt, a young Inupiat man takes a break from moving whale meat into his family's ice cellar in Utqiagvik, Alaska. Ice cellars are large underground freezers, often generations old, dug deep into the permafrost. However, as permafrost thaws, the cellars may become unusable, either physically damaged or warming to the point where it is no longer safe to store food.



Thawing permafrost beneath the earth's surface releases methane gas into Arctic lakes, causing gas bubbles to form in the frozen water. If the gas is released, a small flame can create a brief fire on the lake's surface, as demonstrated by Melanie Engram and Allen Bondo on a pond near Smith Lake in Fairbanks, Alaska.



The Batagaika crater near the Siberian town of Batagay in Russia has been called the "hell crater" or the "gateway to the underworld". Over 90 metres deep and one kilometre long, this thermokarst depression is one of the largest in the world, holding over 200,000 years of history and climate information. It started forming in the 1960s when the permafrost under the area began to thaw after nearby forests were cleared.



The village of Newtok, Alaska, is sinking as the permafrost beneath it thaws. On a summer bird hunt, four Yupik boys cross a flooded walkway. Newtok is the first community in Alaska to start relocating as a result of climate change, pioneering a process that many other Alaskan villages may soon undergo.



The permafrost tunnel near Fairbanks, Alaska, was built by the United States Army Corps of Engineers in the 1960s and is now used to research and study permafrost. The 110-metre-long tunnel walls contain soil, fossils, frozen animal and plant remains, silt, sand, gravel, bedrock, and thousands of years of history.



Animals graze in the Uvs region of Mongolia. In the winter of 2017–2018, extreme weather conditions in Uvs led to a natural disaster known locally as a dzud. Frigid winds from Siberia caused the temperature to drop to below -50 °C. More than 1 million animals froze or starved to death because they were unable to graze. As summers become hotter and drier, less hay is available for livestock to gain the weight they need to survive the harsh winters.



The Alatna River, Alaska. The treeline is the northernmost limit of where trees can grow. As a result of climate change, the treeline is moving north, transforming the tundra landscape and bringing new wildlife species, such as beavers, moose, and snowshoe hares into the Arctic. Scientists are only just beginning to understand what this means for the future, in particular what impact these landscape changes could have on accelerating permafrost thaw.

